

PROPOSED LABEL
FRONT PANEL

TOPSIN® M 4F

Thiophanate-methyl Fungicide

4 Pound per Gallon Flowable

ACTIVE INGREDIENT:

Thiophanate-methyl(dimethyl[1,2-phenylene]
bis(iminocarbonothioyl)]bis(carbamate))* - - - - - 45%

INERT INGREDIENTS: - - - - - 55%

TOTAL: - - - - - 100%

* Also known as dimethyl 4,4'-o-phenylenebis[3-thioallophanate]

© TOPSIN is a registered trademark of Nippon Soda Company, Ltd., and is licensed to Pennwalt Corporation. Covered by one or more of the following U.S. patents: 3,769,308; 3,856,847, 4,020,095; and 4,029,813.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

May be Harmful if Swallowed, Inhaled, or Absorbed Through Skin


Avoid contact with eyes, skin, or clothing. Avoid breathing of spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not contaminate feed or food. Keep children and pets out of treated areas until material is completely dry.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, induce vomiting with a mixture of soap or mustard in warm water. Call a physician immediately. If on skin, wash immediately with soap and water. If in eyes, flush immediately with plenty of water.

ENVIRONMENTAL HAZARD

Do not apply directly to water and wetlands. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.



BEST DOCUMENT AVAILABLE

4581-25 ✓
FRONT PANEL (cont.)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL - Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.

CONTAINER DISPOSAL - Triple rinse container, then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by State and local authorities.

DIRECTIONS FOR USE - See accompanying brochure located on the rear of this package.

EPA Registration No. 4581-_____

EPA Establishment No. 4581-WA-1

Net Contents 2.5 U.S. Gallons

Sold by:

AGCHEM DIVISION - PENNALT CORPORATION
Philadelphia, Pennsylvania

Made in U.S.A.

Printed in U.S.A.

4581-35 ✓

TOPSIN® M 4F

EPA Registration No. 4581-_____

GENERAL INFORMATION

Apply TOPSIN M with ground or aerial equipment, using sufficient volume of spray to provide thorough coverage. Add required amount of TOPSIN M to partially filled tank agitated by mechanical or hydraulic means and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules.

IMPORTANT: If, after using TOPSIN M as recommended, treatment is not effective, a tolerant strain of fungi may be present (consult your local Pennwalt representative). Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other suitable fungicide.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALMONDS - BROWN ROT BLOSSOM BLIGHT (*Monilinia* sp.) - Apply 34 to 45 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage at early pink bud and early petal fall. For aerial or concentrate sprays, use an equivalent amount per acre in an amount of water appropriate for the application equipment.

APPLES - SCAB (*Venturia* sp.), POWDERY MILDEW (*Podosphaera* sp.), SOOTY BLOTCH (*Gloeodes* sp.), FLY SPECK (*Microthyriella* sp.), BITTER ROT (*Gloeosporium* sp.), and BLACK ROT (*Physalospora* sp.) - Apply 23 to 34 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage at 5 to 10 day intervals from green tip through petal fall; continue at 7 to 14 day intervals in cover sprays.

NOTE: For control of APPLE diseases in the Southeastern states of Florida, Georgia, South Carolina, North Carolina, and Alabama, use 67 to 90 fluid ounces of TOPSIN M per acre.

BEANS - WHITE ROT (*Sclerotinia* sp.) and GRAY MOLD (*Botrytis* sp.) - Apply 34 to 45 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage once at 50% to 70% bloom. Or, apply 23 to 34 fluid ounces per acre twice with the first application when 10% to 30% of the plants have an open blossom and a second application 4 to 7 days later or at peak bloom. Do not apply to snap or dry beans within 14 days of harvest, or to lima beans within 28 days of harvest.

CELERY - EARLY BLIGHT (*Cercospora* sp.) and LATE BLIGHT (*Septoria* sp.) - Apply 11 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage at the base of the plants on a 14 to 21 day schedule beginning about 2 weeks after transplanting. Or, apply 11 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage as a foliage spray every 7 to 14 days beginning about 2 weeks after transplanting. Do not apply within 7 days of harvest. Do not make more than 4 applications per season.

PEANUTS - LEAF SPOT (*Cercospora* spp.) - Apply 11 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage. Begin applications 35 days after planting or when disease first appears and repeat at 14 to 21 day intervals as needed. Use the 14 day interval under severe disease conditions. Do not apply within 14 days of harvest.

PECANS - SCAB (*Fusicladium* sp.), BROWN SPOT (*Cercospora* sp.), Downy Spot (*Mycosphaerella* sp.), and POWDERY MILDEW (*Micosphaera* sp.) - Apply 11 to 23 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage; use the higher rate for trees over 30 feet tall and for aerial application in Arkansas, Louisiana, Mississippi, Oklahoma, and Texas. Begin applications when first leaves are showing and repeat at 3 to 4 week intervals until shuck split. Do not apply after shuck split.

SOYBEANS - ANTHRACNOSE (*Colletotrichum* sp.), BROWN LEAF SPOT (*Septoria* sp.), FROG-EYE LEAF SPOT (*Cercospora* sp.), PURPLE SEED STAIN (*Cercospora* sp.), and POD AND STEM BLIGHT (*Diaporthe* sp. and the imperfect stage, *Phomopsis* sp.) - Apply 11 to 23 fluid ounces of TOPSIN M per acre in sufficient water for thorough coverage. Apply from full bloom to when pods are 1/8" to 1/4" in length. Make a second application 14 to 21 days later. Do not make the second application later than 14 days after pods average 1/4" in length or when beans form in the pod. Use the high rate under severe disease conditions. **FOR SEED BEANS ONLY** - For seed quality make a single application at the high rate when beans form in the pod.

Do not make more than 2 applications per year.
Do not graze or feed treated vines or hay to livestock.

STONE FRUITS - PREHARVEST

PEACHES - MONILINIA BROWN ROT BLOSSOM BLIGHT - Apply 11 to 17 fluid ounces of TOPSIN M per 100 gallons at early bloom and again at full bloom (see note).

1581-252

MONILINIA FRUIT BROWN ROT - In addition to bloom sprays, apply 1 or 2 preharvest sprays at 11 to 17 fluid ounces of TOPSIN M per 100 gallons (see note) starting 3 weeks before harvest. If needed under severe disease pressure, apply additional sprays at 10 to 14 day intervals between full bloom and final preharvest sprays. Do not apply within 1 day of harvest.

NOTE: For Florida, Georgia, South Carolina, North Carolina, Mississippi, and Alabama, use 17 fluid ounces per 100 gallons.

PEACH SCAB - Use same schedule as for Monilinia Brown Rot Blossom Blight plus apply 8 to 11 fluid ounces of TOPSIN M per 100 gallons at shuck split and at first cover spray.

CHERRIES - MONILINIA BROWN ROT BLOSSOM BLIGHT - Apply 11 fluid ounces of TOPSIN M at early and again at full bloom.

MONILINIA FRUIT BROWN ROT - In addition to bloom sprays, apply 1 or 2 sprays at 11 fluid ounces of TOPSIN M per 100 gallons starting 3 weeks before harvest. If needed under severe disease pressure, apply additional sprays at 10 to 14 day intervals between full bloom and final preharvest sprays. Do not apply within 1 day of harvest.

COCCOMYCES CHERRY LEAF SPOT - Apply 8 to 11 fluid ounces of TOPSIN M per 100 gallons at petal fall or before, when leaves first unfold, and first, second, and third cover at 10 to 14 day intervals and one spray 14 to 21 days after harvest. Do not apply within 1 day of harvest.

POWDERY MILDEW - Use same schedule as for Monilinia Brown Rot Blossom Blight plus applications of 8 to 11 fluid ounces of TOPSIN M per 100 gallons at shuck fall and first cover.

APRICOTS and NECTARINES - MONILINIA BROWN ROT BLOSSOM BLIGHT - Apply 11 fluid ounces of TOPSIN M per 100 gallons at early bloom and again at full bloom.

MONILINIA FRUIT BROWN ROT - In addition to bloom sprays, make 1 or 2 applications of TOPSIN M at 11 fluid ounces per 100 gallons starting 3 weeks before harvest. If needed under severe disease pressure, apply additional sprays at 10 to 14 day intervals between full bloom and final preharvest sprays. Do not apply within 1 day of harvest.

PRUNES and PLUMS - MONILINIA BROWN ROT BLOSSOM BLIGHT - Apply 11 fluid ounces of TOPSIN M per 100 gallons at early bloom and again at full bloom.

MONILINIA FRUIT BROWN ROT - In addition to bloom sprays, make 1 or 2 applications at 11 fluid ounces per 100 gallons starting 3 weeks before harvest. If needed under severe disease pressure, apply additional sprays at 10 to 14 day intervals

4581-352

between full bloom and final preharvest sprays. Do not apply within 1 day of harvest.

BLACK KNOT - Apply 11 fluid ounces of TOPSIN M per 100 gallons at pre-bloom, petal fall, and at first, second, and third cover sprays at 10 to 14 day intervals. Do not apply within 1 day of harvest.

COCOMOYCES LEAF SPOT - Apply 11 fluid ounces of TOPSIN M per 100 gallons at petal fall, shuck, and at first, second, and third cover sprays at 10 to 14 day intervals and 1 spray 14 to 21 days after harvest. Do not apply within 1 day of harvest.

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON APPLES

DISEASES

**Bitter Rot, Black Rot, Brown Rot, Cedar Apple Rust,
Fly Speck, Powdery Mildew, Quince Rust,
Scab, Sooty Blotch**

Mixture

Rates

TOPSIN M	11 - 17 fl.oz./A	3 - 4 fl.oz./100 gal
+		
Mantozeb 80WP	3.0-4.0 lb/A	12 - 16 oz./100 gal

Use sufficient volume of spray to provide thorough coverage.

Apply at 1/2 inch green tip and repeat at 7 to 14-day intervals or as needed through the cover sprays. One quart spray oil may be added per 100 gallons. Add oil to the spray tank last.

Use the high rate of TOPSIN M and add spray oil to the spray mixture for varieties more susceptible to powdery mildew, and for scab if an application is missed during an infection period (apply as soon as possible after infection period in order to deactivate scab and to prevent further infection).

**Do not apply within 30 days of harvest.
Do not graze treated areas.**

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

**AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102**

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON APPLES

DISEASES

**Apple Scab, Bitter Rot, Black Rot, Fly Speck,
Powdery Mildew, Sooty Blotch**

<u>Mixture</u>	<u>Rates</u>	
TOPSIN M	11 - 17 fl.oz./A	3 - 4 fl.oz./100 gal
+		
Captan 50WP	3.0-4.0 lb/A	12 - 16 oz./100 gal
or		
Captan 80WP	2.0-2.5 lb/A	7.5- 10 oz./100 gal

Use sufficient volume of spray to provide thorough coverage.

Apply at 1/2 inch green tip and repeat at 7 to 14-day intervals or as needed through the cover sprays. Use the high rate of TOPSIN M for varieties more susceptible to powdery mildew. If an application is missed during an infection period, apply the higher rates as soon as possible after the infection period in order to deactivate scab and to prevent further infection.

NOTE: Spray injury may result if Captan is used with, immediately before, or closely following an oil spray.

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON APPLES

DISEASES

Apple Scab, Bitter Rot, Black Rot, Bull's Eye Rot, Cedar Apple
Rust, Fly Speck, Powdery Mildew, Sooty Blotch

<u>Mixture</u>		<u>Rates</u>
TOPSIN M	11 - 17 fl.oz/A	3 - 4 fl. oz/100 gal
+		
Maneb 80WP	6.0-8.0 lb/A	24 - 32 oz./100 gal (1.5 - 2.0 oz./100 gal)

Use sufficient volume of spray to provide thorough coverage.

Begin applications at green tip and repeat at pink and at 7 to 14-day intervals or as needed through the cover sprays. Use the higher rates and closer interval under severe disease conditions.

With Golden Delicious apples, adverse effects on fruit finish and color may result. Use the lower rates of TOPSIN M when finish and color are of primary importance.

Do not apply within 30 days of harvest.

Do not graze livestock in treated areas.

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON APPLES

DISEASES

**Apple Scab, Bitter Rot, Black Rot, Fly Speck,
Powdery Mildew, Sooty Blotch**

<u>Mixture</u>	<u>Rates</u>	
TOPSIN M	11 - 17 fl.oz./A	2 - 4 fl.oz./100 gal
+		
Polyram 80W	3.0-4.0 lb/A	12 - 16 oz./100 gal

Use sufficient volume of spray to provide thorough coverage.

Begin applications at green tip and repeat applications as needed through the cover sprays.

Do not apply within 30 days of harvest.
Do not graze livestock in treated areas.

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON APPLES

DISEASES

**Bitter Rot, Black Rot, Fly Speck, Frog-eye Leaf Spot,
Powdery Mildew, Sooty Blotch,
Apple Scab**

<u>Mixture</u>	<u>Rates</u>	
TOPSIN M	11 - 17 fl.oz./A	3 - 4 fl.oz./100 gal
+		
Sulfur	12 - 30 lb a.i./ A	48 - 120 oz. a.i. (3.0-7.5 lbs a.i./100 gal)

Use sufficient volume of spray to provide thorough coverage.

Begin applications at green tip and repeat at pink and at 7 to 14-day intervals or as needed through the cover sprays. Use the higher rates and closer interval under severe disease conditions.

Observe the more restrictive cautions and limitations on labeling products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON PEANUTS

DISEASES

Ascochyta Web Blotch, Cercospora Leaf Spot, Rust

Mixture

TOPSIN M
+
Mancozeb 80WP

Rates

6 fl.oz./A
1.0 - 2.0 lb/A

Use sufficient volume of spray to provide thorough coverage.

Begin application 35 to 40 days after planting or when disease first appears. Repeat at the following intervals: for Cercospora leaf spot at 10 to 14 days; for rust at 7 to 10 days; and for Ascochyta web blotch at 7 to 14 days.

Spray oil may be added at the rate of 1 pint to 1 quart per acre.

Do not graze or feed treated vines, hay, or hulls to live-stock.

Do not apply within 14 days of harvest.

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

ACCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Register No. 4581-

NOT FOR CALIFORNIA

TANK MIX FOR USE ON STONE FRUITS
(Apricots, Cherries, Nectarines, Peaches, Plums, and Prunes)

DISEASES

Black Knot, Brown Rot Blossom Blight, Cherry Leaf Spot
(*Coccomyces* sp.), Fruit Brown Rot, Peach Scab,
Powdery Mildew, Russet Scab

Mixture

Rates

TOPSIN M	17 - 23 fl.oz./A	6 - 9 fl.oz./100 gal
+		
Captan 50WP	3.0-6.0 lb/A	16 - 32 oz./100 gal (1.0-2.0 lb/100 gal)

Use sufficient volume of spray to provide thorough coverage.

BROWN ROT BLOSSOM BLIGHT - Apply at early bloom stages before infection occurs (apricots - red bud; cherries - early popcorn; nectarines and peaches - pink bud; plums and prunes - green tip). If blossoming is prolonged or conditions favorable to disease persist, make a second application 14 days later.

FRUIT BROWN ROT - After blossom blight sprays, make a pre-harvest application any time from 3 weeks before harvest to 1 day of harvest. Make a second application if conditions favorable for disease persist or harvest is prolonged.

PEACH SCAB, POWDERY MILDEW - Use same schedule as for Brown Rot Blossom Blight and continue at 10 to 14 day intervals to 1 day of harvest. Make an additional application 2 to 3 weeks after harvest.

CHERRY LEAF SPOT - Use same schedule as for Brown Rot Blossom Blight and continue at 10 to 14 day intervals to 1 day of harvest. Make an additional application 2 to 3 weeks after harvest.

RUSSET SCAB (Plums and Prunes) - Use same schedule as for Brown Rot Blossom Blight; include full coverage spray at full bloom.

BLACK KNOT - Apply at first, second, and third cover sprays at 10 to 14-day intervals.

Do not apply within 1 day of harvest.

NOTE: This tank mixture will not control bacterial blast, but will supplement a dormant program for peach leaf curl and shot-hole (Coryneum blight); refer to other fungicide labels for dormant period control information. Do not use this tank mixture with spray oils.

Observe the more restrictive cautions and limitations on labeling products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON STONE FRUITS
(Apricots, Nectarines, and Peaches)

DISEASES

**Brown Rot Blossom Blight, Fruit Brown Rot, Peach Scab,
Powdery Mildew**

<u>Mixture</u>	<u>Rates</u>
TOPSIN M	34 - 45 fl.oz./A 11 - 14 fl.oz./100 gal
+	
Maneb 80WP	6.0-8.0 lb/A 28 - 40 oz./100 gal (1.75-2.5 lb/100 gal)

Use sufficient volume of spray to provide thorough coverage.

BROWN ROT BLOSSOM BLIGHT - Apply at early bloom stages before infection occurs (apricots - red bud; nectarines and peaches - pink bud). For this application only, TOPSIN M plus Maneb 80 may be used in combination with superior-type spray oil. If blossoming is prolonged or conditions favorable for disease persist, make a second application 14 days later.

FRUIT BROWN ROT - After blossom blight sprays, make an additional application 3 weeks before harvest.

PEACH SCAB, POWDERY MILDEW - Use same schedule as for Brown Rot Blossom Blight plus applications at shuck split, shuck fall, and 14 days later.

Do not apply within 14 days of harvest.

* **NOTE:** This tank mixture will not control bacterial blast, but will supplement a dormant program for peach leaf curl and shot-hole (Coryneum blight); refer to other fungicide labels for dormant period control information.

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON STONE FRUITS
(Apricots, Cherries, Nectarines, Peaches, Plums, and Prunes)

DISEASES

Black Knot, Brown Rot Blossom Blight, Fruit Brown Rot,
Leaf Spots, Peach Canker, Peach Scab, Powdery Mildew,
Rust, Shot-Hole Fungus (Coryneum Blight)

<u>Mixture</u>	<u>Rates</u>	
TOPSIN M	17 - 45 fl.oz./A	6 - 14 fl.oz./100 gal
+		
Sulfur	12 - 24 lb a.i./A	64 - 96 oz. a.i. (4-6 lb a.i./100 gal)

Use sufficient volume of spray to provide thorough coverage.

BROWN ROT BLOSSOM BLIGHT - Make application at early bloom and again at full bloom.

FRUIT BROWN ROT - In addition to bloom sprays, apply 1 or 2 sprays starting 3 weeks before harvest. If needed under severe disease pressure, apply additional sprays at 10 to 14-day intervals between full bloom and final preharvest sprays.

POWDERY MILDEW - In addition to bloom sprays, make applications at shuck fall and first cover.

LEAF SPOTS - Make application when leaves first unfold and first, second, and third cover at 10 to 14-day intervals and 1 spray 14 to 21 days after harvest.

SHOT-HOLE, RUST, PEACH CANKER, AND PEACH SCAB - Apply at pink bloom, petal fall, shuck split, shuck fall, and at 10 to 14-day day intervals as long as necessary.

BLACK KNOT - Apply at first, second, and third cover sprays at 10 to 14-day intervals.

Do not apply within 1 day of harvest.

Observe the more restrictive cautions and limitations on labeling of products used in mixtures.

AGCHEM DIVISION - PENNWALT CORPORATION
Three Parkway, Philadelphia, Pennsylvania 19102

SUPPLEMENTAL LABEL

TOPSIN® M 4F
(Thiophanate-methyl Fungicide)

EPA Registration No. 4581-_____

NOT FOR USE IN CALIFORNIA

TANK MIX FOR USE ON STONE FRUITS
(Apricots, Cherries, Nectarines, and Peaches)

DISEASES

Brown Rot Blossom Blight, Cherry Leaf Spot, Fruit Brown Rot,
Peach Scab, Powdery Mildew, Shot-hole Fungus (Coryneum Blight)

CROP	MIXTURE	RATES	
		Per Acre	Per 100 gal
Apricots	TOPSIN M + Ziram	17-45 fl.oz./A	6-14 fl.oz./100gal
		+ 4.5 lb a.i./A	+ 1.5 lb a.i./100 gal
Cherries	TOPSIN M + Ziram	17-45 fl.oz./A	6-14 fl.oz./100gal
		+ 3.5 lb a.i./A	+ 1.14 lb a.i./100 gal
Nectarines + Peaches	TOPSIN M + Ziram	17-45 fl.oz./A	6-14 fl.oz./100gal
		+ 3.5-4.5 lb a.i./A	+ 1.14-1.52 lb a.i./100gal

Use sufficient volume of spray to provide thorough coverage.

BROWN ROT BLOSSOM BLIGHT - Make application at early bloom and again at full bloom.

FRUIT BROWN ROT - In addition to bloom sprays, apply 1 or 2 sprays starting 3 weeks before harvest. If needed under severe disease pressure, apply additional sprays at 10 to 14-day intervals between full bloom and final preharvest sprays.

POWDERY MILDEW - In addition to bloom sprays, make applications at shuck fall and first cover.

CHERRY LEAF SPOT - Make application when leaves first unfold and first, second, and third cover at 10 to 14-day intervals and 1 spray 14 to 21 days after harvest.

SHOT-HOLE, RUST, PEACH CANKER, AND PEACH SCAB - Apply at pink, bloom, petal fall, shuck split, shuck fall, and at 10 to 14-day intervals as long as necessary.

For apricots, do not apply after early cover periods. For cherries, do not apply within 7 days of harvest. For nectarines and peaches, last application should be made no later than at petal fall.

Observe the more restrictive cautions and limitations on labelling of products used in mixtures.

WARRANTY AND DISCLAIMER

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonable fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. PENNWALT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL PENNWALT OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST; OR OTHER EXPENSES, INCURRED IN PLANTING OR HARVESTING.

Pennwalt and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Pennwalt.