

Reg # 4581-359

PM-21

File
127



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAR 3 1994

Gary R. Sandberg
ELF ATOCHEM NORTH AMERICAN INC.
AGRICHEMICALS DIVISION
2000 MARKET ST. 21ST FLOOR
PHILADELPHIA, PA 19103

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 08/13/93 in Response to PR Notice 93-7
EPA Reg. No. 4581-359
MANEB PLUS ZINC F4 FUNGICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is **accepted**. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,


Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

BEST AVAILABLE COPY

tochem

ATO

SHAKE WELL BEFORE USING

PROPOSED LABEL

MANEB PLUS ZINC F4

Fungicide

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 3 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

4581-359

ACTIVE INGREDIENT:

Maneb (Manganese Ethylenebisdithiocarbamate) 37%
(Equivalent to manganese metallic 7.6%)

INERT INGREDIENTS: 63%

TOTAL 100%

This product contains 4 pounds Maneb per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la
explique a usted en detalle. (If you do not understand the
label, find someone to explain to you in detail).

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, induce vomiting immediately by giving two glasses of
water and sticking finger down throat. Call a physician. Never give any-
thing by mouth to an unconscious person.

IF IN EYES, flush eyes with plenty of water for at least 15 minutes. Get
medical attention if irritation persists.

IF ON SKIN, immediately flush skin with plenty of water. Get medical
attention if irritation persists.

See Inside Panel for Additional Precautionary Statements

See following pages for General Information, Directions for Use, and Chemigation
Instructions. Do not apply this product through any irrigation system unless these
instructions are followed.

EPA Registration No. 4581-359

EPA Establishment No. 4581-TX-1

Net Contents 2.5 U.S. Gallon/9.46 Liters

Sold by:

ELF ATOCHEM NORTH AMERICA, INC.—AGCHEM DIVISION
Philadelphia, PA

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

HAZARDOUS IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED.
CAUSES MODERATE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Avoid breathing mist.

Personal Protective Equipment

APPLICATORS AND OTHER HANDLERS MUST WEAR:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear when mixing or loading
- Chemical-resistant apron when cleaning equipment, mixing or loading

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Human flaggers must be in enclosed cabs.

The enclosed cabs must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)]. The handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:
Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, except as specified for the labeled use on cranberries in areas where surface water is present or to riparian areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area of application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It contains specific instructions, exceptions pertaining to the uses on this label about personal protective equipment (PPE), restricted-entry interval, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

BEST AVAILABLE COPY

DIRECTIONS FOR USE AS A SPRAY

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR GROUND APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, minimum of 10 gallons. Increased volume of water may be necessary as foliage density increases.

FOR AIR APPLICATION use at the rate indicated in sufficient water for thorough coverage, a minimum of 3 gallons per acre for field crops and 10 gallons per acre for orchard crops.

When dosage ranges are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate or apply more frequently than the minimum interval given in the directions for that crop.

If this product is used on a crop, no other product containing a different EBOC active ingredient may be used on the same crop during the same growing season.

597

VEGETABLE CROP USES

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
Beans (Dry)	Anthrachnose Downy Mildew Rust	1.2-1.6	Begin when plants are small. Spray on a 5-7 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 30 days of harvest.
Broccoli Brussels Sprouts Cauliflower	Alternaria Leaf Spot Downy Mildew	1.2-1.6	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 7 days of harvest.
Cabbage Kohlrabi Chinese Cabbage (light-headed only)	Alternaria Leaf Spot Downy Mildew	1.2-1.6	Plant beds and direct seeded fields: Apply when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply later than 7 days before harvest.
Chinese Cabbage (loose head) (California and Hawaii Only)	Alternaria Leaf Spot Downy Mildew	0.8-1.2	Plant beds and direct seeded fields: Apply when disease threatens. Spray on a 7-10 day interval. Do not apply more than 7.2 lbs. active ingredient (7.2 qts. product) per acre per season. Do not apply later than 7 days before harvest in California and 10 days before in Hawaii.
Sweet Corn Popcorn Sweet Corn (for seed production)	Common Rust Helminthosporium Blight	1.2	East of the Mississippi (including Arkansas & Louisiana) Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 18 lbs. active ingredient (18 qts. product) per acre per season. Do not feed treated forage to livestock.
		1.2	West of the Mississippi (except Arkansas & Louisiana) Use sufficient water for thorough coverage. Start applications when disease first appears at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 6.0 lbs. active ingredient (6.0 qts. product) per acre per season. Do not feed treated forage to livestock.
Cucumbers	Alternaria (Mosaicvirus) Leaf Spot Angular Leaf Spot Anthrachnose Downy Mildew Pythium Fruit Rot	1.2-1.6	Begin when disease threatens or plants begin to run. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Eggplant	Anthrachnose Septoria Leaf Spot Cladosporium Leaf Mold Early Blight Late Blight Gray Leaf Spot (Stemphylium)	1.2-1.6	Begin at first fruit cluster and repeat at 7-10 day intervals. Do not apply more than 11.2 lbs. active ingredient (11.2 qts. product) per acre per crop. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot Downy Mildew	1.2-1.6	Begin when disease threatens and apply on a 7-10 day interval. Do not use more than 3.2 lbs. active ingredient (3.2 qts. product) per cutting. Do not apply within 10 days of harvest.

VEGETABLE CROP USES (continued)

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
Lettuce (Head and Leaf) Endive	Downy Mildew	1.2-1.6	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming. California—Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient maximum (6.4 qts. product) per acre per season. Do not apply within 14 days of harvest.
Melons including cantaloupes, cantaloups, cantaloups, honeydews, watermelons	Alternaria Leaf Spot Anthrachnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight	1.2-1.6	Apply as soon as plants begin to run or when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Onions (Dry Bulb)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	1.6-2.4	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 24 lbs. active ingredient (24 qts. product) per acre per season. Do not apply to exposed bulbs.
Onions (Green)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	1.6-2.4	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 16.8 lbs. active ingredient (16.8 qts. product) per acre per season. Do not apply to exposed bulbs.
(Furrow Drench)	Smut	2.4	Apply as a furrow drench at time of planting. Onion seed: Use 75 to 25 gallons water per acre. Do not use more than 2.4 lbs. active ingredient (2.4 qts. product) per 29,000 linear feet of row (10 inch spacing). Do not use in California.
Peas	Anthrachnose Cercospora Leaf Spot (Bacterial Spot) Phytophthora Root Rot	1.2-2.4	East of the Mississippi: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 14.4 lbs. active ingredient (14.4 qts. product) per acre per season. Do not apply within 7 days of harvest.
		2-1	West of the Mississippi: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 7 days of harvest.
Potatoes	Early Blight Late Blight	2-1.6	Begin when plants are 2 to 5 inches high. Apply on a 5-10 day interval. Do not apply more than 11.2 lbs. active ingredient (11.2 qts. product) per acre per season. Do not apply within 14 days of harvest except for: Connecticut, Delaware, Florida, Maine, Massachusetts, Michigan, New Hampshire, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Wisconsin where the PH is 3 days. It is recommended that this product be used with an Integrated Pest Management Program. First IPI should occur 14 days before harvest.
Potatoes (Seed Pieces)	Fusarium Seed Piece Decay	8.8 qt. per 10 gal.	Seed piece treatment only. Dip whole or cut tubers. Spread in cool place if hot before planting. Do not use treated seed piece for food, feed, or oil purposes.
Pumpkins	Angular Leaf Spot Downy Mildew	1.2-1.6	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Squash (Summer and Winter)	Anthrachnose Downy Mildew	1.2-1.6	Begin when plants start to run. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Tomatoes (Greenhouse and Field)	Anthrachnose Septoria Leaf Spot Cladosporium Leaf Mold Early Blight Late Blight Gray Leaf Spot (Stemphylium)	1.2-2.4	East of the Mississippi: Begin at first fruit cluster and repeat on a 7-10 day interval. Do not apply within 5 days of harvest. Do not apply more than 16.8 lbs. active ingredient (16.8 qts. product) per acre per crop.
		1.2-1.6	West of the Mississippi: Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient (6.4 qts. product) per acre per season. Do not apply within 5 days of harvest.

BEST AVAILABLE COPY

697

FIELD CROPS

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
Sugarbeets	Cercospora Leaf Spot	1.2-1.6	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 11.2 lbs active ingredient (11.2 qts product) per season. Do not apply within 14 days of harvest.

FRUIT AND NUT CROPS

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
FOR POME FRUITS Use either the "Pre-Bloom" or "Extended Application" schedules. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.			
Apples	Rusts (including Cedar apple) Fabrya Leaf Spot Physcok Scab Sooty Blotch	4.8 or (1.2 qts per 100 gal in dilute spray)	PRE-BLOOM: Begin applications at 1/4 to 1/2 inch green tip and continue on a 7- to 10-day application schedule through bloom. Do not apply more than 19.2 lbs active ingredient (19.2 qts product) per acre per year. Do not graze livestock in treated areas.
		2.4 or (0.6 qts per 100 gal in dilute spray)	EXTENDED APPLICATION OR TANK MIX: For implementation of IPM programs, applications based on tree-row volume or for use as a resistance management tool: begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7- to 10-day schedule through the second cover spray or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not apply more than 16.8 lbs active ingredient (16.8 qts product) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program. Add spreader/sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/curative fungicide for more effective control of diseases.
Almonds	Brown Rot Blossom and Twig Blight Scab Shot-hole Fungus (Leaf Blight)	4.8-6.4 or (1.2-1.6 qts per 100 gal in a dilute spray)	Apply at popcorn, full bloom, and petal fall or every 7 to 10 days if bloom is staggered. Omit petal fall spray if only brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not apply more than 25.6 lbs active ingredient (25.6 qts product) per acre per season. Do not allow livestock to graze in almond orchards.
Bananas	Sagatoka	1.6-2.4	Apply when leaves first appear and repeat as needed on a 14-21 day interval. Do not apply more than 24 lbs active ingredient (24 qts product) per acre per season. May be applied up to the day of harvest.
Cranberries	Fruit Rot	3.8-4.8	Begin applications at mid-bloom and repeat at 7-10 day interval. Do not apply more than 14.4 lbs active ingredient (14.4 qts product) per season. Do not apply within 30 days of harvest.
Grapes	Black Rot Bunch Rot Deadarm Downy Mildew	1.2-3.2 or (0.3-0.8 qts per 100 gal in a dilute spray)	East of the Rocky Mountains: Apply in sufficient water to provide thorough coverage starting when shoots are 1/4 to 1 1/2 inches long and continue at 7-10 day intervals until fruit is set. Do not apply within 66 days of harvest. Do not apply more than 19.2 lbs active ingredient (19.2 qts product) per acre per season.
		1.2-2.0 or (0.3-0.5 qts per 100 gal in a dilute spray)	West of the Rocky Mountains: Apply when shoots are 1/4 to 1 1/2 inches long and continue at 7-10 day intervals. Do not apply more than 6.0 lbs active ingredient (6.0 qts product) per acre per season. Do not apply within 66 days of harvest except in California. California—Do not apply after bloom.
Kadota figs	Surface Mold (Cladosporium) Surface Rot (Alternaria)	0.6 qts 100 gal	Apply when disease threatens in 100-400 gallons of water/acre. Do not apply more than 2.4 lbs active ingredient (2.4 qts product) per acre per season. Do not apply within 10 days of harvest.
Papayas	Anthrachnose Phytophthora Fruit Rot Cercospora Black Spot	1.6-2.0 or (0.4-0.5 qts per 100 gal in a dilute spray)	Apply when disease first threatens and continue at 14-21 day intervals. Do not apply more than 28 lbs active ingredient per season (28 qts product). May be applied up to the day of harvest.

COMMERCIAL ORNAMENTAL CROP USES (NOT FOR HOME GARDEN USE)

DO NOT USE ANY OF THESE CROPS FOR FOOD OR FEED PURPOSES

CROP	DISEASES	RATE QT/100 GAL	DIRECTIONS
Carnations	Alternaria Leaf Spot Anthrachnose Botrytis Blight	0.8-1.2	Begin when new growth starts. Repeat weekly.
Chrysanthemums	Ascochyta Ray Blight Botrytis Petal Spot Septoria Leaf Spot	12.8 fl oz PLUS Captan 50W 3/4 lbs per 100 gal water	Apply twice weekly during the blooming season for Petal Spot and Ray Blight. For Septoria Leaf Spot apply weekly throughout the season.
Dahlias	Alternaria Leaf Spot Botrytis Blight	0.8-1.2	Begin when new growth starts. Repeat weekly.
Dogwood	Anthrachnose	0.8-1.2	Begin when buds open. Repeat when bracts fall 4 weeks later and in later summer.
Gladiolus	Botrytis Blight Curvularia Stemphylium Leaf Spots	1.2	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Grasses Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.	Brown Patch	4.8 fl oz in sufficient water/1000 sq ft	Begin at first sign of disease. Repeat at 7- to 14-day intervals.
	Dollar Spot	9.6-12.8 fl oz in sufficient water/1000 sq ft	Apply as for Brown Patch, above.
	Melting-Out	4.8-6.4 fl oz in sufficient water/1000 sq ft	Apply as for Brown Patch, above.
	Leaf, Stem and Sheath Rusts	2.4	Begin when rust pustules are first seen. Repeat at 7- to 14-day intervals. Do not graze treated areas. Do not feed clippings to livestock.
Lilies	Botrytis Blight	0.8-1.2	Begin with new growth. Repeat weekly.
Pansies	Anthrachnose	0.8-1.2	Begin with new growth. Repeat weekly.
Peonies	Alternaria Leaf Spot Botrytis Blight Phytophthora Blight	0.8-1.2	Apply to foliage and soil in early spring and early fall and 7- to 10-day intervals during the growing season.
Roses	Black Spot Cercospora Leaf Spot (Texas) Rust (California)	0.8-1.2	Begin when first leaves unfold. Repeat at 7- to 10-day intervals. For Aerial Application, use 0.8 to 1.2 qts product in a minimum of 3 gallons spray solution per acre.
Snapdragons	Rust	0.8-1.2	Begin with emergence. Repeat weekly.
Zinnias	Alternaria Blight Leaf Spot Botrytis Blight	0.8-1.2	Begin with emergence. Repeat weekly.

CHEMIGATION NOT FOR THIS USE IN CALIFORNIA GENERAL INSTRUCTIONS

Apply this product only through sprinkler irrigation systems including center pivot, lateral move, and low, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

BEST AVAILABLE COPY

727

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

INSTRUCTIONS FOR SPRINKLER (OVERHEAD) IRRIGATION:

Observe the requirements in the System Requirements section above.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply MANEB PLUS ZINC F4 only through systems containing anti-siphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

MANEB PLUS ZINC F4 may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

Check local restrictions and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

SPRAY PREPARATION:

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

Prepare a suspension of MANEB PLUS ZINC F4 in a mix tank. Fill the tank with $\frac{1}{2}$ or $\frac{3}{4}$ the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of MANEB PLUS ZINC F4 and then the remaining volume of water.

APPLICATION INSTRUCTIONS:

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy. Start sprinkler and then uniformly inject the suspension of MANEB PLUS ZINC F4 into the irrigation water line so as to deliver the desired rate per acre. The suspension of MANEB PLUS ZINC F4 should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

NOTE: When treatment with MANEB PLUS ZINC F4 has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage Instructions: Store in the original container. Keep away from fire and sparks. Store in a cool place, but not below 32°F. Do not allow product to become overheated in storage as this may bring on chemical changes which will impair the fungicide effectiveness of MANEB PLUS ZINC F4 Fungicide. Keep container closed when not in use. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Instructions: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

Elf Atochem North America warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. ELF ATOCHEM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL ELF ATOCHEM OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION OR CUSTOMERS LABOR COST, OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

Elf Atochem and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Elf Atochem.

Made and Printed in U.S.A.

5-Z324S-03 C3 (3/83)

BEST AVAILABLE COPY

(8/93)