

JUL 26 1994

Gary R. Sandberg
Elf Atochem North America, Inc.
Three Parkway, Philadelphia, PA 19102

Dear Mr. Sandberg:

Subject: Label Amendment - PR Notice 93-3/93-8
Maneb Plus Zinc F4
EPA Registration Number 4581-359
Your Submission Dated August 17, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit three (3) copies of final printed labeling for our records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 156, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely Yours,

Steve Robbins *SR*
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

CONCURRENCES								
SYMBOL	7505C							
SURNAME	Robbins							
DATE	7/26/94							

elf atochem

ATO

SHAKE WELL BEFORE USE

PROPOSED LABEL

MANEB PLUS ZINC F4

Fungicide

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUL 26 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
4581-359

ACTIVE INGREDIENT:

Maneb (Manganese Ethylenebisdithiocarbamate) 37%
(Equivalent to manganese metallic 7.6%)

INERT INGREDIENTS: 63%

TOTAL 100%

This product contains 4 pounds Maneb per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES, flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN, wash with plenty of soap and water. Get medical attention if irritation persists.

See Inside Panel for Additional Precautionary Statements

EPA Registration No. 4581-359

EPA Establishment No. 4581-TX-1

Net Contents 2.5 U.S. Gallon/9.46 Liters

BEST AVAILABLE COPY

Sold by:

ELF ATOCHEM NORTH AMERICA, INC.—AGCHEM DIVISION
Philadelphia, PA

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED. CAUSES MODERATE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Avoid breathing mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

This product is toxic to fish.

Except as specified for the labeled use on cranberries, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

GENERAL INFORMATION

Do not use this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas within 24 hours of application unless wearing the protective clothing listed below. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

After sprays have dried, do not enter or allow entry into treated areas or areas where there is a danger of drift until the 24-hour reentry interval has expired unless wearing the personal protective equipment listed on this label.

Keep all unprotected persons, children, livestock, and pets away from treated area or where there is danger of drift.

Do not rub eyes or mouth with hands. See Statement of Practical Treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PERSONAL PROTECTIVE EQUIPMENT

HANDLERS (MIXERS, LOADERS, AND APPLICATORS) AND EARLY REENTRY WORKERS MUST WEAR THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT: coveralls over long-sleeved shirt and long pants, shoes, socks, and chemical-resistant gloves. During mixing and loading, a chemical-resistant apron and goggles or a face shield must also be worn. For agricultural workers, when completely enclosed cabs with positive pressure filtration or an enclosed cockpit for aerial application are used, a long-sleeved shirt and long pants may be worn in place of the protective clothing described above. Chemical-resistant gloves must be available in the cab or cockpit and worn upon exiting. The gloves must be kept in an enclosed container in the cab or cockpit to prevent contamination of the inside of the cab or cockpit.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face, and arms with soap and water before eating, smoking or drinking. Always wash hands and arms with soap and water before using the toilet.

After work take off all clothes and shoes. Shower using soap and water. Wear only clean clothes. Do not use contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment drenched with maneb must be destroyed according to state and local regulations.

DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

During aerial application, human flaggers are prohibited unless in totally enclosed vehicles.

REENTRY STATEMENT

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, such warnings shall be given in a language customarily understood by the workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by the workers. Written warnings must include the following information: "CAUTION. Area treated with MANEB PLUS ZINC F4 Fungicide on (date of application). Do not enter without protective clothing within 24 hours after application."

DIRECTIONS FOR USE AS A SPRAY

FOR GROUND APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, minimum of 10 gallons. Increased volume of water may be necessary as foliage density increases.

FOR AIR APPLICATION use at the rate indicated in sufficient water for thorough coverage, a minimum of 3 gallons per acre for field crops and 10 gallons per acre for orchard crops.

When dosage ranges are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate or apply more frequently than the minimum interval given in the directions for that crop.

If this product is used on a crop, no other product containing a different EBDC active ingredient may be used on the same crop during the same growing season.

VEGETABLE CROP USES

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
Beans (Dry)	Anthracnose Downy Mildew Rust	1.2-1.6	Begin when plants are small. Spray on a 5-7 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 30 days of harvest.
Broccoli Brussels Sprouts Cauliflower	Alternaria Leaf Spot Downy Mildew	1.2-1.6	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 7 days of harvest.
Cabbage Kohlrabi Chinese Cabbage (light-headed only)	Alternaria Leaf Spot Downy Mildew	1.2-1.6	Plant beds and direct seeded fields: Apply when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply later than 7 days before harvest.
Chinese Cabbage (loose head) (California and Hawaii only)	Alternaria Leaf Spot Downy Mildew	0.8-1.2	Plant beds and direct seeded fields: Apply when disease threatens. Spray on a 7-10 day interval. Do not apply more than 7.2 lbs. active ingredient (7.2 qts. product) per acre per season. Do not apply later than 7 days before harvest in California and no days before in Hawaii.
Sweet Corn Popcorn Sweet Corn (used for seed production)	Downy Mildew Helminthosporium Blight	1.2	East of the Mississippi (including Arkansas, Louisiana) Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3-10 day intervals. Add recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 18 lbs. active ingredient (18 qts. product) per acre per season. Do not feed treated forage to livestock.
		1.2	West of the Mississippi (except Arkansas & Louisiana) Use sufficient water for thorough coverage. Start applications when disease first appears at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 6.0 lbs. active ingredient (6.0 qts. product) per acre per season. Do not feed treated forage to livestock.
Cucumbers	Alternaria (Blasporium) Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Pythium Fruit Rot	1.2-1.6	Begin when disease threatens or plants begin to run. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Eggplant	Anthracnose Septoria Leaf Spot Cladosporium Leaf Mold Early Blight Late Blight Gray Leaf Spot (Strombosphaera)	1.2-1.6	Begin at first fruit cluster and repeat at 7-10 day intervals. Do not apply more than 11.2 lbs. active ingredient (11.2 qts. product) per acre per crop. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot Downy Mildew	1.2-1.6	Begin when disease threatens and apply on a 7-10 day interval. Do not use more than 3.2 lbs. active ingredient (3.2 qts. product) per cutting. Do not apply within 10 days of harvest.

BEST AVAILABLE COPY

VEGETABLE CROP USES (continued)

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
Lettuce (Head and Leaf) Endive	Downy Mildew	1.2-1.6	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs active ingredient (9.6 qts. product) per acre per season. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and blanching. California—Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient maximum (6.4 qts. product) per acre per season. Do not apply within 14 days of harvest.
Melons including: cantaloupes, cantaloups, cantaloupes, honeydews, watermelons	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight	1.2-1.6	Apply as soon as plants begin to run or when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Onions (Dry Bulb)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	1.6-2.4	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 24 lbs. active ingredient (24 qts. product) per acre per season. Do not apply to exposed bulbs.
Onions (Green)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	1.6-2.4	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 16.8 lbs. active ingredient (16.8 qts. product) per acre per season. Do not apply to exposed bulbs.
(Furrow Dranch)	Smut	2.4	Apply as a furrow dranch at time of planting. Use 75 to 25 gallons of water per acre. Do not use more than 2.4 lbs. active ingredient (2.4 qts. product) per 29,000 linear feet of row (10 inch spacing). Do not use in California.
Peanut	Anthracnose Cercospora Leaf Spot (Ringspot) Phytophthora Blight Root Rot	1.2-2	East of the Mississippi Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 14.4 lbs active ingredient (14.4 qts. product) per acre per season. Do not apply within 7 days of harvest.
Potatoes	Early Blight Late Blight	2-1	West of the Mississippi Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (9.6 qts. product) per acre per season. Do not apply within 14 days of harvest.
Potatoes (Seed Pieces)	Fusarium Seed Piece Decay	0.8 qt. per 10 gal.	Seed piece treatment only. Dip whole or cut tubers. Spread in cool place if held before planting. Do not use treated seed piece for food, feed, or oil purposes.
Pumpkins	Angular Leaf Spot Downy Mildew	1.2-1.6	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Squash (Summer and Winter)	Anthracnose Downy Mildew	1.2-1.6	Begin when plants start to run. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 qts. product) per acre per season. Do not apply within 5 days of harvest.
Tomatoes (Greenhouse and Field)	Anthracnose Septoria Leaf Spot Cladosporium Leaf Mold Early Blight Late Blight Gray Leaf Spot (Stemphylium)	1.2-2.4	East of the Mississippi Begin at first fruit cluster and repeat on a 7-10 day interval. Do not apply within 5 days of harvest. Do not apply more than 16.8 lbs active ingredient (16.8 qts. product) per acre per crop.
		1.2-1.6	West of the Mississippi Spray on a 7-10 day interval. Do not apply more than 6.4 lbs active ingredient (6.4 qts. product) per acre per season. Do not apply within 5 days of harvest.

FIELD CROPS

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
Sugarbeets	Cercospora Leaf Spot	1.2-1.6	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 11.2 lbs active ingredient (11.2 qts. product) per season. Do not apply within 14 days of harvest.

FRUIT AND NUT CROPS

CROPS	DISEASES	RATE QTS/ACRE	DIRECTIONS
FOR POME FRUITS Use either the "Pre-Bloom" or "Extended Application" schedules. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.			
Apples	Rusts (including Cedar apple) Fabra Leaf Spot Physcok Scab Sooty Blotch	4.8 or (1.2 qts. per 100 gal. in dilute spray)	PRE-BLOOM: Begin applications at 1/4 to 1/2 inch green tip and continue on a 7- to 10-day application schedule through bloom. Do not apply more than 19.2 lbs. active ingredient (19.2 qts. product) per acre per year. Do not graze livestock in treated areas.
		2.4 or (0.6 qts. per 100 gal. in dilute spray)	EXTENDED APPLICATION OR TANK MIX: For a implementation of IPM programs, applications based on tree-row volume or for use as a resistance management tool: Begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7- to 10-day schedule through the second cover spray or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not apply more than 16.8 lbs. active ingredient (16.8 qts. product) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program. Add spreader/sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/curative fungicide for more effective control of diseases.
Almonds	Brown Rot Blossom and Twig Blight Scab Shot-hole Fungus (Leaf Blight)	4.8-6.4 or (1.2-1.6 qts. per 100 gal. in a dilute spray)	Apply at popcorn, full bloom, and petal fall or every 7 to 10 days if bloom is staggered. Omit petal fall spray if only brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not apply more than 25.6 lbs. active ingredient (25.6 qts. product) per acre per season. Do not allow livestock to graze in almond orchards.
Dananas	Sigatoka	1.6-2.4	Apply when leaves first appear and repeat as needed on a 14-21 day interval. Do not apply more than 24 lbs. active ingredient (24 qts. product) per acre per season. May be applied up to the day of harvest.
Cranberries	Fruit Rot	3.0-4.2	Begin applications at mid-bloom and repeat at 7-10 day interval. Do not apply more than 14.4 lbs. active ingredient (14.4 qts. product) per season. Do not apply within 30 days of harvest.
Grapes	Black Rot Bunch Rot Deadarm Downy Mildew	1.2-3.2 or (0.3-0.8 qts. per 100 gal. in a dilute spray)	East of the Rocky Mountains Apply in sufficient water to provide thorough coverage starting when shoots are 1/4 to 1 1/2 inches long and continue at 7-10 day intervals until fruit is set. Do not apply within 85 days of harvest. Do not apply more than 19.2 lbs. active ingredient (19.2 qts. product) per acre per season.
		1.2-2.0 or (0.3-0.5 qts. per 100 gal. in a dilute spray)	West of the Rocky Mountains Apply when shoots are 1/4 to 1 1/2 inches long and continue at 7-10 day intervals. Do not apply more than 6.0 lbs. active ingredient (6.0 qts. product) per acre per season. Do not apply within 85 days of harvest except in California. California—Do not apply after bloom.
Kadota Figs	Surface Mould (Cladosporium) Surface Rot (Alternaria)	0.6 qts. / 100 gal.	Apply when disease threatens in 100-400 gallons of water per acre. Do not apply more than 2.4 lbs. active ingredient (2.4 qts. product) per acre per season. Do not apply within 10 days of harvest.
Papayas	Anthracnose Phytophthora Fruit Rot Cercospora Black Spot	1.6-2.0 or (0.4-0.5 qts. per 100 gal. in a dilute spray)	Apply when disease first threatens and continue at 14-21 day intervals. Do not apply more than 28 lbs active ingredient per season (28 qts. product). May be applied up to the day of harvest.

BEST AVAILABLE COPY

**COMMERCIAL ORNAMENTAL CROP USES
(NOT FOR HOME GARDEN USE)
DO NOT USE ANY OF THESE CROPS FOR FOOD OR FEED PURPOSES**

CROP	DISEASES	RATE QT/100 GAL	DIRECTIONS
Carnations	Alternaria Leaf Spot Anthracnose Botrytis Blight	0.8-1.2	Begin when new growth starts Repeat weekly
Chrysanthemums	Ascochyta Ray Blight Botrytis Petiole Spot Septoria Leaf Spot	12.8 fl oz PLUS Captan 50W 3/4 lbs per 100 gal water	Apply twice weekly during the blooming season for Petiole Spot and Ray Blight. For Septoria Leaf Spot apply weekly throughout the season.
Dahlias	Alternaria Leaf Spot Botrytis Blight	0.8-1.2	Begin when new growth starts Repeat weekly.
Dogwood	Anthracnose	0.8-1.2	Begin when buds open. Repeat when bracts fall 4 weeks later and in later summer.
Glaucolus	Botrytis Blight Curvularia and Stemphylium Leaf Spots	1.2	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Grasses Do not use on pasture or range grasses. Do not use clippings to stock. Do not use if grass is green for seed.	Brown Patch	4.8 fl oz in sufficient water/1000 sq ft	Begin at first sign of disease. Repeat at 7- to 14-day intervals.
	Dollar Spot	9.6-12.8 fl oz in sufficient water/1000 sq ft	Apply as for Brown Patch, above
	Melting-Out	4.8-6.4 fl oz in sufficient water/1000 sq ft	Apply as for Brown Patch, above
	Leaf, Stem and Sheath Rusts	2.4	Begin when rust pustules are first seen. Repeat at 7- to 14-day intervals. Do not graze treated areas. Do not feed clippings to livestock.
Lilies	Botrytis Blight	0.8-1.2	Begin with new growth. Repeat weekly.
Panicle	Anthracnose	0.8-1.2	Begin with new growth. Repeat weekly.
Panicle	Alternaria Leaf Spot Botrytis Blight Phytophthora Blight	0.8-1.2	Apply to foliage and soil in early spring and early fall and 7- to 10-day intervals during the growing season.
Petas	Black Spot Cercospora Leaf Spot (Texas) Rust (California)	0.8-1.2	Begin when first leaves unfold. Repeat at 7- to 10-day intervals. For Aerial Application, use 0.8 to 1.2 qts. product in a minimum of 3 gallons spray solution per acre.
Snapdragons	Rust	0.8-1.2	Begin with emergence. Repeat weekly.
	Alternaria Blight Leaf Spot Botrytis Blight	0.8-1.2	Begin with emergence. Repeat weekly.

**CHEMIGATION
NOT FOR THIS USE IN CALIFORNIA
GENERAL INSTRUCTIONS**

Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end row, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

INSTRUCTIONS FOR SPRINKLER (OVERHEAD) IRRIGATION:

Observe the requirements in the System Requirements section above.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply MANEB PLUS ZINC F4 only through systems containing anti-siphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

MANEB PLUS ZINC F4 may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

Check local restrictions and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

SPRAY PREPARATION:

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

Prepare a suspension of MANEB PLUS ZINC F4 in a mix tank. Fill the tank with 1/2 or 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of MANEB PLUS ZINC F4 and then the remaining volume of water.

APPLICATION INSTRUCTIONS:

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy. Start sprinkler and then uniformly inject the suspension of MANEB PLUS ZINC F4 into the irrigation water line so as to deliver the desired rate per acre. The suspension of MANEB PLUS ZINC F4 should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

NOTE: When treatment with MANEB PLUS ZINC F4 has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage Instructions: Store in the original container. Keep away from fire and sparks. Store in a cool place, but not below 32°F. Do not allow product to become overheated in storage as this may bring on chemical changes which will impair the fungicide effectiveness of MANEB PLUS ZINC F4 Fungicide. Keep container closed when not in use. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Instructions: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

El Alchem North America warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. EL ALCHEM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL EL ALCHEM OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS, LABOR COST, OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

El Alchem and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of El Alchem.

Made and Printed in U.S.A.

5-Z324S-03 C3 (3/93)

BEST AVAILABLE COPY

(8/93)