

Pp

4581-371

100 100

AUG 7 1993

Mr. Gary R. Sandberg
Elf Atochem N.A., Inc.
2000 Market Street
Agchem Division, 21st Floor
Philadelphia, PA 19103

Dear Mr. Sandberg:

Subject: Maneb 75DF Dry Flowable Fungicide
EPA Reg. No. 4581-371
Your Submission of November 30, 1993

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration or reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- a. Add the heading "Foliar Applications" above the section beginning "When EBDC Products Used ---". Following these sections, add the heading "Seed Treatment" under which you must add "In addition to the maximum number of foliar applications permitted by the formula stated above, a single application or seed treatment may be made on crops which have registered seed treatment uses".


- b. The amended labeling does not contain the required Worker Protection Standard labeling. Assure that the WPS labeling which was accepted for this product is incorporated into this label.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


Steve Robbins
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)


Enclosure

7505C:C.Grable:cg:8/17/94

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


Steve Robbins
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

7505C:C.Grable:cg:8/17/94

elf atochem

ATO

MANEB 75DF

DRY FLOWABLE FUNGICIDE

ACCEPTED
with COMMENTS
in EPA Letter Dated

AUG 17 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

4581-371

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate) 75.0%
(Manganese equivalent as metallic 16.5%)

INERT INGREDIENTS: 25.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, induce vomiting immediately by giving two glasses of water and sticking finger down throat. Call a physician. Never give anything by mouth to an unconscious person.

IF INHALED, remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

IF ON SKIN, immediately flush skin with plenty of water. Get medical attention if irritation persists.

See inside panel for additional Precautionary Statements.

See following pages for General Information, Directions for Use, and Chemigation Instructions. Do not apply this product through any type of irrigation system unless these instructions are followed.

EPA Registration No 4581-371

EPA Establishment No. 33910-HL-01

Net Contents _____ lbs. (_____ kg)

Sold by
ELF ATOCHEM NORTH AMERICA, INC.—AGCHEM DIVISION
Philadelphia, PA

BEST AVAILABLE COPY

C
C
C
D
Dr
GI
Gr
Dr
pa
gr
ter
to
Dr
gr
for
LW
Pa
Pe
Ro
Sna
Zer
App
move
produ
Cro
nonun
If y
specia
A p
under
neces
Do r
a syst
15 ser
days o
Svs
follow
The
drain
from b
The
valve t

COMMERCIAL ORNAMENTAL CROP
(NOT FOR HOME GARDEN USE)
DO NOT USE ANY OF THESE CROPS FOR FOOD OR FEED PURPOSES

CROP	DISEASES	RATE LBS/100 GAL	DIRECTIONS
Carnations	Alternaria Leaf Spot Anthracnose Botrytis Blight	1.0-1.5	Begin when new growth starts. Repeat weekly
Chrysanthemums	Ascochyta Ray Blight Botrytis Petal Spot Septoria Leaf Spot	0.5 lbs or 227 grams	Apply twice weekly during the blooming season for Petal Spot and Ray Blight. For Septoria Leaf Spot apply weekly throughout the season.
Dahlias	Alternaria Leaf Spot Botrytis Blight	1.0-1.5	Begin when new growth starts. Repeat weekly.
Dogwood	Anthracnose	1.0-1.5	Begin when buds open. Repeat when bracts fall 4 weeks later and in late summer.
Gladiolus	Botrytis Blight Curvularia and Sclerotium Leaf Spots	1.5	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Grasses Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed	Brown Patch	0.2 lbs or 90 grams	Begin at first sign of disease. Repeat at 7 to 14 day intervals.
	Dollar Spot	0.4-0.5 lbs or 170-227 grams	Apply as for Brown Patch, above
	Melting-Out	0.2-0.3 lbs or 85-113 grams	Apply as for Brown Patch, above
	Leaf, Stem and Stripe Rusts	3.0	Begin when rust pustules are first seen. Repeat at 7 to 14 day intervals. Do not graze treated areas. Do not feed clippings to livestock
Lilies	Botrytis Blight	1.0-1.5	Begin with new growth. Repeat weekly
Pansies	Anthracnose	1.0-1.5	Begin with new growth. Repeat weekly
Peonies	Alternaria Leaf Spot Botrytis Blight Phytophthora Blight	1.0-1.5	Apply to foliage and soil in early spring and early fall and 7 to 10 day intervals during the growing season
Roses	Black Spot Cercospora Leaf Spot (Texas) Rust (California)	1.0-1.5	Begin when first leaves unfold. Repeat at 7 to 10 day intervals
Snappedragons	Rust	1.0-1.5	Begin with emergence. Repeat weekly
Zinnias	Alternaria Blight Leaf Spot Botrytis Blight	1.0-1.5	Begin with emergence. Repeat weekly

CHEMIGATION NOT FOR THIS USE IN CALIFORNIA ... GENERAL INSTRUCTIONS

Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

... SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

INSTRUCTIONS FOR SPRINKLER (OVERHEAD) IRRIGATION:

Observe the requirements in the System Requirements section above.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply MANEB 75DF only through systems containing anti-siphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

MANEB 75DF may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

Check local restrictions and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

SPRAY PREPARATION

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

Prepare a suspension of MANEB 75DF in a mix tank. Fill the tank with 1/2 or 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of MANEB 75DF and then the remaining volume of water.

APPLICATION INSTRUCTIONS:

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy. Start sprinkler and then uniformly inject the suspension of MANEB 75DF into the irrigation water line so as to deliver the desired rate per acre. The suspension of MANEB 75DF should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

NOTE: When treatment with MANEB 75DF has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage Instructions: Store in the original container in a dry area. If allowed to become wet the product will deteriorate and represent a fire hazard. Keep away from sources of ignition (e.g. sparks and open flame). Close bag when not in use. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Instructions: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

ELI Atochem North America warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. ELI ATOCHEM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL ELI ATOCHEM OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS, LARDER COST, OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

ELI Atochem and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of ELI Atochem.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN OR INHALED. Avoid contact with skin or clothing. Avoid breathing spray mist or dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

Do not apply directly to water; except as specified for the labeled use on cranberries to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

GENERAL INFORMATION

MANEB 75DF may be used as a spray for the control of many fungus plant diseases. It mixes readily with water, suspends well, and adheres to sprayed surfaces with a minimum of runoff.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WORKER SAFETY RULES

Keep all unprotected persons, children, livestock, and pets away from treated area or where there is danger of drift. Do not rub eyes or mouth with hands. See Statement of Practical Treatment.

PERSONAL PROTECTIVE EQUIPMENT

ALL AGRICULTURAL WORKERS/HANDLERS (MIXERS, LOADERS, AND APPLICATORS) APPLYING EBDC'S AND EARLY REENTRY WORKERS MUST WEAR THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT: coveralls over long-sleeved shirt and long pants, shoes, socks, and chemical-resistant gloves. During mixing and loading, a chemical-resistant apron and goggles or a face shield must also be worn. For agricultural workers, where completely enclosed cabs with positive pressure filtration or an enclosed cockpit for aerial application are used, a long-sleeved shirt and long pants may be worn in place of the protective clothing described above. Chemical-resistant gloves must be available in the cab or cockpit and worn upon exiting. The gloves must be kept in an enclosed container in the cab or cockpit to prevent contamination of the inside of the cab or cockpit.

During aerial application, human flaggers are prohibited unless in totally enclosed vehicles.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before eating, smoking or drinking. Always wash hands and arms with soap and water before using the toilet. After work take off all clothes and shoes. Shower using soap and water. Wear only clean clothes. Do not use contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment drenched with maneb must be destroyed according to state and local regulations.

DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

REENTRY STATEMENT

Do not enter treated areas within 24 hours after application unless wearing the protective clothing listed on this label. Because certain states may require more restrictive reentry intervals for various crops treated with this product consult your State Department of Agriculture for further information.

After sprays have dried, do not enter or allow entry into treated areas or areas where there is a danger of drift until the 24-hour reentry interval has expired unless wearing the personal protective equipment listed on this label.

Written or oral warnings must be given to workers who are expected to in a treated area or in an area about to be treated with this product. When oral warnings are given, such warnings shall be given in a language customarily understood by the workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by the workers. Written warnings must include the following information: **CAUTION** Area treated with MANEB 75DF fungicide on (date of application). Do not enter without protective clothing within 24 hours after application.

INSTRUCTIONS FOR APPLICATION

MANEB 75DF is recommended for control of the diseases on the crops listed below.

SPRAYS

Place MANEB 75DF into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on product labels prior to spray preparation.

FOR GROUND APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, minimum 10 gallons. Increased volume of water may be necessary as foliage density increases.

To prepare mixture, sift in the required amount of product when the agitator paddles or the jet or recirculating pipe from pump are covered. Fill the tank with water when the product is thoroughly mixed. Three to four level teaspoonfuls of MANEB 75DF per one gallon of water is equivalent to 1 1/2 to 2 pounds per 100 gallons.

FOR AIR APPLICATION use at rate indicated in sufficient water for thorough coverage, a minimum of 3 gallons per acre for field crops and 10 gallons per acre for orchard crops.

When dosage ranges are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate or apply more frequently than the minimum interval given in the directions for that crop.

Where EBDC Products Used Allow The Same Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient (maneb, mancozeb, or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

VEGETABLE CROP USES

CROPS	DISEASES	RATE LBS/ACRE	DIRECTIONS
Beans (Dry)	Anthrachnose Downy Mildew Rust	1.5-2.0	Begin when plants are small. Spray on a 5-7 day interval. Do not apply more than 9.6 lbs. active ingredient (12.8 lbs. product) per acre per season. Do not apply within 30 days of harvest.
Broccoli Brussels Sprouts Cauliflower	Alternaria Leaf Spot Downy Mildew	1.5-2.0	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12.8 lbs. product) per acre per season. Do not apply within 7 days of harvest.
Cabbage Kohlrabi Chinese Cabbage (tight-headed only)	Alternaria Leaf Spot Downy Mildew	1.5-2.0	Plant beds and direct seeded fields. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12.8 lbs. product) per acre per season. Do not apply later than 7 days before harvest.
Chinese Cabbage (loose head)	Alternaria Leaf Spot Downy Mildew	1.0-1.5	Plant beds and direct seeded fields. Spray on a 7-10 day interval. Do not apply more than 7.2 lbs. active ingredient (9.6 lbs. product) per acre per season. Do not apply later than 7 days before harvest in California and 10 days before in Hawaii.
Sweet Corn Popcorn Sweet Corn used for seed production	Common Rust Helminthosporium Blight	1.5	East of the Mississippi (including Arkansas & Louisiana) Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 18 lbs. active ingredient (24.0 lbs. product) per acre per season. Do not feed treated forage to livestock.
		1.5	West of the Mississippi (except Arkansas & Louisiana) Use sufficient water for thorough coverage. Start applications when disease first appears at 3-10 day interval. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 18.0 lbs. active ingredient (24.0 lbs. product) per acre per season. Do not feed treated forage to livestock.
Cucumbers	Alternaria (Macrosporium) Leaf Spot Angular Leaf Spot Anthrachnose Downy Mildew Pythium Fruit Rot	1.5-2.0	Begin when disease threatens or plants begin to run. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (12.8 lbs. product) per acre per season. Do not apply within 5 days of harvest.

VEGETABLE CROP USES *continued*

FRUIT AND NUT CROPS

CROPS	DISEASES	RATE LBS/ACRE	DIRECTIONS
Eggplant	Anthr. necrose Septoria leaf Spot Cladosporium Leaf Mold Early Blight Late Blight Gray Leaf Spot (Stemphylium)	1.5-2.0	Begin at first fruit cluster and repeat at 7-10 day intervals. Do not apply more than 11.2 lbs. active ingredient (14.9 lbs. product) per acre per crop. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot Downy Mildew	1.5-2.0	Begin when disease threatens and apply on a 7-10 day interval. Do not use more than 3.2 lbs. active ingredient (4.2 lbs. product) per cutting. Do not apply within 10 days of harvest.
Lettuce (Head and Leaf) Endive	Downy Mildew	1.5-2.0	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12.8 lbs. product) per acre per season. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming. California—Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient maximum (8.5 lbs. product) per acre per season. Do not apply within 14 days of harvest.
Melons including cantaloupes, casabas, crenshaws, honeydews, watermelons	Alternaria Leaf Spot Anthr. necrose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight	1.0-2.0	Apply as soon as plants begin to run or when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (17.1 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Onions (Dry Bulb)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	2.0-3.0	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 24 lbs. active ingredient (32.0 lbs. product) per acre per season. Do not apply to exposed bulbs.
Onions (Green)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	2.0-3.0	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 16.8 lbs. active ingredient (22.4 lbs. product) per acre per season. Do not apply to exposed bulbs.
(Furrow Drench)	Smut	3.0	Apply as a furrow drench at time of planting onion seeds. Use 75 to 125 gallons of water per acre. Do not use more than 2.4 lbs. active ingredient (3.2 lbs. product) per 29,000 linear feet of row (18 inch spacing). Do not use in California.
Peppers	Anthr. necrose Cercospora Leaf Spot (Frog-eye spot) Phytophthora Blight Ripe Rot	1.5-3.0 1.5-2.0	East of the Mississippi: Begin when disease threatens. Spray on a 7-day interval. Do not apply more than 14.4 lbs. active ingredient (19.2 lbs. product) per acre per season. Do not apply within 7 days of harvest. West of the Mississippi: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12.8 lbs. product) per acre per season. Do not apply within 7 days of harvest.
Potatoes	Early Blight Late Blight	1.5-2.0	Begin when plants are 2 to 6 inches high. Apply on a 5-10 day interval. Do not apply more than 11.2 lbs. active ingredient (14.9 lbs. product) per acre per season. Do not apply within 14 days of harvest except for Connecticut, Florida, Maine, Massachusetts, New Hampshire, New York, Pennsylvania, Vermont, Wisconsin where the PHI is 3 days. It is recommended that this product be used with an Integrated Pest Management Program. Vine kill should occur 14 days before harvest.
Potatoes (Seed Pieces)	Fusarium Seed Piece Decay	1.0 lb. per 10 gal.	Seed piece treatment only. Dip whole or cut tubers. Spread in cool place if held before planting. Do not use treated seed piece for food feed, or oil purposes.
Pumpkins	Angular Leaf Spot Downy Mildew	1.5-2.0	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (17.1 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Squash (Summer and Winter)	Anthr. necrose Downy Mildew	1.5-2.0	Begin when plants start to run. Spray on a 7-10 day interval. Do not apply more than 12.8 lbs. active ingredient (17.1 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Tomatoes (Greenhouse and Field)	Anthr. necrose Septoria Leaf Spot Cladosporium Leaf Mold Early Blight Late Blight Gray Leaf Spot (Stemphylium)	1.5-3.0 1.5-2.0	East of the Mississippi: Begin at first fruit cluster and repeat on 7-10 day intervals. Do not apply within 5 days of harvest. Do not apply more than 16.8 lbs. active ingredient (22.4 lbs. product) per acre per crop. West of the Mississippi: Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient (8.5 lbs. product) per acre per season. Do not apply within 5 days of harvest.

CROPS	DISEASES	RATE LBS/ACRE	DIRECTIONS
FOR PCME FRUITS Use either the "Pre-Bloom" or "Extended Application" schedules. DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES.			
Apples	Rusts (including Cedar apple fabrea leaf Spot Hyspeck Scab Sooty Blotch)	6.0 or (1.5 lbs. per 100 gal. in dilute spray) 3.0 or (0.75 lbs. per 100 gal. in dilute spray)	PRE-BLOOM: Begin applications at 1/4 to 1/2 inch green tip and continue on a 7- to 10-day application schedule through bloom. Do not apply more than 19.2 lbs. active ingredient (25.6 lbs. product) per acre per year. Do not graze livestock in treated areas. EXTENDED APPLICATION OR TANK MIX: Applications based on tree-row volume or for use as a resistance management tool. Begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7- to 10-day schedule through the second cover spray; or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not apply more than 16.8 lbs. active ingredient (22.4 lbs. product) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program. Add spreader/sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/curative fungicide for more effective control of diseases.
Almonds	Brown Rot Blossom and Twig Blight Scab Shot-hole Fungus (Leaf Blight)	6.0-8.0 or (1.5-2.0 lbs. per 100 gal. in a dilute spray applied to run-off)	Apply at popcorn, full bloom, and petal fall or every 7 to 10 days if bloom is staggered. Omit petal fall spray if only brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not apply more than 25.6 lbs. active ingredient (34.1 lbs. product) per acre per season. Do not allow livestock to graze in almond orchards. Adjust the spray concentration in concentrate spray for different sized trees.
Bananas	Sigatoka	2.0-3.0	Apply when leaves first appear and repeat as needed on a 14-21 day interval. Do not apply more than 24 lbs. active ingredient (32 lbs. product) per acre per season. May be applied up to the day of harvest.
Cranberries	Fruit Rot	4.8-6.0	Spray on a 14-21 day interval. Do not apply more than 14.4 lbs. active ingredient (19.2 lbs. product) per season. Do not apply within 30 days of harvest.
Grapes	Black Rot Bunch Rot Deadarm Downy Mildew	1.5-4.0 or (0.375-1.0 lbs. per 100 gal. in a dilute spray) 1.5-2.5 or (0.375- 0.63 lbs. per 100 gal. in a dilute spray)	East of the Rocky Mountains: Apply in sufficient water to provide thorough coverage starting when shoots are 1/2 to 1 1/2 inches long and continue at 7-10 day intervals until fruit is set. Do not apply within 66 days of harvest. Do not apply more than 19.2 lbs. active ingredient (25.6 lbs. product) per acre per season. West of the Rocky Mountains: Apply when shoots are 1/2 to 1 1/2 inches long and continue at 7-10 day intervals. Do not apply more than 6.0 lbs. active ingredient (8.0 lbs. product) per acre per season. Do not apply within 66 days of harvest except in California. California—Do not apply after bloom.
Kadota figs	Surface Mold (Cladosporium) Surface Rot (Alternaria)	0.75 lbs./ 100 gal.	Apply when disease threatens in 100-400 gallons of water/acre. Do not apply more than 2.4 lbs. active ingredient (3.2 lbs. product) per acre per season. Do not apply within 10 days of harvest.
Papayas	Anthr. necrose Phytophthora Fruit Rot	2.0-2.5 or (0.5- 0.625 lbs. per 100 gal. in a dilute spray)	Apply when disease first threatens and 14-21 day intervals. Do not apply more than 28 lbs. active ingredient per season (37.0 lbs. product). May be applied up to the day of harvest.

FIELD CROPS

CROPS	DISEASES	RATE LBS/ACRE	DIRECTIONS
Corn	Leaf Blight Gray Leaf Spot (Stemphylium)	1.5-2.0	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 11.2 lbs. active ingredient (14.9 lbs. product) per acre per season. Do not apply within 14 days of harvest.