

4581-373

4-11-2007

112



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Timothy M. Formella
Manager, Product Registration
Cerexagri, Inc.
630 Freedom Business Center, Suite402
King of Prussia, PA 19406

APR 11 2007

Dear Mr. Formella:

Subject: Microthiol Disperss
EPA Reg. No.4581-373
Your Submission of January 24, 2007

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable. A stamped copy of the label is enclosed for your records. Submit one copy of the final printed label before you release the product for shipment.

Sincerely yours,

A handwritten signature in cursive script, appearing to read "Mary L. Waller".

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure

7505P:CGrable:cg:4/11/07

2/12

ACCEPTED
with COMMENTS
In EPA Letter Dated

APR 11 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

4581-373

MICROTHIOL® DISPERS®

MICRONIZED WETTABLE SULFUR

For Organic Production

ACTIVE INGREDIENT:

| | |
|-------------------------|--------|
| Sulfur | 80.0% |
| OTHER INGREDIENTS | 20.0% |
| TOTAL | 100.0% |

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

EPA Reg. No. 4581-373

EPA Est. No. 44420-FR-1

Net Contents: _____

Cerexagri, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)
CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling, and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- shoes plus socks
- chemical resistant gloves made of any waterproof material
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For non-agricultural turf and grass uses (golf courses, home lawns, and landscapes around buildings) do not enter or allow entry into treated area until sprays have dried.

APPLICATION INSTRUCTIONS

Apply through any type of handheld, knapsack, mechanical or pneumatic spraying equipment.

MIXING: Pour the required amount of MICROTHIOL DISPERSS into the indicated amount of water while stirring. MICROTHIOL DISPERSS disperses instantly.

Dilute Application:

Field: Specified rate in 20 to 60 gallons of water per acre.

Orchard: Specified rate in 100 to 800 gallons of water per acre.

Concentrated Application:

Field: Specified rate in 5 to 10 gallons of water per acre.

Orchard: Specified rate in 20 to 100 gallons of water per acre.

Aerial Application:

Field: Specified rate in 3 to 20 gallons of water per acre. Orchard: Specified rate in 10 to 20 gallons of water per acre.

COMPATIBILITY

MICROTHIOL DISPERSS is compatible with most Bordeaux mixtures, copper fungicides, liquid fertilizers, organic fungicides, insecticides, acaricides and herbicides.

USE INSTRUCTIONS

Some crops may be damaged by sulfur under certain climatic conditions, such as when temperature is high. Cerexagri does not recommend application if temperature will exceed 90° F within the three days following spraying, due to the risk of crop injury. Crops grown in greenhouses may be more sensitive to sulfur injury, so the lowest labeled rate (or below labeled rates) should be tried initially. Do not use on any crop unless sulfur has been shown to be safe in your locality. Do not allow spray to drift onto sulfur-sensitive crops such as apricots, cranberries, spinach and d'Anjou pears.

Do not use within two weeks of an oil spray treatment, except for Dormant, Delayed Dormant, or Postharvest applications with a suitable oil. For citrus, do not apply within 21 days of an oil spray. Spreader/stickers are not required for use with MICROTHIOL DISPERSS. If spreader/stickers are used with MICROTHIOL DISPERSS, foliar injury may be enhanced under high temperatures. When growing crops for processing, consult the processor before applying sulfur.

NOTE: The application rates specified for each crop are the maximum that may be applied. These amounts may be reduced if experience or local practice indicates that lesser amounts may be effective. Consult State Agriculture Experiment Station or State Extension Service specialist for rates recommended in your local area.

VEGETABLE CROPS

| Crop | Pest | Rate Lbs/A | Directions |
|-------------------------------|-----------|---------------|--|
| Artichokes (all varieties) | Leaf Spot | 10-20 | Begin when disease first appears and repeat at 7- to 10-day intervals or as needed. |
| Asparagus | Rust | 10-30 | Apply after cutting, irrigate and cultivate before applying. Repeat at 7- to 10-day intervals, or as needed, through fern growth season. |

| Crop | Pest | Rate Lbs/A | Directions |
|--|--|---------------|---|
| Beans, Peas | Leaf Spot Powdery Mildew Red Spider Mite Rust | 3 - 10 | Apply at early leaf stage and repeat at 10- to 14-day intervals or as needed. |
| Broccoli, Brussels sprouts, Cabbage, Cauliflower Kohlrabi | Powdery Mildew Red Spider Mite | 3-10 | Begin in early leaf stage. Repeat every 10 to 14 days or as needed. |
| Carrots | Petrobia Mite Powdery Mildew | 3-10 | Apply at early leaf stage and repeat every 14 days or as needed. |
| Celery | Powdery Mildew Red Spider Mite Rust | 4-6 | Apply at early leaf stage and repeat every 10 to 14 days or as needed. |
| Cowpeas | Rust | 2-9 | Begin applications soon after seedlings emerge. Repeat at 7- to 10-day intervals as needed throughout the season. |
| Cucumbers | Powdery Mildew | 2-4 | Apply when disease first appears and repeat as necessary. Do not apply more than 2 pounds MICROTHIOL DISPERSS per acre per application during periods of high temperatures. |
| Eggplant | Powdery Mildew | 4-6 | Apply when first true leaves appear. Repeat at weekly intervals or as needed. |
| Endive Escarole Swiss Chard Salsify | Powdery Mildew Red Spider Mite Rust | 4-6 | Apply at early leaf stage and repeat every 10 to 14 days or as needed. |
| Greens: Collards, Kale, Mustard, Beet, Turnip | Powdery Mildew Red Spider Mite Rust | 3-10 | Apply at early leaf stage and repeat every 14 days or as needed |
| Lettuce (Head and Leaf) | Powdery Mildew Red Spider Mite Rust | 5-10 | Apply at early leaf stage and repeat every 14 days or as needed. Thorough coverage is required. |
| Melons | Powdery Mildew | 5-10 | Apply when disease first appears and repeat as needed. Sulfur can injure plants, especially when temperature reaches 90° F. Do not use on sulfur sensitive varieties. |
| Okra | Leaf Spot Mites Powdery Mildew | 3-10 | Apply at early leaf stage and repeat every 14 days or as needed. |
| Onions, Garlic, Dry Onions Dry Shallots | Petrobia Mite Powdery Mildew | 3-10 | Apply when disease first appears and repeat as needed. |
| Potatoes Sweet Potatoes | Leaf Spot Powdery Mildew Red Spider Mite | 5 | Apply at early leaf stage and repeat every 14 days or as needed. |
| Peppers | Broad Mite | 3-10 | Apply at early leaf stage and repeat every 14 days or as |

6/12

| Crop | Pest | Rate Lbs/A | Directions |
|---|---|---------------|---|
| | Leaf Spot Powdery Mildew Red Spider Mite | | needed. |
| Spinach | Powdery Mildew Rust | 4-6 | For use on spinach to be processed only. Do not use on fresh spinach as sulfur has been shown to have phytotoxic effects on certain varieties of spinach. Spray on a small area first to test for phytotoxicity before making a full application. Apply when disease first appears and repeat at 7- to 10-day intervals or as needed. |
| Squash (including Pumpkins, Summer and Winter Squash) | Powdery Mildew | 5-10 | Apply when disease first appears and repeat as needed. Sulfur can injure plants, especially when temperature reaches 90° F. Do not use on sulfur sensitive varieties. |
| Table Beets | Powdery Mildew Red Spider Mite | 3-5 | Apply at first indication of problem with or without other pesticides. Repeat as necessary. |
| Tomatoes | Powdery Mildew Russet Mite Two-Spotted Mite | 5-10 | Begin application when symptoms first appear and repeat at 7- to 10-day intervals or as needed. Thorough coverage is required. |
| Turnips Parsnips Horseradish Radishes Rutabagas | Powdery Mildew Red Spider Mite Rust | 3-10 | Apply at early leaf stage and repeat every 14 days or as needed. |

FRUIT AND NUT CROPS

| Crop | Pest | Rate Lbs/A | Directions |
|---|--|---------------|---|
| Almonds | Brown Rot Flat Mite Leaf Spot Red Spider Mite Rust Silver Mite | 10-20 | Apply at dormant through harvest, alone or in tank-mixes with other fungicides/miticides for disease protection, mite suppression, and resistance management. Repeat applications at 7- to 10-day intervals or as needed. |
| Apples, Pears | Blister Mite Powdery Mildew Red Spider Mite Scab Silver Mite Two-Spotted Mite | 10-20 | Apply at prebloom and petal fall. Apply with cover sprays throughout the season. Do not apply to d'Anjou pears. |
| Apples, Pears (Dormant, Delayed Dormant or | Blister Mite Pear Psylla Red Spider Mite Two-Spotted | 10-20 | Apply with a suitable spray oil postharvest and during the dormant periods of bud development. Do not apply to sensitive varieties of pears except at these stages. |

7/12

| Crop | Pest | Rate Lbs/A | Directions |
|--------------|---|---------------|---|
| Postharvest) | Mite | | |
| Avocados | Brown Mite | 10-20 | Apply as needed. |
| Bananas | Mites | 10-30 | Apply as needed. |
| Citrus | Broad Mite Bud Mite Citrus Rust Mite Flat Mite Red Spider Mite Silver Mite Six-spotted Mite Thrips Yuma Spider Mite | 10-15 | Apply as needed. Do not apply within 21 days of an oil spray. |
| Dates | Banks Grass Mite | 20 | Apply in sufficient water to provide thorough coverage of the date clusters. Begin applications as mites or mite webbing are first observed. Repeat applications every 14 days or as necessary, until dates are mature and are no longer subject to mite feeding injury. |
| Figs | Mites, including Eriophyid Mites, Fig Mite | 5 | Apply at a minimum of 100 gallons per acre. Begin applications in the spring, and continue on a 3-4 week schedule or as necessary. |
| Grapes | Blister Mite Bud Mite Phomopsis Powdery Mildew Red Spider Mite | 3-10 | Apply throughout the season at 7- to 14-day intervals or as needed. Thorough coverage is required. Note: Use with caution on Concord, Labrusca type, and other sulfur sensitive grape cultivars as injury can occur after bloom. |
| Mangos | Powdery Mildew | 10-30 | Begin application before bloom. Repeat at bloom, after fruit set, and 3 weeks later or as necessary. |
| Olives | Olive Mites | 10-30 | Apply as needed. |
| Papaya | Mites | 10-30 | Apply when mites first appear and repeat as necessary to maintain control. Do not apply in combination with emulsifiable pesticide formulations as foliar injury may occur. Do not apply to low vigor or moisture stressed plants. Do not apply prior to or during periods of excessively high temperatures, periods of low rainfall or within two weeks of an oil spray. |
| Pecans | Eriophyid Mites (including Hickory and Pecan) Flat Mite Leaf Spot Powdery Mildew Red Spider Mite Silver Mite Sooty Mold Two-Spotted Mite | 5-10 | Apply throughout the season as needed. Note: Some varieties of pecans are sensitive to sulfur sprays under certain conditions. Do not apply unless varieties are known to be tolerant of sulfur. |

| Crop | Pest | Rate Lbs/A | Directions |
|--|---|---------------|---|
| Pineapple | Mites | 10-30 | Apply as needed. |
| Pistachios | Mites | 10-20 | Begin applications in June-July. Make additional applications as necessary. When temperatures exceed 90° F, lower rates and more frequent applications are advised in order to avoid crop injury. |
| Pomegranate | Mites | 3-10 | Begin applications in May or June. Make additional applications on a 3-4 week schedule, or as necessary. Use higher rates if past mite damage has been high. If temperatures exceed 90° F, lower rates and more frequent applications are advised in order to avoid crop injury. |
| Quince | Brown Rot Powdery Mildew Scab | 10-30 | Begin applications at early leaf stage and repeat as necessary. Do not use on sulfur sensitive varieties. |
| Stone Fruit: Peaches, Plums, Cherries, Nectarines, Prunes | Brown Rot Flat Mite Leaf Spot Powdery Mildew Red Spider Mite Rust Silver Mite | 10-20 | Apply at dormant through harvest, alone or in tank-mixes with other fungicides/miticides for disease protection, mite suppression, and resistance management. Repeat applications at 7- to 10-day intervals or as needed. For Brown Rot of fruit, apply when fruit starts to ripen. Do not apply to apricots. |
| Walnuts, Macadamias | Almond Mite Broad Mite Pacific Mite Red Spider Mite Two-Spotted Mite | 10-20 | Apply throughout the season as needed. Note: Some varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless varieties are known to be tolerant of sulfur. |

FIELD CROPS

| Crop | Pest | Rate Lbs/A | Directions |
|--|--|---------------|---|
| Alfalfa Including: Seed Alfalfa | Atlantic Mite Red Spider Mite | 8-10 | Apply throughout the season as necessary. |
| Cereals: Corn, Wheat, Barley, Oats, Rye, Sorghum | Atlantic Mite Banks Grass Mite Pacific Mite Red Spider Mite Two-spotted mite | 8-10 | Apply when mites or symptoms first appear and repeat applications as necessary. |
| Clover | Mites Powdery Mildew | 2-9 | Apply as needed. |
| Cool Season Forage Grasses (including; Timothy, Orchard Grass, | Mites (including Banks Grass Mite) | 5-10 | Apply at first sign of mites. Continue applications at 7-day intervals, or as needed, until mites are below economic levels. The addition of adjuvants (spreaders, surfactants, etc.) can increase the chance of crop injury, and is not recommended. |

9/12

| Crop | Pest | Rate Lbs/A | Directions |
|---|--|---------------|---|
| Fescues, Sudan Grass, Bermuda Grass, Bluegrass, and Bromegrass) | | | For aerial applications: Use the specified rate in a minimum of 3 gallons of water per acre. For ground applications: Use the specified rate in a minimum of 20 gallons of water per acre. |
| Cotton | Atlantic Mite Red Spider Mite Two-Spotted Mite | 5-10 | Apply throughout the season as necessary. |
| Flax | Powdery Mildew | 2-9 | Apply at first sign of disease. Repeat at 7- to 10-day intervals or as necessary. |
| Peanuts | Leaf Spot Powdery Mildew Red Spider Mite Rust Two-Spotted Mite | 5-8 | Apply at early leaf stage and repeat applications as necessary. |
| Soybeans | Leaf Spot Powdery Mildew Rust | 10-15 | Apply at early leaf stage and repeat at 7- to 14-day intervals or as needed. |
| Sugar Beets | Powdery Mildew Red Spider Mite | 5-10 | Apply as soon as symptoms appear and repeat every 7 to 14 days or as needed. |
| Sugarcane | Mites | 8-15 | Apply when mite pressure is imminent and repeat applications as necessary. |

SMALL FRUITS AND BERRIES

| Crop | Pest | Rate Lbs/A | Directions |
|--|---|---------------|---|
| Blackberry, Boysenberry, Dewberry, Loganberry, Blueberry, Gooseberry, Huckleberry, Currant | Powdery Mildew | 6-15 | Begin application before blossoms open and continue at 10-day intervals or as needed. |
| Raspberries | Powdery Mildew Red Berry Mite | 6-15 | Apply each week from first bloom to fruit set. Repeat as the weather requires. |
| Strawberries | Powdery Mildew Red Spider Mite Two-Spotted Mite | 5-10 | Begin application at early leaf stage and repeat as needed. Do not use on sulfur-sensitive varieties. |

MISCELLANEOUS CROPS

| Crop | Pest | Rate | Directions |
|------|------|------|------------|
|------|------|------|------------|

| | | Lbs/A | |
|--------------------------|---|-------------------------------------|--|
| Spearmint, Peppermint | Powdery Mildew | 4-6 | Apply when mint is 5-6" tall or when disease appears. Repeat twice, at 30-day intervals. Do not apply within 30 days of harvest. |
| Hops | Red Spider Mite | 2-6 | Begin applications at early leaf stage, and repeat as needed. |
| Ornamentals, Roses | Leaf Spot Powdery Mildew Two-Spotted Mite | 3-10 by ground or 3-20 by air | Apply to ensure complete plant coverage. Begin when disease first appears and repeat at 5-10 day intervals. |

TURF AND GRASS

| Crop | Pest | Rate Lbs/A | Directions |
|---|--|---------------|---|
| Turf and Grass (Commercial, including; golf courses, sod farms, and grass grown for seed) | Fusarium Patch (suppression) Powdery Mildew Rust Take-all (suppression) | 3-10 | Applications should begin at the very first sign of disease, or as a preventative when conditions favor disease. These are explosive diseases and are very difficult to control once fully established. Repeat applications as needed. Tank mixing with systemic fungicides will improve control and help in a resistance management program with these fungicides. |

DIRECTIONS FOR USE THROUGH CHEMIGATION SYSTEMS

GENERAL INSTRUCTIONS

Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, hand move; or drip (mini-micro sprinklers, strip tubing, trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

PRECAUTION: Corrosion of aluminum and carbon/galvanized steel irrigation sprinkler systems may be experienced with the use of sulfur fungicides. If the user elects to apply this product through such systems, it is essential that all application equipment containing this product be thoroughly flushed with clean water after each day's use. Continue to operate system with clean water until all product has cleared the last sprinkler head.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

MICROTHIOL DISPERSS may be applied in conjunction with chemically neutral liquid fertilizers.

SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

APPLICATION INSTRUCTIONS

Observe the requirements in the System Requirements section.

Apply MICROTHIOL DISPERSS only through systems containing anti-siphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

SPRAY PREPARATION:

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water. Prepare a suspension of MICROTHIOL DISPERSS in a mix tank. Fill the tank with 1/2 or 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of MICROTHIOL DISPERSS and then the remaining volume of water.

Sprinkler Irrigation- Notes:

Observe all System Requirements and Application Instructions above.

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy. Start sprinkler and then uniformly inject the suspension of MICROTHIOL DISPERSS into the irrigation water line so as to deliver the desired rate per acre. The suspension of MICROTHIOL DISPERSS should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. When treatment with MICROTHIOL DISPERSS has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

Check local restrictions and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a closed package in a cool, dry, secure place. Do not store near flammable materials. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then dispose of bag in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains or other influencing factors in the use of the product. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Cerexagri, Inc., Manufacturer and Seller harmless for any claims relating to such factors. Cerexagri, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CEREXAGRI, INC. AND MANUFACTURER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, in no event shall Cerexagri, Inc., Manufacturer or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CEREXAGRI, INC., MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CEREXAGRI, INC., MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Cerexagri, Inc., Manufacturer and Seller offer this product, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Cerexagri, Inc.

Microthiol and Disperss are registered trademarks of Cerexagri, Inc.

Rev. 01/24/07