.4581 -	388
---------	-----

3-23-2001

	-388		-23-200						Y	
Please read instructions on I	Ur Environmental	United States nental Protection Age Washington, DC 20460		orm Approv	Registra	Registration Amendment		OPP Identifier Number		
		Application	n for Pesticide	Section	n l					
1. Company/Product Numbe 4581-388	1. Company/Product Number			2. EPA Product Manager Miller				3. Proposed Classification		
4. Company/Product (Name) Aquathol Super K)		PM# 23				None	nes	stricted	
5. Name and Address of Ap Cerexagri, Inc. 2000 Market Street, 2 Philadelphia, PA 19	21st Floor	de)	(b)(i), my pi to: EPA Reg.	roduct is si	. In accorda	tical in co	ımpositio			
			Section - II	taine					===	
Amendment - Explain Resubmission in resp Notification - Explain Explanation: Use addition Notification amendment to: sites that can be irrigated im I certify that this notification USC Sec. 1001 to willfully mad CFR 152.46, this product	below. hal page(s) if necessary Add "Other" instead of mediately after treatmen is consistent with the proake any false statement	r. (For section "Inert" Ingredier t. ovisions of PR N to EPA. I furthe	Age "M Oth I and Section II.) hts, 2) Add "First Aid" ins Notice 98-10 and EPA re or understand that if this	ency tetter d e Too" Apple er - Explain stead of "State gulations at a	below. tement of Pract 40 CFR 152.46	ical Treatm	MAF	s a violatio Notice 98-	ence to	
1. Material This Product Wil	1 D - D - d 1 1		Section - III			·				
Child-Resistant Packaging Yes No Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container		ging Vo. per container	2. Type of	Metal Plastic Glass Paper	Specify)_			
3. Location of Net Contents	Information	4. Size(s) Reta	ail Container	5.1	Location of La	bel Directi	០ពន			
Label 6. Manner in Which Label is	Container Affixed to Product	Lithogr Paper Stencii	aph plued ed	Other						
			Section - IV							
1. Contact Point (Complete	items directly below for	or identific etio i	of individual to be co	ntacted, if n	ecessary, to p	rocess this	application	on.)	_	
Name Christopher Davis			Title Registration Manage	г		Telephor 215-419		lude Area	Code)	
•	ments I have made on ny knowlingly false or l law.		all attachments thereto			•	Recai			
2. Signatore) _~		3. Title Registration Manager				, , , , , , ,	,))))))	
4. Typed Narhe Christopher Davis		5	2/28/0)					, , ,		

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

White - EPA File Copy (original)

AQUATHOL° SUPER K GRANULAR AQUATIC HERBICIDE

ACTIVE INGREDIENT:

NOTIFICATION

MAR 2 3 2001

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID

IF IN EYES, hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED, Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth or do not induce vomiting.

IF ON SKIN, Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See side panel for additional Precautionary Statements.

EPA Registration No. 4581-388

EPA Establishment No. 39578-TX-1

Cerexagri, Inc. 2000 Market Street Philadelphia, PA 19103 Net Weight: 25 lbs.

^{* 7-}oxabicyclo [2.2.1]heptane-2,3-dicarboxylic acid equivalent 44.7%

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS) DANGER

CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE. MAY BE FATAL IF SWALLOWED. HARMFUL IF ABSORBED THROUGH THE SKIN. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles or face shield). Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

Avoid contact with or drift to other crops or plants as injury may result.

Do not use water from treated areas for irrigation, for agricultural sprays on food crops or for domestic purposes within 7 days of treatment.

Do not use fish from treated areas for food or feed within 3 days of treatment.

GENERAL INFORMATION

AQUATHOL SUPER K is a granular aquatic herbicide for use in ponds and lakes which, under field test conditions has shown to be effective against a broad range of aquatic plants with a margin of safety to fish. Dosage rates indicated for the applications of AQUATHOL SUPER K are measured in "Parts Per Million" (ppm). 1 ppm as a dosage rate means that there would be 1 part of AQUATHOL SUPER K 's active ingredient in 1,000,000 parts of water. Only 0.5 to 5 ppm are generally required for aquatic weed control, whereas some fish species are tolerant to approximately 100 ppm or over. For best results treat areas of one acre or more and/or margins of at least 100 feet in large bodies of water.

Thoroughly clean application equipment immediately after use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY:

AQUATHOL SUPER K is a contact herbicide, consequently, do not apply before weeds are present. Treat as early as possible after weeds are present and are actively growing. If an entire pond is treated at one time, or if the dissolved oxygen level is low, at the time of application, decay of weeds may remove enough oxygen from the water causing fish to suffocate. Water containing very heavy vegetation should be treated in sections to prevent suffocation of fish. Sections should be treated 5-7 days apart.

Carefully measure size and depth of area to be treated and determine proper amount of

AQUATHOL SUPER K to apply. For best results apply when water is quiescent and/or flows are minimal.

Apply AQUATHOL SUPER K as evenly as possible over treated areas. A cyclone spreader is useful for this purpose. In instances where the nuisance weed(s) to be controlled is an exposed surface problem (i.e., some of the broad-leaved pond weeds) coverage is important.

Necessary approval and/or permits should be obtained in states where required.

HOW TO DETERMINE DOSAGE RATE (Active Ingredient)

AQUATHOL SUPER K is recommended for the control of the following aquatic weeds at the rates indicated. Since AQUATHOL SUPER K 's active ingredient is water soluble and tends to diffuse from the treated area, select the dosage rate applicable to the area to be treated. Use the lower rate in each range when the growth is young and growing and/or where the weed stand is not heavy. Marginal treatments of large bodies of water require highest rates as indicated.

WEEDS CONTROLLED AND AQUATHOL SUPER K DOSAGE RATE CHART

	RATES					
Aquatic Weed	Entire Pond/Lake or Large Area Treatment	Lbs. per Acre Ft.	Spot or Lake Margin Treatment	Lbs. per Acre Ft.		
Bur Reed, Sparganium spp.	3.0-4.0 ppm	13.2-17.6 lbs.	4.0-5.0 ppm	17.6-22.0 lbs.		
Coontail, Ceratophyllum spp.	1.0-2.0 ppm	4.4-8.8 lbs.	2.0-3.0 ppm	8.8-13.2 lbs.		
Horned Pondweed, Zannichellia palustris	1.0-2.0 ppm	4.4-8.8 lbs.	2.0-3.0 ppm	8.8-13.2 lbs.		
Hydrilla, Hydrilla verticillata	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.		
Hygrophila, Hygrophila polysperma	4.0-5.0 ppm	17.6-22.0 lbs.	5.0 ppm	22.0 lbs.		
Milfoil, Myriophyllum spp.	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.		
Naiad, Najas spp.	1.0-3.0 ppm	4.4-13.2 lbs.	2.0-4.0 ppm	8.8-17.6 lbs.		
Pondweed, Potamogeton spp. Including:	0.5-3.0 ppm	2.2-13.2 lbs.	1.5-4.0 ppm	6.6-17.6 lbs.		
American, P. nodosus	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6,lb\$.		
Largeleaf (Bass Weed), P. Amplifolius	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.		
Curlyleaf, P. crispus	0.5-1.5 ppm	2.2-6.6 lbs.	1.5-3.0 ppm	6.6-13:2 lbs.		

Flatstem, P. zosterformis	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.
Floating-leaf, P. natans	1.0-2.0 ppm	4.4-8.8 lbs.	2.0-3.0 ppm	8.8-13.2 lbs.
Illinois, P. Illinoensis	1.5-2.5 ppm	6.6-11.0 lbs.	2.5-3.5 ppm	11.0-15.4 lbs.
Narrowleaf, P. pusillus	1.0-2.0 ppm	4.4-8.8 lbs.	2.0-3.0 ppm	8.8-13.2 lbs.
Slender, P. filiformis	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.
Sago, P. pectinatus	1.0-2.0 ppm	4.4-8.8 lbs.	2.0-3.0 ppm	8.8-13.2 lbs.
Variable Leaf, P. diversifolius	1.0-2.0 ppm	4.4-8.8 lbs.	2.0-3.0 ppm	8.8-13.2 lbs.
Parrot Feather, Myriophyllum aquaticum	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.
Water Stargrass, Heteranthera spp.	2.0-3.0 ppm	8.8-13.2 lbs.	3.0-4.0 ppm	13.2-17.6 lbs.

Approximate Pounds of Aquathol Super K For One Acre*
Treatment Dosage in Pounds For Various Concentrations In PPM

Treatment besage in Founds For Various confermations in Frim								
Depth	0.5 ppm	1.0 ppm	1.5 ppm	2 ppm	3 ppm	4 ppm	5 ppm	
1 Ft. Deep	2.2	4.4	6.6	8.8	13.2	17.6	22	
2 Ft. Deep	4.4	8.8	13.2	17.6	26.4	35.2	44	
3 Ft. Deep	6.6	13.2	19.8	26.4	39.6	52.8	66	
4 Ft. Deep	8.8	17.6	26.4	35.2	52.8	70.4	88	
5 Ft. Deep	11	22	33	44	66	88	110	

^{*} One acre equals approximately 208' x 208'

Where the area being treated is greater than those listed in the charts, proceed as follows:

- a. Compute the approximate surface acreage
- b. Determine the average depth
- c. Multiply a. by b. to determine total number of acre/feet
- d. Multiply the pounds required at the 1 foot depth under the rate to be used by the number of acre/feet to determine the total quantity to be used.

Approximate Pounds of Aquathol Super K For 1000 Square Feet Treatment Dosage in Pounds For Various Concentrations In PPM

Depth	0.5 ppm	1.0 ppm	1.5 ppm	2 ppm	3 ppm	4 ppm	5 ppm
1 Ft. Deep	0.05	0.1	0.15	0.2	0.3	0.4	0.5
2 Ft. Deep	0.10	0.2	0.3	0.4	0.6	0.8	1.0
3 Ft. Deep	0.15	0.3	0.45	0.6	0.9	1.2	1.5
4 Ft. Deep	0.2	0.4	0.6	0.8	1.2	1.6	2.0
5 Ft. Deep	0.25	0.5	0.75	1.0	1.5	2.0	2.5

Where the depth is greater than 5 feet, multiply the depth by the appropriate rate for 1 ft depth to determine the amount of product required per 1000 square feet.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage Instructions: Store in the original container, preferably in a locked storage area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal Instructions: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

Cerexagri, Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. CEREXAGRI MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL CEREXAGRI OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COSTS; OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING. Cerexagri and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Cerexagri.