

11776
JAN 23 197
FIS

Dr. MAYFIELD

WARFARIN CONCENTRATE FOR KILLING RATS AND MICE 1/4 Pound

E. P. A. Registration Number 4644

Active Ingredients: WARFARIN (3-(α -acetylbenzyl) - 4-hydroxycoumarin).....	0.5%
Inert Ingredients	99.5%
Total	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Right Panel for Additional Cautions
Manufactured for

DR. MAYFIELD LABORATORIES
CHARLES CITY, IOWA

DIRECTIONS FOR RATS: Mix thoroughly one part Dr. Mayfield WARFARIN CONCENTRATE with 19 parts, by weight, of a suitable bait material, such as fresh corn meal, rolled oats, nut crumbs, poultry mash, etc. If very dusty, add up to 5% of a bland, edible oil, such as mineral or corn oil. THIS 1/4 LB. PACKAGE MAKES 5 LBS. OF MIXED BAIT.

Four ounces of bait per bait location should be minimum amount placed.

An uninterrupted supply of bait should be maintained for approximately 10 days.

When perishable baits, such as ground meat, fish, vegetables, etc. are used, place bait in late afternoon and replace daily. Dogs and cats will actively search out the baits of this type. Therefore, baits must be placed so as to positively prevent feeding by these animals.

Protect from children, pets, and livestock by means of cages or bait boxes where necessary.

Replace baits as consumed. Never use moldy, sour or decomposed baits. Never let a bait station remain empty.

Near dumps and other sources of continual infestation, use permanent bait stations, and replenish as needed.

PA 12573

DIRECTIONS FOR follow the same g but use smaller ba are more difficult complete control n

Place table spoon foot intervals.

An uninterrupted maintained for a

Do not reuse con

CAUTION: WARFA ingredient an ant if taken by human may reduce the cl and serious hemor give a tablespoonf water and repeat u physician immediat

NOTE FOR PHYSIC been known to blood transfusions infection and are indicated a in the a. under st D

1116
JAN 23 197

Dr. MAYFIELD

WARFARIN CONCENTRATE FOR KILLING RATS AND MICE 1/4 Pound

E. P. A. Registration Number 4644-1

Active Ingredients: WARFARIN (3-(a-acetylbenzyl) - 4-hydroxycoumarin).....	0.5%
Inert Ingredients	99.5%
Total	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Right Panel for Additional Cautions
Manufactured for

DR. MAYFIELD LABORATORIES
CHARLES CITY, IOWA

DIRECTIONS FOR RATS: Mix thoroughly one part Dr. Mayfield WARFARIN CONCENTRATE with 19 parts, by weight, of a suitable bait material, such as fresh corn meal, rolled oats, nut crumbs, poultry mash, etc. If very dusty, add up to 5% of a bland, edible oil, such as mineral or corn oil. THIS 1/4 LB. PACKAGE MAKES 5 LBS. OF MIXED BAIT.

Four ounces of bait per bait locations should be minimum amount placed.

An uninterrupted supply of bait should be maintained for approximately 10 days.

When perishable baits, such as ground meat, fish, vegetables, etc. are used, place bait in late afternoon and replace daily. Dogs and cats will actively search out the baits of this type. Therefore, baits must be placed so as to positively prevent feeding by these animals.

Protect from children, pets, and livestock by means of cages or bait boxes where necessary.

Replace baits as consumed. Never use moldy, sour or decomposed baits. Never let a bait station remain empty.

Near dumps and other sources of continual intestation, use permanent bait stations and replenish as needed.

PA 12573

DIRECTIONS FOR follow the same ; but use smaller b are more difficu complete control i

Place tablespo foot intervals.

An uninterupte maintained for a

Do not reuse co

CAUTION: WARF ingredient an an if taken by human may reduce the c and serious hemo give a tablespoof water and repeat u physician immedic

NOTE FOR PHYSI been known to blood transfusions injections and are indicated as in the by overdoses of D

DIRECTIONS FOR RATS: Mix thoroughly one part Dr. Mayfield WARFARIN CONCENTRATE with 19 parts, by weight, of a suitable bait material, such as fresh corn meal, rolled oats, nut crumbs, poultry mash, etc. If very dusty, add up to 5% of a bland, edible oil, such as mineral or corn oil. THIS 1/4 LB. PACKAGE MAKES 5 LBS. OF MIXED BAIT.

Four ounces of bait per bait locations should be minimum amount placed.

An uninterrupted supply of bait should be maintained for approximately 10 days.

When perishable baits, such as ground meat, fish, vegetables, etc. are used, place bait in late afternoon and replace daily. Dogs and cats will actively search out the baits of this type. Therefore, baits must be placed so as to positively prevent feeding by these animals.

Protect from children, pets, and livestock by means of cages or bait boxes where necessary.

Replace baits as consumed. Never use moldy, sour or decomposed baits. Never let a bait station remain empty.

Near dumps and other sources of continual infestation, use permanent bait stations and replenish as needed.

PA 12573

DIRECTIONS FOR MICE: Follow the same general directions as for rat bait, but use smaller baits and mix of them. Mice are more difficult to control than rats and complete control may require longer.

Place table spoon amount of bait at 6-12 foot intervals.

An uninterrupted supply of bait should be maintained for at least 15 days.

Do not reuse container. Destroy when empty.

CAUTION: WARFARIN contains as its active ingredient an anti-coagulant chemical which if taken by humans, domestic animals, or pets may reduce the clotting ability of the blood and serious hemorrhage may result. If eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call physician immediately.

NOTE FOR PHYSICIAN: When a human has been known to have ingested WARFARIN, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

ACCEPTED

MAY 1957

TECH-1