

ACC...
 Dec 16, 1967
 X6402

HOUSEFLIES: For reduction of fly populations, use 2 gallons per 100 gallons of water or one pint in 6 gallons of water. In dairy barns, horse and sheep barns, poultry houses, apply spray thoroughly to wet ceilings, walls and other places where flies congregate.

FOR A BAIT SPRAY: Use 2 gallons with 2 gallons unsulfurized molasses or corn syrup per 100 gallons of water. Repeat applications as necessary.

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Wash thoroughly after using. Do not apply to milking cows or to animals under one month of age. Do not apply to cows within two weeks of freshening. Avoid contamination of feed and food stuffs.

NOTE TO PHYSICIAN: Malathion, upon repeated prolonged careless use, may cause cholinesterase inhibition. Atropine is antidotal.

DR. MAYFIELD

MALATHION EMULSIFIABLE CONCENTRATE

**"CAUTION" "See left panel for additional CAUTIONS"
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS ONE PINT

For the control of lice and ticks on cattle. For the control of lice on hogs. For the control of lice, red mites, and northern fowl mites on chickens, ducks, geese, and turkeys. For use as a fly spray in horse, cattle, and sheep barns, and poultry houses.

ACTIVE INGREDIENTS:

Malathion (O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	57%
Aromatic petroleum hydrocarbon	36%
INERT INGREDIENTS	7%
Total	100%

U.S.D.A. Registration number 4644-2
Manufactured for

DR. MAYFIELD LABORATORIES

CHARLES CITY IOWA

NOTE: Do not contaminate milk, milking equipment, feed or drinking water.

DIRECTIONS

For the control of lice and ticks on cattle and lice on swine, one pint to twelve gallons of water or one gallon to 100 gallons of water as a complete coverage spray. Repeat application only if needed.

For use as a spray in poultry houses for the control of flies, lice, red mites, and northern fowl mites, apply four tablespoons per gallon or one pint to eight gallons of water. Spray liberally to walls, ceilings, roosts, nests, and adjacent areas, taking care to force spray into all cracks and crevices.

For use as a roost paint for the control of lice and red mites, apply four tablespoons per gallon of water. Brush this dilution as a roost paint at the rate of one pint per 150 feet of roost. Repeat as necessary.