



ACCEPTED

8-7-78



93% SULPHUR DUST

DIRECTIONS FOR USE

FOR BROWN ROT ON PEACHES: Use as a dust 40 to 50 lbs. per acre. Begin applications when buds show pink and continue at 10-14 day intervals until 1 week before harvest. Sulphur may burn foliage when the temperature is high. Do not make sulphur applications at such times.

FOR LEAF SPOT ON PEANUTS: Use as a dust 15 to 20 lbs. per acre. For further information as to time of application and amount consult Federal or State Experimental Stations. Start first application when disease first appears. Repeat applications at 14 day intervals during season and increase rate per acre as plants increase in size.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACCEPTED
J. M. ...
ED UNDER NO. 12130-24
FOR ECONOMIC POISON REGISTER
FUNGICIDE AND RODENTICIDE ACT
UNDER THE FEDERAL INSECTICIDE

**KEEP OUT OF THE REACH OF CHILDREN
CAUTION:**

ACTIVE INGREDIENT:

Sulphur 93.0%

INERT INGREDIENTS:

7.0%

100.0%

DO NOT RE-USE EMPTY CONTAINER. Destroy it by burying with Waste or Burning. Stay Away From Smoke or Fumes.

FARM CHEMICALS, INC.

Manufactured By

Raeford, North Carolina

Aberdeen, North Carolina

Lbs.

Net Contents

EPA Reg. No. 12130-24