

Teats, Udders, and Flanks of Cows: Immediately prior to milking wash the teats, udder, and flank of each cow with a separate clean paper towel soaked in a solution of 1/2 to 1 1/2 fluid ounces of Udder Sanitizer in 10 quarts of lukewarm (90° F.) water. Wipe dry with a clean paper towel.

Teat Cup Sanitizer: Teat cups should be thoroughly rinsed in clear water and then dipped in a solution of 1/2 fluid ounce of Udder Sanitizer in 10 quarts of water before milking each cow.

Teat Dip Directions: After each milking dip teats in a solution of 1/2 fluid ounce of Udder Sanitizer in 10 quarts of water.

Sanitizing Milking Equipment and Bulk Coolers with Udder Sanitizer:

1. Thoroughly wash all milking equipment with lukewarm (90° F.) water to remove all traces of milk.
2. Wash all milking equipment with a sanitizing solution containing 1 fluid ounce of Udder Sanitizer in 5 gallons of lukewarm (90° F.) water.
3. Allow sanitizing solution to remain in contact with milking equipment for at least 2 minutes.
4. Rinse milking equipment with warm water.
5. Before milking and after milking equipment has been re-assembled, rinse all milking equipment with sanitizing solution [1 fluid ounce of Udder Sanitizer to each 5 gallons of lukewarm (90° F.) water].
6. Allow excess sanitizing solution to drain off, but do not rinse.
7. For bulk coolers use a brush or spray.

Note: The yellow color of the sanitizing solution is an indication of sanitizing activity. If yellow color disappears prepare a fresh sanitizing solution.

CAUTION: Udder Sanitizer is a concentrated product. It must be properly diluted before use.

DILUTION TABLE:

1/2 fluid ounce to 5 gallons of water	12.5 P.P.M. Titratable Iodine
1 fluid ounce to 5 gallons of water	25 P.P.M. Titratable Iodine
3 fluid ounces to 10 quarts of water	150 P.P.M. Titratable Iodine
6 fluid ounces to 10 quarts of water	300 P.P.M. Titratable Iodine

Follow State and local health department regulations for cleaning and sanitizing.

ANCHOR LABORATORIES, INC.

A Division of PHILIPS ROXANE, Inc.
St. Joseph, Missouri 64502, U.S.A.

100-0574

ANCHOR

**Detergent - Sanitizer - Germicide
Disinfectant**

Active Ingredients:

Nonylphenoxy (ethyleneoxy) ethanol-iodine complex	13.76%
Phosphoric Acid	5.95%
Inert Ingredients:	80.29%

TOTAL 100.00%

(Udder Sanitizer supplies a minimum of 1.75% titratable iodine.)

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

See back panels for additional cautions.

ONE FL. GAL.

General Sanitation and Disinfection (Barns, Stalls, Pens, Poultry Houses, etc.):

1. Remove all livestock or poultry from buildings to be treated with the disinfectant.
2. Remove all excess filth and dirt.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Wash with warm water containing a synthetic detergent. Do not use soap.
5. Rinse with warm water (90° F.)
6. Wash with a solution of Udder Sanitizer—3 to 6 fluid ounces in 5 gallons of water.
7. Ventilate buildings, coops, and other closed spaces. Do not house poultry and animals or employ equipment until treatment has been absorbed, set or dried.
8. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be thoroughly rinsed with potable water prior to use.

Note: After contact for 2 minutes aluminum, copper, brass and zinc fittings and utensils should be rinsed with warm water.

FOR CLEANING SOWS: Scrub down the flanks and udder with a solution of 1 fluid ounce of UDDER SANITIZER in 1 gallon of lukewarm (90° F.) water, before placing the sow in farrowing pen or crate. Use a soft-bristled brush. Let sow drain dry. Rinse sow just before farrowing.

SHOE BATH: Place the shoe bath, containing 3 fluid ounces of UDDER SANITIZER in each gallon of water, inside the doorway. Before entering the building scrape extraneous dirt from the shoes, then step into the shoe bath and remain for 30 seconds. Use fresh solution daily.

CAUTION: Avoid contact with eyes. Avoid skin contact with concentrate. In case of contact, flush with large amounts of water. Call a physician for treatment of the eyes. Avoid contamination of food and feedstuffs. Harmful if swallowed. Do not use in water over 120° F. when preparing solutions. Do not apply to hot surfaces.

Avoid contamination with soap, anionics, or alkaline materials. A concentrated product—use only as directed—Must be diluted before using. Rinse empty container thoroughly with water and discard it.

Store in a cool place

Keep from freezing For external use only

Keep tightly closed when not in use
Hazardous—Not for human use

E.P.A. Reg. No. 4691-63