

Active Ingredients:

Nonylphenoxypoly (ethyleneoxy)
ethanol-iodine complex
Gamma isomer of benzene hexachloride
from lindane
Butoxypolypropylene glycol

1.5%
3.0%
30.0%
59.5%
100.0%

Inert Ingredients:

Water 1.5%

CAUTION: Harmful if swallowed. Avoid contact with the eyes and mucous membranes. Avoid contact with the human skin. In case of contact, wash immediately with soap and water. May be absorbed through the skin. Avoid contamination of feed and foodstuffs. Do not use on dogs, cats, or other household pets or on dairy animals. Do not use on newborn animals. Do not use on any animals which will be slaughtered for food.

Hazardous - Not for human use. **IDE ACT**
Store at a temperature not exceeding 75° F.
Do not reuse empty container. Destroy by crush-
ing or burning in a safe place.

ANCHOR LABORATORIES, INC.

A Division of PHILIPS ROXANE, Inc.
St. Joseph, Missouri 64502, U.S.A.

416-0574

ACCEPTED
ANCHOR
ANTISEPTIC-INSECTICIDAL
PROTECTANT DRESSING
For Abrasions, Minor Burns,
Superficial Wounds, and
use following Castration
and Branding

CAUTION: Keep out of the reach of children. See side panels for additional cautions.

NET WT. 16 OZ. (1 LB.)

USES: For topical treatment of abrasions, minor burns, superficial wounds; as a protectant following castration and branding of horses.

Kills face flies, house flies and stable flies on contact on and around the wound.

DIRECTIONS: Stir well before use. Thoroughly cover the entire injured area and approximately 1 inch around the injured tissues, using a small soft brush or other applicator. In cold weather having the product at room temperature (75° F.) will facilitate use. Apply immediately after branding or castration to aid in preventing infection. Gently cleanse injured areas with warm water and cotton gauze to remove debris and dead tissue, pat dry with a clean cloth before applying Wound Guard Smear. Repeat as necessary at 3- to 5-day intervals.

Not to be used in deep wounds, in body cavities, or on extensive burns.

CAUTION: In case of deep or puncture wounds consult a veterinarian. Keep tightly closed when not in use.

For external animal use only
E.P.A. Reg. No. 4691-70-AA

EPA Est. 4691-MO-1