

Methoxychlor & Malathion Dairy Cattle Dust is a combination of two highly effective fly-control chemicals. To prevent the number of horn flies from increasing rapidly on dairy animals begin dusting animals in the spring when flies first appear.

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

DAIRY ANIMALS—HORN FLIES (on beef cattle also) Use 2 heaping tablespoonfuls (2 oz.) per animal for control of horn flies. Dust on poll, neck, and back. Rub in lightly to insure that the material is worked down beneath the hair. If application is made by dust gun, apply against the grain of hair to force material into the hair. Do not repeat applications within a 3-week time period. Do not contaminate water troughs. Do not apply during lactation or within 2 weeks of freshening.

BEEF CATTLE—LICE, TICKS, AND HORN FLIES: For control of lice, ticks, and horn flies make liberal application to all parts of the animal's body. Rub into hair as application is made. Repeat treatment as necessary to control infestation.

HOGS—LICE: Make liberal application of dust to hogs, pens, and litter. One treatment may be sufficient; if not, repeat in 10 days or thereafter as needed for control. May be dusted on baby pigs and farrowing-house litter. Apply ¼ to ½ tablespoonful on pigs less than one month of age.

SHEEP—LICE AND KEDS: Make thorough and liberal application to entire animal. Treatment just after shearing is preferred. Repeat as necessary for control.

CAUTION

Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed or foodstuffs, or feed or water troughs. Keep out of the reach of children.


Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product as specified on this label.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ANCHOR LABORATORIES, INC.

A Division of PHILIPS ROXANE, Inc.
St. Joseph, Missouri 64502, U.S.A



ACTIVE INGREDIENTS:

Technical Methoxychlor 5.0%
Malathion 4.0%

INERT INGREDIENTS

91.0%

Total 100.0%

*Equivalent to 4.4% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% other isomers and related compounds

**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

E.P.A. Reg. No. 4691-95-AA

CAUTION: Keep out of reach of children.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Code 411-811

ANCHOR™

Methoxychlor & Malathion

**DAIRY CATTLE
DUST**

FOR HORN FLY CONTROL

On Dairy and Beef Cattle and Sheep
Also Controls Lice and Ticks

NET WEIGHT 2 LB.