

**KILLS RESISTANT AND HARD-TO-KILL ANIMAL PARASITES
NO WAITING PERIOD BETWEEN LAST APPLICATION AND SLAUGHTER
ONE QUART MAKES 25 GALLONS OF SPRAY**

Code 412-011

DIRECTIONS FOR USE

Do not use on dairy animals or in dairy barns.

ON CATTLE: To control horn flies, cattle lice, ear ticks, Lone Star ticks, Gulf Coast ticks, winter ticks, and cattle fever ticks mix one quart of LIVESTOCK SPRAY & BACKRUBBER with 25 gallons of water and stir well to form a creamy emulsion. Spray animals with a strong spray to penetrate to the skin. Wet all portions of the skin thoroughly. Approximately 2 quarts of spray are required for a full-grown cow, and proportionately less for smaller animals. Be sure not to overdose young animals. Repeat as necessary but not more often than once every two weeks. One spraying with LIVESTOCK SPRAY & BACKRUBBER will control horn flies up to 3 weeks and a single application will control lice.

ON SHEEP AND GOATS: To control sheep ticks (keds), sheep lice, goat lice, and wool maggots use one quart of LIVESTOCK SPRAY & BACKRUBBER with 25 gallons of water, mix well, and spray animals thoroughly. Repeat application at two-week intervals if necessary. Spray animals right after shearing. When dipping animals be sure to wet them thoroughly. Always stir dipping vat well just before using. Post-shearing treatment will also aid in screwworm control. Do not apply to dairy goats.

ON HOGS: To control hog lice use one quart of LIVESTOCK SPRAY & BACKRUBBER to 25 gallons of water, mix well, and spray with a strong spray to thoroughly wet the skin.

BACKRUBBER APPLICATION: For control of horn flies and lice on cattle dilute 1 part of LIVESTOCK SPRAY & BACKRUBBER with 9 parts of oil. Apply with conventional cattle oilers. Do not use on dairy animals. Do not spray animals with this solution.

E.P.A. Reg. No. 4691-97

ANCHOR LABORATORIES, INC.

A Division of PHILIPS ROXANE, Inc.
St. Joseph, Missouri 64502, U.S.A.

0172

ANCHOR

**Delnav
LIVESTOCK SPRAY
& BACKRUBBER**


ACTIVE INGREDIENTS:

Dioxathion [2,3-p-dioxanedithiol S, S-bis
(O, O-diethyl phosphorodithioate)]..... 10.5%
Related compounds 4.5%
Aromatic petroleum derivatives..... 75.0%
INERT INGREDIENTS 10.0%
100.0%

WARNING: Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.

NET CONTENTS ONE GALLON



AS A PREMISE SPRAY FOR KENNELS AND LAWNS: To control fleas and lice in kennels, dog runs, lawns, and outdoor areas dilute 1 quart of LIVESTOCK SPRAY & BACKRUBBER with 7½ gallons of water. Thoroughly spray places where animals sleep and dog-run areas. Apply 1 gallon of dilute mixture per 1,000 square feet for fleas and ticks on lawns. Do not treat pets, pet bedding, or animals with this concentration. Repeat as necessary.

LIVESTOCK SPRAY & BACKRUBBER DILUTION TABLE

Amount of Livestock Spray & Backrubber	VOLUME OF WATER			
	Cattle Spray or Dip	Sheep and Goats Spray or Dip	Hog Lice	Premise Spray for Fleas and Ticks
½ Pint	6 gallons	6 gallons	6 gallons	2 gallons
1 Quart	25 gallons	25 gallons	25 gallons	7½ gallons
1 Gallon	100 gallons	100 gallons	100 gallons	30 gallons

WARNING: May be fatal if swallowed, inhaled, or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Do not contaminate feed and foodstuffs. Wash hands thoroughly after using. Do not use or store near heat or open flame. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Wear clean clothing.

ANTIDOTE:

External—Wash with soap and water.

Internal—Call a physician immediately; large doses of atropine are antidotal. Give emetic of mustard.

Toxic to fish and wildlife. Do not contaminate water by cleaning equipment or disposal of waste and containers.

Do not reuse container. Destroy when empty.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.