

Code 412-611

DIRECTIONS
BEFORE USE READ CAUTION AND NOTICE
STATEMENTS CAREFULLY

ANIMAL TREATMENT

To control cattle and hog lice, ticks on beef cattle, and sheep ticks on sheep, thoroughly cover animals with a spray consisting of 1 quart MANGE & LOUSE SPRAY mixed with 100 gallons of water (1 tablespoonful of product in 1 1/2 gallons of water). To control psoroptic mites (responsible for sheep scab), thoroughly cover the animal with a spray consisting of 1 quart MANGE & LOUSE SPRAY mixed with 52 gallons of water (2 tablespoonfuls of product in 1 1/2 gallons of water). Repeat applications as required.

Caution: Do not spray sheep or beef cattle within 30 days of slaughter. Do not use on lactating dairy animals. Do not dip animals. Do not treat or dip calves under 3 months of age.



Sarcoptic mange on hogs may be effectively reduced or controlled by the use of 1 quart of MANGE & LOUSE SPRAY in 60 gallons of water (1 tablespoonful of product in 1 gallon of water). A second application should follow the first by one week.

Caution: Avoid spraying feed, forage, or feed and drinking troughs. Do not treat animals under 3 months of age. Spray or dip application to young animals (over 3 months of age) should be limited to the 0.03% lindane concentration. Do not treat sows within 2 weeks of farrowing or at least 3 weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not spray hogs within 60 days of slaughter.

ANCHOR LABORATORIES, INC.

A Division of PHILIPS ROXANE, Inc.
St. Joseph, Missouri 64502, U.S.A.

0673



LINDANE MANGE &
LOUSE SPRAY

ONE POUND LINDANE PER GALLON

ACTIVE INGREDIENTS:	
Gamma isomer of benzene hexachloride from lindane	12.5%
Xylene	82.5%
INERT INGREDIENTS:	5.0%
Total	100.0%

CAUTION: Keep out of the reach of children.
See side panels for additional cautions.

FOR APPLICATION ON ANIMALS
AND BUILDINGS

NET CONTENTS ONE QUART

RESIDUAL SPRAY

Flies, mosquitoes, and gnats can be controlled by spraying walls, ceilings, window ledges, and other resting places of insects in barns and other buildings with a solution of MANGE & LOUSE SPRAY and water. Usually 1 quart of MANGE & LOUSE SPRAY diluted with 10 gallons of water (2 tablespoonfuls of product in 2 1/2 pints of water) will cover 5,000 square feet of surface, or 500 square feet per gallon of diluted material. For roaches, use 1 quart of MANGE & LOUSE SPRAY diluted with 6 1/2 gallons of water (2 tablespoonfuls of product in 1 1/2 pints of water). Cover all feed troughs. Remove animals from barn while spraying.

Caution: Do not use in poultry houses. Residual spray treatment is not to be used in dairy barns or milk rooms.

WARNING

MAY BE ABSORBED THROUGH THE SKIN. AVOID INHALATION AND SKIN CONTACT. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE ON HOUSEHOLD PETS OR HUMANS. HARMFUL IF SWALLOWED. KEEP OUT OF THE REACH OF CHILDREN.

DO NOT USE ON FEED OR FORAGE TO BE FED TO DAIRY ANIMALS, OR ANIMALS BEING FINISHED FOR SLAUGHTER. DO NOT CONTAMINATE MILK OR DRINKING WATER.

DO NOT CONTAMINATE WATER SUPPLIES BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES AND CONTAINERS.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY. KEEP AWAY FROM HEAT, SPARKS, AND OPEN FLAME.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 4691-98