

a manna a construction of the construction of the

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For use on Beef Cattle, Dairy Cattle, Horses, Sheep, Goats, and Swine

WORM AND BLOWFLY INFESTATIONS:

Apply with a swab or brush, working into deep pockets. Also spread smear around and over drainage area below wound. Repeat applications at 5- to 7-day intervals until wound is healed. Freely bleeding wounds may require several applications.

TO PREVENT SCREW WORM AND **BLOWFLY INFESTATIONS:**

Apply directly to all open or fresh wounds resulting from dehorning, castration, shear cuts, etc. Also treat navels of newborn animals. Freely bleeding wounds may require several applications. Repeat applications at 5- to 7-day intervals until wound is healed.

Notice: Solier makes no warranty, expressed or implied, con-cerning the use of this product other than inducated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.


ANCHOR LABORATORIES, INC.

A Division of PHILIPS ROXANE, Inc.

St. Joseph, Missouri 64502, U.S.A.

EPA Reg. No. 4691-101

EPA Est. 4691-MO-1



DIRECTIONS FOR USE **GENERAL CLASSIFICATION**

a violation of Federal law to use this product In a manner inconsistent with its labeling

se on Beef Cattle, Dairy Cattle, Horses, Sheep, Goats, and Swine

SCREW WORM AND BLOWFLY INFESTATIONS:

Apply with a swab or brush, working into deep pockets. Also spread smear around and over drainage area below wound. Repeat applications at 5- to 7-day intervals until wound is healed. Freely bleeding wounds may require several applications.

TO PREVENT SCREW WORM AND **BLOWFLY INFESTATIONS:**

Apply directly to all open or fresh wounds resulting from dehorning, castration, shear cuts, etc. Also treat navels of newborn animals. Freely bleeding wounds may require several applications. Repeat applications at 5- to 7-day intervals until wound is healed.

Notice: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 4691-101 EPA Est. 4691-MO-1

> ANCHOR LABORATORIES, INC. A Division of PHILIPS ROXANE, Inc. St. Joseph, Missouri 64502, U.S.A.



Active Ingredients: Ronnel [0,0-dimethyl 0-(2,4,5-

trichiorophenyi) phosphorothiosts] 5.0% Xylene Inert Ingredients TOTAL 100.0% KEEP OUT OF THE REACH OF CHILDREN.

CAUTION:

IF SWALLOWED - Do not induce vomiting. Consult your physician immediately. IF ON SKIN — Wash area with soap and water and rinse with a large volume of water. IF IN EYES — Immediately flood eyes with copious amounts of water and consult your physician.

Precautionary Statements



Active Ingredients: Ronnei [0.0-dimethyl 0-(2,4,5-TOTAL 100.0%

CAUTION:

IF SWALLOWED --- Do not induce vomiting. Consult your physician immediately. IF ON SKIN — Wash area with soap and water and rinse with a large volume of water. IF IN EYES --- Immediately flood eyes with copious amounts of water and consult your physician.

> See Side Panels For Additional Precautionary Statements

> > Net Weight 14 Ounces

0376



Ronnel

SUNEW WUNN SHERE

STATEMENT OF PRACTICAL TREATMENT

See Side Panels For Additional

Net Weight 14 Ounces

PRECAUTIONARY STATEMENTS CAUTION

Avoid contact with skin, eyes and clothing. Wash hands thoroughly after using. Harmful if swallowed. Avoid contami-

nation of feed and foodstuffs. NOTE TO PHYSICIAN: This material is a cholinesterase in-hibitor. Treat symptomatically. Atropine is an autidola. Ronnel is a cholinesterase inhibitor. Do not use this product on

animals simultaneously or within a few days opfore or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Atropine is an antidote. Consult a veterinarian at first sign of an adverse reaction. ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not use near lakes, streams, or ponds or where run off is likely to occur.

Apply this product only as specified on this label.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT CONTAMINATE WATER, FODD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PRO-HIBITED.

PESTICIDE DISPOSALI PESTICIDE THAT CANNOT BE USED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL: OFFER FOR RECYCLING, RECONDI-TIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

GENERAL: CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.



Ronnel

KEEP OUT OF THE REACH OF CHILDREN.

STATEMENT OF PRACTICAL TREATMENT

PRECAUTIONARY STATEMENTS CAUTION

Avoid contact with skin, eyes and clothing. Wash hands thoroughly after using. Harmful if swallowed. Avoid contamination of feed and foodstuffs.

NOTE TO PHYSICIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote. Ronnel is a cholinesterase inhibitor. Do not use this product on

animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals. Atropine is an antidote. Consult a veterinarian at first sign of an adverse reaction.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not use near lakes, streams, or ponds or where run off is likely to occur.

Apply this product only as specified on this label.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PRO-HIBITED.

PESTICIDE DISPOSAL: PESTICIDE THAT CANNOT BE USED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL: OFFER FOR RECYCLING, RECONDI-TIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

GENERAL: CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.