



ANCHOR PERMECTRIN[®] II

10% EMULSIFIABLE INSECTICIDE
LONG LASTING LIVESTOCK AND PREMISE SPRAY

ACTIVE INGREDIENTS

Permethrin
(3-phenoxyphenyl) methyl (±)-cis, trans-3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate* 10.00%
INERT INGREDIENTS 90.00%
TOTAL 100.00%

*Cis/trans ratio: Min. 35% (±) cis, and max. 65% (±) trans isomers.
Contains 0.75 lbs. permethrin per gallon.

CAUTION: Keep out of the reach of children.
See side panels for additional cautions.

NET CONTENTS: 1 QUART
(0.946 liters)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION: May be harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. In case of eye contact, immediately flush eyes with plenty of water. In case of skin contact wash skin with soap and water. Get medical attention if irritation persists. If swallowed call a physician. Vomiting should be supervised by a physician because of possible pulmonary damage due to aspiration of the solvent. **ENVIRONMENTAL HAZARDS**
This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.
PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE AND DISPOSAL

Keep container sealed when not in use. Do not store near food, feed, or bedding. In case of spill on floor or paved surface, wash up with sand, earth, or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with state and local regulations. Do not reuse container. Triple rinse empty container and add rinse to mixture of spray area. Do not dump contents in any body of water. Do not use for any other purpose than that for which it is labeled. See side panels for additional cautions.

Directions for use cont'd FOR USE ON

Work and pets and their premises for control

of flies, lice, mites and ticks. Mix PERMECTRIN[®] II and apply the undiluted material to animals and pest breeding or resting surfaces at the rates shown below. These dilutions and rates will provide most efficient pest control under conditions of heavy pressure when good contact is achieved. Timing and frequency of application should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray feed, food, or water. Retreat as needed but not more often than once every two weeks.

PESTS	PERMECTRIN II DILUTIONS FOR USE	HOW TO APPLY
PESTS ON FARM PREMISES (Barns, Stables, Feedlots, Stock pens, Poultry and Livestock Housing)		
Houseflies, Stableflies, Lesser houseflies and other manure breeding flies. Also in control of cockroaches, mosquitoes and spiders	Mix 1 qt in 5 gal of clean water. Use undiluted in mist blower or space spray. Use 1 qt in 6 gal of oil.	Spray all surfaces to runoff with diluted emulsion using 1 gal per 750 sq. ft. Mist or fog 4 oz per 1000 sq. ft.
PESTS ON LARGE ANIMALS (Dairy or Beef Cattle and Horses)		
Face flies, Horn flies, Mosquitoes, Stableflies	Mix 1 qt in 20 gal of clean water for backrubbers	Animals self apply. Recharge backrubber area as needed.
Face flies, Horn flies, Mosquitoes, Stableflies, Manure flies, Lice, Mites, Ticks	Mix 1 qt in 200 gal of clean water for whole body area spraying	Spray thoroughly over entire animal. For Lice or Mites a second treatment is recommended 14-21 days later.
Ear Ticks, Face flies, Horn flies	Mix 2 oz in 1 gal of clean water for spot applications	Apply 1/2 oz per ear or 2-4 oz per face or 12-16 oz along the backbone

© 1991 Anchor Laboratories, Inc.
EPA Reg. No. 4691-
EPA EST No. 70-KY-1

ANCHOR LABORATORIES, INC.

A Division of PHILIPS DOWNEY, Inc.
St. Joseph, Missouri 64501, U.S.A.

0353

ACCEPTED
OCT 11 1993
9091-108