

Code 401-411

ANCHOR PERMECTRIN II

10% EMULSIFIABLE INSECTICIDE
LONG LASTING LIVESTOCK AND PREMISE SPRAY

ACTIVE INGREDIENTS

Permethrin
(3-phenoxyphenyl) methyl (±)-cis, trans-3-(2,2-dichloroethenyl)-2,
2-dimethylcyclopropanecarboxylate* 10.00%
INERT INGREDIENTS 90.00%
TOTAL 100.00%

*Cis/trans ratio: Min. 35% (±) cis, and max. 65% (±) trans isomers.

Contains 0.75 lbs. permethrin per gallon.

CAUTION: Keep out of the reach of children.
See side panels for additional cautions.

NET CONTENTS: 1 QUART
(0.946 liters)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION: May be harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. In case of eye contact, immediately flush eyes for at least 15 minutes with water. In case of skin contact wash skin with soap and water. Get medical attention if irritation persists. If swallowed call a physician. Vomiting should be supervised by a physician because of possible pulmonary damage due to aspiration of the solvent.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE AND DISPOSAL

KEEP CONTAINER SEALED WHEN NOT IN USE. PROTECT FROM FREEZING.

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. In case of spill on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with state and local regulations.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinse liquids that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Do not reuse container. Triple rinse residue from empty container and add rinsate to mixture in spray tank. Dispose of container in sanitary landfill in accordance with state and local regulations or by other approved State and local procedures.

Distributed by

ANCHOR LABORATORIES, INC.

A Division of FMC CORPORATION
St. Joseph, Missouri 64502 U.S.A.

0353

GENERAL USE CLASSIFICATION

Mange Mites

Mix 1 qt in 100 gal. water.

Spray or dip animals. Retreat after 14 days spraying walls & floor and replace bedding to kill late hatching developing stages. Spray, paint or dip to apply 1 pint per dog or pig, especially around ears. Spray 1-2 oz per bird or 1 gal. per 100 birds directed toward the vent area.

Managers, Hog Lice, Flies, Ticks

Mix 1 qt in 100 gal clean water for area or spot application.

Poultry Mites

Mix 1 qt in 50 gal. clean water to spray birds and areas.

Directions for use cont'd FOR USE ON

Horses, beef and dairy cattle, swine, poultry and pets and their premises for control of flies, lice, mites and ticks. Mix PERMECTRIN II and apply the use-diluted material to animal's and pest breeding or resting surfaces at the rates shown below. These dilutions and rates will provide most efficient pest control under conditions of heavy pressure when good contact is achieved. Timing and frequency of application should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray feed, food or water. Retreat as needed but not more often than once every two weeks. Wash udders thoroughly before milking.

PESTS	PERMECTRIN II DILUTIONS FOR USE	HOW TO APPLY
PESTS ON FARM PREMISES (Stables, Barns, Feedlots, Stables, Poultry and Cattle Housing)		
Houses, Stables, Lower Horse and Cattle Housing, and other structures	Mix 1 qt in 25 gal of clean water	Spray or brush on surfaces with diluted mixture using 1 gal per 100 sq ft.
Stables, horse and cattle housing, and other structures	Mix 1 qt in 25 gal of clean water	Mix or spray 1 qt per 100 sq ft.
PESTS ON LARGE ANIMALS (Dogs, Cats, and Horses)		
Face, Neck, and Horses	Mix 1 qt in 10 gal of clean water	Wash and spray. Retreat after 14 days if needed.
Face, Neck, and Horses	Mix 1 qt in 200 gal clean water for whole body, and area spraying	Spray thoroughly, cover entire animal for head to tail. Retreat after 14 days if needed.
Ear Ticks, Face, Neck, and Horses	Mix 2 oz in 1 gal of clean water for spot application	Spray 1-2 oz per ear or 1-2 gal per 100 sq ft of body and bedding.
PESTS ON SWINE, POULTRY AND SMALL ANIMALS		
Mange Mites	Mix 1 qt in 100 gal water.	Spray or dip animals. Retreat after 14 days spraying walls & floor and replace bedding to kill late hatching developing stages.
Managers, Hog Lice, Flies, Ticks	Mix 1 qt in 100 gal clean water for area or spot application.	Spray, paint or dip to apply 1 pint per dog or pig, especially around ears.
Poultry Mites	Mix 1 qt in 50 gal clean water to spray birds and areas.	Spray 1-2 oz per bird or 1 gal per 100 birds directed toward the vent area.

Do not ship swine for slaughter within 5 days of last treatment.

401-103

BEST DOCUMENT AVAILABLE