

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION:

Keep out of reach of children. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT: In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water. Use this product only as specified on this label. Do not contaminate water, streams or creeks.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL: Open as needed. Keep container sealed when not in use. Do not store near food or feed. Remove dust before slaughter. Dispose of in accordance with state and local regulations. Do not reuse container.

EPA Est. No. 10543-1-1001
EPA Reg. 4691-110

Distributed by
ANCHOR LABORATORIES, INC.
A Division of PHILLIPS DORANE, Inc.
St. Joseph, Missouri 64502, U.S.A.

ANCHOR

**PERMECTRIN
0.25% DUST**

ACTIVE INGREDIENTS: Permethrin -
* (3-phenoxyphenyl) methyl (+) cis/
trans 3-(2-dichlorobenzoyl)-2, 2-
dimethyl cyclopropanecarboxylate 0.25%
INERT INGREDIENTS 99.75%
100.00%

U.S. Patent No. 4,024,163
* Cis/trans ratio: min. 35% (+) cis and max. 35% (+) trans

For Control of HORN FLIES & LICE on BEEF & DAIRY CATTLE—As an aid in reducing FACE FLY population on BEEF and DAIRY CATTLE—Control of Lice on Swine.

Use In Dust Bags, Shaker Can, or Mechanical Dust Applicator.

CAUTION: Keep out of reach of children
See side panels for warning statements

NET WT 12½ LBS

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

BEEF & DAIRY CATTLE

Horn Flies, Lice, Face Flies: Place contents of this package in any commercially available dust bag. Suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed, or minerals. Bags may also be suspended in feeding sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the stall through which cows leave the milking barn. In all cases the bags should be adjusted so that the bottom of the bag will hang four to six inches below the feeding of the cattle. For reduction of face flies, the bags must be located so the animals will be forced to use them daily and hung at a height to insure that the faces of the cattle will be dusted.

Horn Flies, Lice—Direct Application: Apply 2 to 4 ounces of dust per animal by shaker can over the head, neck, shoulders, back and tailhead. Repeat as necessary.

SWINE

Lice on Swine—Direct Application: Apply not more than 1 oz. (3 level tablespoons) per head as a uniform coat to the head, shoulders, and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.

Bedding Treatment: Apply 2 oz. (6 level tablespoons) uniformly over each 33 square feet of fresh, dry bedding by use of a shaker can or a suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.

In severe infestations both individual animals and the bedding may be treated as directed above.

Do not ship animals for slaughter within 5 days of last treatment.
*NOTE: 1 oz. of dust material is approximately three level tablespoons.

BEST DOCUMENT AVAILABLE

ACCEPTED
JUN 24 1983
U.S. Dept. of Agriculture
Federal Inspection Service
as required for this
commodity under
EPA Reg. 105. 4691-110