

ACCEPTED
 SEP 15 1984
 4891-110

Code 404-011

ANCHOR
**PERMECTRIN™ 0.25%
 DUST**

For Control of **HORN FLIES & LICE** on **DEEF & DAIRY CATTLE**--As an aid in reducing **FACE FLY** population on **DEEF and DAIRY CATTLE**--Control of **LICE** on **SWINE** and **CONTROL OF NITS ON POULTRY**.
 Use in dust bags, shaker can, or mechanical dust applicator

ACTIVE INGREDIENT:	
Permethrin (3-phenoxyphenyl) methyl (±)- cis, trans-3-(2,2-dichloroethenyl)-2,2- dimethylcyclopropanecarboxylate	0.25%
INERT INGREDIENTS	99.75%
	100.00%

U.S. Patent No. 4,024,183
 *Cis/trans ratio: min. 35% (±) cis and max. 35% (±) trans

CAUTION: Keep out of reach of children
 See side panels for warning statements

NET WT 12-1/2 LBS

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND
 DOMESTIC ANIMALS
 CAUTION**

Harmful if inhaled. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT: In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water. Use this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL: Open as needed. Keep container sealed when not in use. Do not store near food or feed. Remove dust before slaughter. Dispose of in a sanitary landfill or by incineration in accordance with state and local regulations. Do not reuse container.

EPA Est. No. 10543-MO-01
 EPA Reg. No. 4091-110

Distributed by
ANCHOR LABORATORIES, INC.
 A Division of PHILIPS ROXANE, Inc.
 St. Joseph, Missouri 64502, U.S.A.

DIRECTIONS FOR USE:

GENERAL USE CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DEEF & DAIRY CATTLE

Horn Flies, Lice, Face Flies: Place contents of this package in any commercially available dust bag, suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the exit through which cows leave the milking barn. In all cases, the bags should be adjusted so that the bottom of the bag will hang four to six inches below the top line of the cattle. For reduction of face flies, the bags must be located so the animal's will be forced to use them daily and hung at a height to insure that the faces of the cattle will be dusted.

Horn Flies, Lice-Direct Application: Apply 2.0 ounces of dust per animal by shaker can over the head, neck, shoulders, back and tailhead. Repeat as necessary.

SWINE

Lice on Swine-Direct Application: Apply not more than 1 oz (3 level tablespoons) per head as a uniform coat to the head, shoulders and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary, but not more often than once every ten days.

In severe infestations, both individual animals and the bedding may be treated as directed above.

Do not ship animals for slaughter within 5 days of last treatment.
NOTE: 1 oz of this material equals approximately three level tablespoons.

POULTRY

Female Nits-Direct Application: apply not more than 1 cc/100 birds directly last to thoroughly cover vent area.

3. N/A