



A13 Code 401-811

**ANCHOR.**

**PERMECTRIN™ 25%  
WETTABLE POWDER**  
Long-lasting Premise Fly Spray  
and Livestock Insecticide

**ACTIVE INGREDIENTS**  
Permethrin (3-phenylphenyl methyl (1)-  
cis,trans-3,4,5-trichlorobenzyloxy)-2,  
8-dimethylcyclohexanecarboxylate 75.00%  
**INERT INGREDIENTS** 15.00%  
TOTAL 100.00%  
\*Contains 100% 25% (1) cis, and 75% (1) trans isomers.  
EPA EST. NO. 279-MJ-1  
EPA Reg. No. 4691-

**Precautionary Statements**  
**Hazards to Humans and Domestic Animals**  
**WARNING**  
May Cause Eye Irritation  
Harmful if Swallowed or Inhaled

Avoid breathing dust or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling.

**STATEMENT OF PRACTICAL TREATMENT: IF IN EYES.** Flush with water for 15 minutes. Get medical attention if irritation persists. **IF SWALLOWED:** Drink for 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. **DO NOT add directly to water.**

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**STORAGE AND DISPOSAL**

Mix as needed, do not store diluted material. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep up and remove to chemical waste area. Dispose of in accordance with state and local regulations. Do not reuse container. Triple rinse residue from empty container and add rinse to mixture in spray tank. Dispose of container in sanitary landfill in accordance with state and local regulations.

Distributed by  
**ANCHOR LABORATORIES, INC.**

A Division of PHILIPS ROXANE, Inc.  
St. Joseph, Missouri 64502, U.S.A.

0383

**CAUTION:** Keep Out of Reach of Children.  
See side panels for additional precautions and directions.

**NET CONTENTS**  
**ONE POUND (16 ounces)**  
**(454 Grams)**

BEST DOCUMENT AVAILABLE

4691-112

# **DIRECTIONS FOR USE SPECIMPH LABEL**

It is a violation of both State and Federal Law to use this product in a manner inconsistent with its labeling. For use on cattle, horses, swine, sheep, dogs, poultry, and other premises for control of flies, cockroaches, mosquitoes, and ticks. Mix 25% WP and apply the use-diluted spray mix to animals and pest infested or resting surfaces such as walls, ceilings, walls, and floors at the rates shown below. These dilutions will provide efficient control under conditions of heavy pest pressure when good contact is achieved. Strong spray mixes completely covering interior surfaces protected from sun, dust, rain and washing should remain effective for up to 10 weeks. The same levels will generally be effective on exterior surfaces for 4-6 weeks, but provide 7-10 weeks control of northern house flies on poultry. As conditions vary widely, variations in effective intervals are common and retreatment should be applied as needed but never more than once every two weeks. Manure removal should precede spraying and control of flies is continuous in manure pile breeding sites. Do not spray food or water. Most animals should not be slaughtered within 14 days of last treatment.

| PESTS TO BE CONTROLLED                                                   | WHERE AND WHEN TO APPLY OR REAPPLY                                                                                                                                    | HOW TO MIX SPECIMPH 25% WP                                                                                                | HOW TO APPLY SPRAY MIXES                                                                                  |
|--------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| <b>SEE FRONT PANEL FOR STATES WHICH HAVE APPROVED USE ON HUMANS ONLY</b> |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| <b>INDOOR PREMISES ONLY - Remove food and water</b>                      |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| House flies, Cockroaches, Mosquitoes, Ticks                              | All surfaces in rooms and rooms containing animals, including houses, horse stalls & other sheep sheds, kennels & other animal sheds. Retreat at 2-10 weeks.          | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. | Apply to surfaces in rooms containing animals using 1 qt/90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. |
| <b>ONLY ON EXTERIOR PREMISES - Remove animals, feed and water</b>        |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| House flies, Mosquitoes, Ticks, and Fleas                                | Walls, doors, fences, & other surfaces around farm buildings, animal pastures, kennels, and other areas. Feed & water should be removed after 4-6 weeks of treatment. | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. | Apply to surfaces in rooms containing animals using 1 qt/90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. |
| <b>POULTRY IN PASTURE OR BARN - Remove feed and water</b>                |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| House flies, Cockroaches, Mosquitoes, Ticks                              | Ceilings, walls, and floors. Repeat at 7-10 weeks or as needed.                                                                                                       | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water.                                                              | Spray to surfaces in rooms containing animals using 1 qt/90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. |
| <b>SEE FRONT PANEL FOR STATES WHICH HAVE APPROVED USE ON ANIMALS</b>     |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| <b>TO BEET, DIRT, &amp; LACTATING DAIRY CATTLE AND HORSES IN STALLS</b>  |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| Face flies, House flies, Mosquitoes, Ticks, and Fleas                    | Face & legs. Repeat every 3-5 weeks or as needed.                                                                                                                     | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water.                                                              | Spray to surfaces in rooms containing animals using 1 qt/90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. |
| Face flies, Mosquitoes, Ticks                                            | Spot treat ears, face, and neck. Repeat at 3-5 weeks or as needed.                                                                                                    | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water.                                                              | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water.                                              |
| <b>TO SWINE, SHEEP, OR DOGS IN STALLS - Remove feed and water</b>        |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| House flies, Mosquitoes, Ticks, and Fleas                                | Animals. Repeat at 3-5 weeks or as needed.                                                                                                                            | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water.                                                              | Spray to surfaces in rooms containing animals using 1 qt/90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. |
| <b>TO Poultry in HOUSES OR COUSINS - Remove feed and water</b>           |                                                                                                                                                                       |                                                                                                                           |                                                                                                           |
| House flies                                                              | Walls, doors, and floors. Repeat at 7-10 weeks or as needed.                                                                                                          | Apply 1 qt. 90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water.                                                              | Spray to surfaces in rooms containing animals using 1 qt/90 gal or 1 lb 10 gal or 1/2 lb 20 gal of water. |