

4691-113

# NEW

LONGER, Floppy  
Design Directs  
More Insecticide  
to Sites Pests  
Prefer

# ANCHOR

## PERMECTRIN™ POULTRY MITE STRIPS

### 10% PERMETHRIN INSECTICIDE STRIPS

FOR USE AS

AN AID TO CONTROL AND PREVENT LICE, HOUSEFLIES, STABLE FLIES  
AND NORTHERN FOWL MITES ON POULTRY.

**ACTIVE INGREDIENTS:**

Permethrin (3-phenoxyphenyl) methyl (+)-  
cis, trans-3-(2,2-dichloroethenyl)-2,  
2-dimethylcyclopropanecarboxylate\*

INERT INGREDIENTS..... 90.00%  
TOTAL 100.00%

\*Cis/trans ratio: Min. 35% (+) cis, and max. 65% (+) trans isomers

KEEP OUT OF REACH OF CHILDREN

**CAUTION**  
CONTENTS  
In EPA Reg. 4691-113

EPA Est. No. 42836-TX-01  
EPA Reg. 4691-113  
Batch No.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION:** Wash thoroughly with soap and water after handling, and before eating or smoking.


**STATEMENT OF PRACTICAL TREATMENT:** In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

**ENVIRONMENTAL HAZARDS:** Do not use on or near fish. Do not apply directly to water.

**PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

**STORAGE AND DISPOSAL:** Open as needed. Keep container sealed when not in use. Do not store near food or feed. Remove tapes before slaughter. Wrap used strips and dispose of in trash containers.

See back panel for additional directions for use.



- Minimizes Labor of application
- Helps increase egg size and production rates
- Helps prevent weight loss caused by flying pests
- No withdrawal required FOR EGGS OR MEAT

**NET CONTENTS:**  
1000 STRIPS PER PACKAGE  
1000 ATTACHMENT CLIPS  
**NET WEIGHT:**  
17.5 GRAMS PER STRIP  
19 KG PER PACKAGE

ANCHOR  
PERMECTRIN POULTRY MITE STRIPS  
10% PERMETHRIN INSECTICIDE STRIPS

**DIRECTIONS FOR USE**


IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

ATTACH ONE STRIP PER CAGE TO PREVENT NORTHERN FOWL MITES. Strips remain effective for 10 to 12 months. Replace as necessary.


Attach two strips per cage to control lice AND Northern fowl mites and as an aid to control stable flies and houseflies.

Use one strip per cage attached inside colony cages CONTAINING UP TO 8 BIRDS PER CAGE. They should be suspended from the top center of the cage using pin rings, large paper clips or other strong attachment devices. PERMECTRIN POULTRY MITE STRIPS are tough and resist tearing out but have a hole at both ends so that they may be reversed and resuspended if they are torn out.

They must be suspended at hen body height inside poultry cages to prevent northern fowl mite infestations most efficiently.



0487



Distributed by  
**ANCHOR LABORATORIES, INC.**  
A Division of PHILIPS ROXANE, Inc.  
St. Joseph, Missouri 64502, U.S.A.