



Permethrin™ CDS Pour-On Insecticide

Synergized Formula Concentrated Dose

Controls lice and flies on cattle

Apply as a pour-on, mist spray or back rubber on beef cattle, lactating and non-lactating dairy cattle and calves.

Active Ingredients

Permethrin (3-phenoxyphenyl) methyl (±)-cis, trans-3 (2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate*	7.4%
Piperonyl Butoxide Technical**	7.4%

Inert Ingredients***	85.2%
----------------------------	-------

Total	100.0%
-------	--------

*cis/trans ratio: Min. 35% (±) cis and max. 65% (±) trans.

**Equivalent to Min. 5.92% (butylcarbityl) (6-propylpiperonyl) ether and 1.48% related compounds.

***Contains petroleum distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements.

08301-01

**NET CONTENTS:
1 Pint (480 mL)**

Code 408-311

ACCEPTED

08 MAR 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 4691-121

Bar Code
#0 12313 40831 2
for position only

ECL INSIDE PANEL #1:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If on Skin: Wash with plenty of soap and water. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and other aquatic invertebrates. Do not add directly to water. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flames.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ECL INSIDE PANEL #2:

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: This is a highly concentrated formulation. Use only with an accurate measuring container or applicator.

EXCEPT FOR BACK RUBBER / READY TO USE - No dilution necessary.

Apply to: Lactating and Non-lactating Dairy Cattle. Beef Cattle and Calves

Target Insects	Application Instructions
Lice, Horn Flies, Face Flies. Aids in control of Horse Flies, Stable Flies, House Flies, Mosquitoes and Black Flies. Controls Horn Flies up to 6 weeks after application.	<p>Dosage: Apply 2 mL per 100 lbs body wt of animal, up to a maximum of 20 mL for any one animal when applied as pour-on or spray.</p> <p>Pour-On: Pour correct dose along back and down face.</p> <p>Ready-To-Use Spray: Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears.</p> <p>Back Rubber Use: Mix 64 mL (2.1 oz) per gallon of mineral oil. Keep rubbing device charged. Results improved by daily forced use.</p>



ECL INSIDE PANEL #3:

Apply to: Premises - in and around Beef, Dairy, Swine and Poultry Premises, Animal Hospital Pens and Kennels and "outside" Meat Processing Premises

Target Insects	Application Instructions
House Flies, Stable Flies, Face Flies, Gnats, Mosquitoes, Black Flies, Fleas, Little House Flies (<i>Fannia</i> spp.). Aids in control of Cockroaches, Ants, Spiders and Crickets.	Ready-To-Use Spot or Premise Spray: Use undiluted in a mist sprayer. Apply directly to surface to leave a residual insecticidal coating, paying particular attention to areas where insects crawl or alight. 500 mL will treat approximately 7,200 square feet.

For cattle, repeat treatment as needed, but not more than once every two weeks. For optimum lice control, two treatments at a 14-day interval are recommended.

Special Note: PERMECTRIN™ CDS Pour-On is not effective in controlling cattle grubs.

PERMECTRIN™ CDS Pour-On is an oil-base, ready-to-use product that may leave an oily appearance on the hair coat of some animals.

PERMECTRIN™ CDS Pour-On should be used in an integrated pest management system which may involve repeat treatments and the use of other pest control practices.

ECL INSIDE PANEL #4:

NOTICE OF WARRANTY

BOEHRINGER INGELHEIM ANIMAL HEALTH, INC., MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS OF ANY PARTICULAR PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

EPA Reg. No.: 4691-121

08304-01


EPA Est. No.: 62478-FL-1A

44616-MO-1B

56493-KS-1D

First letter of run code on container denotes Est. No. of manufacturer
Mfd. & Printed in U.S.A.

Mfd. for **Anchor Division,**
Boehringer Ingelheim Animal Health, Inc.
St. Joseph, MO 64506, U.S.A.



4 7 4

ECL OUTSIDE PANEL:

INDICATIONS

Apply as a pour-on, mist spray or back rubber on beef cattle, lactating and non-lactating dairy cattle and calves.

Apply 2 mL per 100 lbs body wt of animal, up to a maximum of 20 mL for any one animal for control of lice, horn flies and face flies on cattle. Also aids in control of horse flies, stable flies, house flies, mosquitoes and black flies. Mix 64 mL (2.1 oz) per gallon of mineral oil for back rubber use.

Use undiluted in a mist sprayer for control of flies, gnats, mosquitoes and fleas in and around beef, dairy, swine and poultry premises, animal hospital pens and kennels and "outside" meat processing premises.

See chart inside for complete directions.

PRECAUTIONARY STATEMENTS

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish and other aquatic invertebrates.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flames.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

KEEP CONTAINER SEALED WHEN NOT IN USE.

See label for complete directions on storage and disposal.

EPA Reg. No.: 4691-121

08304-01

EPA Est. No.: 62478-FL-1A

44616-MO-1B

56493-KS-1D

First letter of run code on container denotes Est. No. of manufacturer
Mfd. & Printed in U.S.A.



Mfd. for Anchor Division,
Boehringer Ingelheim Animal Health, Inc.
St. Joseph, MO 64506, U.S.A.

