

Ectrin Insecticide

Water
Dispersible
Liquid
Livestock and
Premise
Spray

Active Ingredient:

Permethrin w/w
(CAS No. 51630-58-1) ... 10.0%

Other
Ingredients* 90.0%

Total 100.0%

*This product contains 0.8 lbs. of
Ectrin Insecticide per gallon

Store above 32°F

SHAKE WELL
BEFORE USE



Boehringer
Ingelheim

Keep Out of
Reach of Children
CAUTION

See back panel
for additional
precautionary
statements and
Directions for Use.



MAY 23 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 4691-141

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

General Directions

Ectrin Insecticide WDL is suitable for use in conventional power or low pressure knapsack sprayers. Follow the Use Directions to select the recommended dilution for a specific insect use. Shake Well Before Use.

Livestock Use Directions

Animal	Insect	Applica- tion Method	Dilution Ectrin WDL in Water	Remarks
Swine	Lice, Mange Mites	Spray	1 qt./50 gal. or 1 oz./1 1/4 gal.	Wet entire animal with up to 8 oz. of dilution with particular attention to heavily infested areas. Repeat in fourteen days if necessary. Do not apply within 1 day of slaughter.
	Lice	Pour- on***	1 qt./25 gal. or 2 oz./1 1/4 gal.	Apply 4 oz. of dilution on head and down midline of animal. Repeat in fourteen days if necessary. Do not apply within 1 day of slaughter.

Premise Use Directions

Animal	Insect	Applica- tion Method	Dilution Ectrin WDL in Water	Remarks
Horse barns, swine buildings	House flies, Fannia	Spray	1 qt./25 gal. or 2 oz./1 1/4 gal.	Spray to point of runoff (approximately 1 gal./750 sq. ft. of surface area). Repeat in fourteen days if necessary.*
			1 qt./10 gal. or 3 1/2 oz./1 gal.	For longer residual control, apply as above. Repeat in thirty days if necessary.*

*Spray ceilings, walls and walkways. Use proper sanitation and manure management to reduce fly-breeding sites. Do not treat horse barns if horses are intended for slaughter.

**Repeat application in 30 days if necessary. Do not apply Ectrin Insecticide more often than 2 times in the spring and 2 times in the fall pest seasons. (One application after shearing is usually adequate for keds control.) Do not apply within 2 days of slaughter.

***For pour-on mixtures, add 2 fl. oz. of a wetting agent per 5 gal. of water. A household detergent such as Joy or Ivory can be used.

Precautionary Statements

Hazards to Humans

Caution

Harmful if swallowed or absorbed through skin. Avoid contact with skin or eyes. Avoid breathing vapors and spray mist. Wear goggles, protective clothing and gloves while handling. Remove and wash contaminated clothing before reuse.

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-821-7467 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Contains petroleum distillate - vomiting may cause aspiration pneumonia.	

Environmental Hazards

This product is highly toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Boehringer Ingelheim Vetmedica, Inc.

St. Joseph, MO 64506-2002 U.S.A.

(800) 821-7467

EPA Reg. No. 4691-141

EPA Est. No. 4691-KS-01

655002L-01-9908

Animal	Insect	Applica- tion Method	Dilution Ectrin WDL in Water	Remarks
Horses	Horn flies, Face flies, House flies, Stable flies	Spray	1 qt./25 gal. or 2 oz./1 1/4 gal.	Apply 8 oz. as light spray with attention to head and legs. Do not treat animals intended for slaughter. Repeat as necessary.
Sheep, Goats (non- lactating)	Keds, Lice	Spray	1 qt./100 gal. or 1 oz./3 gal.	Wet animal with up to 1 qt. of dilution.**
		Pour on***	1 qt./12 1/2 gal. or 4 oz./1 1/4 gal.	Apply up to 4 oz. of dilution down midline of back.**
		ULV Spray	1 qt./1 gal. or 1 oz./4 oz.	Mist 1/2 oz. of dilution per animal.**

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a secure area above 32°F.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse container. Triple rinse, then puncture and dispose of container in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Licensed under U.S. Pat. No. 4,062,968 to Sumitomo Chemical Company.