

# Clean-up™ Pour-on Insecticide with IGR

A NOVEL READY-TO-USE SUSPENSION FOR CONTROL OF  
ECTOPARASITES ON LACTATING AND NON-LACTATING DAIRY CATTLE,  
BEEF CATTLE AND CALVES

- Has a zero day pre-slaughter interval and milk discard time.
- May be used as a component of an integrated pest management program, either alone or in combination with other insecticides as an aid to prevent the development of insecticide resistance.
- Dual mode of action delays development of insecticide resistance.

### Active Ingredients

Diflubenzuron [CAS# 35367-38-5] .....5.00%  
 Permethrin [CAS# 52645-53-1].....5.00%

**Other Ingredients\*** .....90.00%  
**Total** .....100.00%

\* Contains petroleum distillates.


SHAKE WELL BEFORE USE

**Keep Out of Reach of Children**

**CAUTION**


See side panel for additional precautionary statements

Net Contents: \_\_ fluid ounces



Boehringer Ingelheim Vetmedica, Inc.  
 St. Joseph, MO 64506-2002

EPA Reg. No. 4691-167  
 EPA Est. No. 4691-KS-01



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

**Personal Protective Equipment (PPE)**

Applicators and handlers must wear:

- 1) Long-sleeved shirt and long pants
- 2) Shoes and socks
- 3) Chemical-resistant gloves

**FIRST AID**

**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-821-7467 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and other aquatic invertebrates. Do not add directly to water, or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**READY TO USE – NO DILUTION NECESSARY**

TARGET ANIMAL	TARGET PEST	APPLICATION INSTRUCTIONS
Lactating and non-lactating dairy cattle, beef cattle and calves	Sucking Lice, Chewing Lice, Houseflies, Stable Flies, Horse Flies, Black Flies, Horn Flies, Deer Flies, Face Flies, and Cattle Ticks.	<b>Pour-on:</b> Apply 3 mL per 100 lbs body weight of animal, up to maximum of 30 mL for any one animal. Pour along back line and down face. Product may be either administered directly from graduated tip-and-pour container, graduated applicator gun, or graduated ladle/dipper. Product is not effective in controlling cattle grubs; therefore, the product can be used on cattle at any time of the year without fear of host-parasite reactions commonly associated with grub treatment products.
Horses	Houseflies, Stable Flies, Horse Flies, Black Flies, Horn Flies, Deer Flies, Face Flies, Sucking Lice, Chewing Lice, and Cattle Ticks.	<b>Pour-on:</b> Apply 3 mL per 100 lbs body weight of animal, up to maximum of 30 mL for any one animal. Pour along back line and down face of <b>MATURE</b> horses. Product may be either administered directly from graduated tip-and-pour container, graduated applicator gun, or graduated ladle/dipper. Do not use this method of application on foals.
Horses and Foals	Houseflies, Stable Flies, Horse Flies, Black Flies, Horn Flies, Deer Flies, Face Flies, Sucking Lice, Chewing Lice, and Cattle Ticks.	<b>Wipe-on:</b> Apply 3 mL per 100 lbs body weight of animal, up to maximum of 30 mL for any one animal. Dampen an applicator mitt, cloth or toweling with Clean-up Pour-on Insecticide with IGR. Rub over hair with special attention to the legs, shoulders, neck and facial areas where pests appear or tend to congregate. Apply directly onto attached ticks.  <b>Ready-To-Use Spray:</b> Use Clean-up Pour-on Insecticide with IGR undiluted in a mist sprayer, applying 3 mL per 100 lbs body weight of animal, up to maximum of 30 mL for any one animal. Direct spray to the legs, shoulders, neck and facial areas where pests appear or tend to congregate. Apply directly onto attached ticks.

**Instructions for all methods of application:** Keep away from eyes and mucous membranes of animal being treated. Repeat treatment as needed, but do not apply more than once every two weeks.

For net contents of 8 oz.:	This container will treat up to 8 animals.
For net contents of 1 pint:	This container will treat up to 16 animals.
For net contents of 1 quart:	This container will treat up to 32 animals.
For net contents of ½ gallon:	This container will treat up to 63 animals.
For net contents of 1 gallon:	This container will treat up to 126 animals.
For net contents of 2.5 gallons:	This container will treat up to 315 animals.
For net contents of 5 gallons:	This container will treat up to 631 animals.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Keep container sealed when not in use. Do not store near food or feed.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For Containers 1 Gallon and Under:** Do not reuse empty container. Rinse thoroughly, wrap in several layers of newspaper, and place in trash.

**WARRANTY AND LIMITATION OF DAMAGES**

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. **Seller makes no other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty.** In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which claim is made.

(Label Date: 1/27/05)