### **GENERAL DIRECTIONS**

Precautions in using: During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11 for protection against Ciodrin insecticide.

#### **FLY CONTROL**

Dairy and Beef Cattle: To control Horn Flies, Face Flies, House Flies, Stable Flies, Lice and the Lone Star Tick: Mix 1 pt. with 6-12 gals. water. Thoroughly apply from 2-3 qts. of diluted spray per animal. Do not apply more often than once per week.

Premise Spray. For Fly Control: Mix 2½ pts. in 12 gals. water. Thoroughly treat at the rate of 1-gal. of diluted spray per 1,000 sq. ft. of surface to aid in reducing populations of house flies and stable flies around barns, on walls, eaves, fences, in loafing sheds and corrals. Repeat later as needed for control.

#### LICE and TICKS

Dairy and Beef Cattle, Sheep, Goats and Hogs: For application in Fall and Winter: Mix 1 pt. in 6-12 gals. water and apply thoroughly to cattle at the rate of 1-gal of diluted spray per animal. Use approximately ½ gal. diluted spray per head on small animals. Repeat application 14 days later.

#### **BACKRUBBER & FACE TREATER USE**

Dairy and Beef Cattle: In back rubbers for Horn Flies, or face treaters for Face Flies: Use a 1% solution by diluting 1 pt. to 2% gals. of fuel oil, kerosene or mineral seal oil. The face rubber mechanism must be so designed that one end is



# CIODRIN®150

## Emulsifiable Spray

For control of Horn Flies, Stable Flies and Face Flies on Beef and Dairy Cattle

ACTIVE INGREDIENTS:	By Weight
Dimethyl phosphate of alpha-methylbe	
3-hydroxycis-crotonate*	
Aromatic Petroleum Hydrocarbons	75.6%
INERT INGREDIENTS:	
Total	100.0%

\*Ciodrin® Insecticide

OLORADO INTERNATIONAL CORPORATION

low enough (not more than 15-20 inches above the ground) to allow cattle to rub their faces. Sheet canvas type face treaters should be located at barn door openings, in loafing areas or around salt licks where animals may freely treat themselves as needed for face fly control. This product is only effective for face fly control on those cattle that rub the face and head on the face treater.

NOTE: Do not contaminate feed or foodstuffs or drinking water. Do not apply under, in or around poultry houses.

WARNING: Hazardous if swallowed, inhaled, or absorbed through skin. Avoid breathing vapors. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing, in case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Call a physician in all cases of suspected poisoning. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ENVIRONMENTAL CAUTION: This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift of spray from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not re-use empty container. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as: weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-119

EPA Est. 4715-CO-1