



25% MALATHION WETTABLE POWDER

ACTIVE INGREDIENT:	By Weight
Malathion: 0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.....	25%
INERT INGREDIENTS:.....	75%
Total.....	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not re-use empty container. Destroy by burning (stay away from smoke or fumes) or bury with waste. Bury ashes at least 18-inches deep in non-cropland away from water supplies.

NET WEIGHT: POUNDS

COLORADO INTERNATIONAL CORP., Commerce City, Colo. 80022

DIRECTIONS FOR USE

Rates given are to be mixed with 100 gallons of water. Contact your Cooperative Agricultural Extension Service for timing of application. Not for use in control of Phosphate resistant Red Spider Mites or Pear Psylla.

APPLES

Aphids: 2-2½ lbs. Make 1 or more applications as needed.

Red Spider Mites: 2-2½ lbs. Make at least 2 applications 10-12 days apart during the summer months. **NOTE:** Do not apply within 3 days of harvest. Malathion may injure McIntosh and Cortland Varieties.

PEARS

Red Spider Mites: 2-2½ lbs. Make 2 applications 10-12 days apart when infestations warrant. Pear Psylla: 2-2½ lbs. Apply 1 or more sprays as needed for control. **NOTE:** Do not apply within 1 day of harvest.

CABBAGE, BROCCOLI, KALE, MUSTARD, TURNIPS

Aphids, Imported Cabbageworms, Cabbage Loopers: 2 lbs. Make full coverage application and repeat as needed for control. **NOTE:** Do not apply within 7 days of harvest on cabbage, kale, and mustard; 3 days for broccoli and turnips.

POTATOES

Aphids, Leafhoppers: 2½ lbs. Make 1 or more applications as needed for control.

TOMATOES

Aphids: 2 lbs. Make 1 or more applications as needed. Spider Mites: 2 lb. Apply as needed. **NOTE:** Do not apply within one day of harvest.

ORNAMENTALS

Lace Bugs, Bagworms: 4 lbs. Tent Caterpillars: 2 lbs. Aphids, Mealybugs, Spider Mites, Whiteflies: 2½ lbs. This spray will also control the Leafhoppers and Thrips that may be present. Soft Brown Scale, Juniper Scale: 6 lbs. Make a thorough application and repeat as necessary. For Scale: Spray after the crawlers have settled on the foliage. **NOTE:** Injury may result to certain ferns and some species of Crassula as well as Canaerti Junipers.

BEEF CATTLE

Lice: 16 lbs. One treatment may be sufficient. Repeat only if needed for control. Horn Fly: 16-24 lbs. Apply a complete coverage spray. Repeat application at 2-week intervals, if needed. **NOTE:** Do not apply to dairy animals. Do not treat calves under one-month of age. Avoid contamination of feed, food, food containers and water troughs.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as: weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.