


1(of 2)



ACCEPTED
AUG 8 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 171-128

4% MALATHION DUST

POULTRY — CATTLE — SWINE AND LISTED VEGETABLES

ACTIVE INGREDIENTS:	BY WEIGHT
Malathion: O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate	4.0%
INERT INGREDIENTS:	96.0%
TOTAL	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or absorbed through the skin. Avoid breathing in dust. Avoid contact with the skin. Wash thoroughly after using. Avoid contamination of feed and food stuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may be a cholinesterase inhibitor.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Shrimp and crayfish may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams, ponds. Keep out of tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bee. Exposed to direct treatment or residual spray. For more information may be obtained from your State Pesticide Applicator's Office. See label.

CONTAINER DISPOSAL: Do not reuse empty container. Dispose of by incineration or by burning. Stay away from open flames. Bury under soil if incineration is not possible. Do not reuse from water supplies.

NET WEIGHT 1 POUNDS

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

DIRECTIONS FOR USE

Begin applications when insects first appear and repeat as required. Dust plants thoroughly. Observe day intervals between first application and harvest indicated by number in () following crop.

ALFALFA (0): Armyworms: Dust plant thoroughly. Make thorough application when larvae are small. Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittle Bugs, Vetch Bruchids: Dust plant thoroughly. Spotted Alfalfa Aphid: Dust plant thoroughly. May be applied on day of harvest or grazing. Note: During bloom apply only in the evening or early morning when bees are not active.

BEANS (1): Mexican Bean Beetle, Leafhoppers, Aphids: Dust plant thoroughly.

BLACKBERRIES (1), BOYSENBERRIES (1), DEWBERRIES (1), LOGANBERRIES (1), RASPBERRIES (1): Aphids, Spider Mites, Thrips, Leafhoppers, Rose Scale: Dust plant thoroughly.

BLUEBERRIES (1): Blueberry Maggots: Dust plant thoroughly.

BRUSSELS SPROUTS (7), CABBAGE (7), KALE (7), TURNIP (5), MUSTARD (7): Aphids, Imported Cabbage Worm, Cabbage Loopers: Dust plant thoroughly. Note: For control of Cabbage Looper, Malathion should be used in combination with other recommended insecticides.

BRUSSELS SPROUTS (7): Aphids: Dust plant thoroughly.

CARROTS (7): Aphids, Leafhoppers: Dust plant thoroughly.

CAULIFLOWER (7): Aphids: Dust plant thoroughly.

CLOVER (0): Aphids, Mites: Dust plant thoroughly. Note: Do not apply to clover in bloom. May be applied on day of harvest or grazing.

CRANBERRIES (3): Leafhoppers, Black Headed Fireworms: Dust plant thoroughly. Note: Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1): Aphids, Spider Mites, Leafhoppers: Dust plant thoroughly. Note: Do not apply on cucurbits unless plants are dry.

GRAPES (3): Drosophila Spp. (Vinegar or grape fly): Dust plant thoroughly. Spider Mites: Dust plant thoroughly.

LETTUCE (Leaf Lettuce 14) (Head Lettuce 7): Aphids, Mites, Cabbage Loopers: Dust plant thoroughly.

ONIONS (Green 3): Thrips: Dust plant thoroughly.

ORNAMENTALS: Aphids, Spider Mites, Bagworms: Dust thoroughly. Note: Applications of this material may cause injury to certain ferns such as Boston Maidenhair and Pteris, as well as on certain species of Brassica, and on foliage of Thuja.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Dust plant thoroughly. Armyworms: Dust plant thoroughly. Note: Make thorough applications when larvae are small and repeat as necessary to maintain control.

PEA (3): Pea Aphid: Dust plant thoroughly. Note: Do not apply within 7 days of harvest if vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Dust plant thoroughly.

SPINACH (7): Aphids: Dust plant thoroughly.

STRAWBERRIES (1): Aphids, Spider Mites, Thrips: Dust plant thoroughly.

TOBACCO (1): Aphids: Dust plant thoroughly.

TOMATO (1): Aphids, Spider Mites, Russet Mites: Dust plant thoroughly.

POULTRY, CATTLE AND SWINE

DIRECT APPLICATION TO BEEF AND DAIRY CATTLE: Horn Fly (1), Eye Flies (1), animal body, neck and neck of the animal. Do not apply to dairy cattle later than 6 hours before milking or during milking. Ideal time for application is immediately after milking. Repeat applications at 10-14 day intervals. Note: Do not treat animals under one year of age when applying dusts. Avoid contamination of feed and water troughs and watering troughs. Do not use on lactating animals.

DIRECT APPLICATION TO BEEF CATTLE AND NON-LACTATING DAIRY CATTLE: Horn Fly (1), Eye Flies (1), animal body, neck and neck of the animal. Do not apply to dairy cattle later than 6 hours before milking or during milking. Ideal time for application is immediately after milking. Repeat applications at 10-14 day intervals. Note: Do not treat animals under one year of age when applying dusts. Avoid contamination of feed and water troughs and watering troughs. Do not use on lactating animals.

POULTRY: Turkey (1), Duck (1), Goose and Turkey, Northern Fowl Mite, Poultry Lice and Chicken Red Mite (1). Do not apply to birds under 10 days of age. Do not apply to waterfowl later than 14 weeks of age. Do not apply to waterfowl later than 14 weeks of age. Do not apply to poultry for Northern Fowl Mite and Chicken Red Mite. Do not apply to waterfowl later than 14 weeks of age. Do not apply to poultry for Northern Fowl Mite and Chicken Red Mite.

PREMISE TREATMENT: Spot treat 10-20 sq. ft. of floor and floor space and 10-20 sq. ft. of walls and ceiling.


SWINE: Lice (1), Fleas (1), and other parasites. Do not apply. Repeat application in 10-14 days.

NOTICE: Always consult the label of this product and refer to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and time of application, may vary, seller makes no claims other than those stated on this label.

EPA Reg. No. 171-128

EPA Est. No. 121-CC-1

1 (of 2)



ACCEPTED
AUG 8 1975
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No. 128

4% MALATHION DUST

POULTRY — CATTLE — SWINE AND LISTED VEGETABLES

ACTIVE INGREDIENTS:	BY WEIGHT
Malathion: O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate	4.0%
INERT INGREDIENTS:	96.0%
TOTAL	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or absorbed through the skin. Avoid breathing or dust. Avoid contact with the skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams or ponds. Keep out of tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from

DIRECTIONS

Begin applications when insects first appear and observe day interval between applications.

ALFALFA (0): Armyworms: Dust plant thoroughly when insects are small. Aphids, Potato Leafhoppers, Spider Mites, Bruchids: Dust plant thoroughly. Spotted Alfalfa on day of harvest or grazing. Note: During bloom when bees are not active.

BEANS (1): Mexican Bean Beetle, Leafhoppers.

BLACKBERRIES (1), BOYSENBERRIES (1), RASPBERRIES (1): Aphids, Spider Mites, Thoroughly.

BLUEBERRIES (0): Blueberry Maggots: Dust thoroughly.

BROCCOLI (3), CABBAGE (7), KALE (7), T. Cabbage Worm, Cabbage Loopers: Dust plant thoroughly. Malathion should be used in combination with other insecticides.

BRUSSELS SPROUTS (7): Aphids: Dust plant thoroughly.

CARROTS (7): Aphids, Leafhoppers: Dust plant thoroughly.

CAULIFLOWER (7): Aphids: Dust plant thoroughly.

CLOVER (0): Aphids, Mites: Dust plant thoroughly. May be applied on day of harvest or grazing.

CRANBERRIES (3): Leafhoppers, Black Headed Crane: Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1): Thoroughly. Note: Do not apply on cucurbits until after harvest.

GRAPES (3): Drosophila Spp. (Vinegar or pom) Dust plant thoroughly.

LETTUCE (Leaf Lettuce 14) (Head Lettuce 7): Thoroughly.

ONIONS (Green 3): Thrips: Dust plant thoroughly.

ORNAMENTALS: Aphids, Spider Mites, Bagworms: this material may cause injury on certain ferns, as on certain species of Crassula, and on Canaertia.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Dust plant thoroughly. Note: Make thorough applications necessary to maintain control.

PEA (3): Pea Aphid: Dust plant thoroughly. No vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Dust plant thoroughly.

SPINACH (7): Aphids: Dust plant thoroughly.

STRAWBERRIES (3): Aphids, Spider Mites: Dust plant thoroughly.

TOBACCO (7): Aphids: Dust plant thoroughly.

TOMATO (1): Aphids, Spider Mites, Russet Mite: Dust plant thoroughly.

POULTRY, CATTLE

DIRECT APPLICATION TO BEEF AND DAIRY ANIMALS: Apply on back and neck of the animal before milking or during milking. Ideal time is before milking.

POULTRY — CATTLE — SWINE AND LISTED VEGETABLES

ACTIVE INGREDIENTS:	BY WEIGHT
Malathion: O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate	4.0%
INERT INGREDIENTS:	96.0%
TOTAL	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with the skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams or ponds. Keep out of tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes. Bury ashes at least 18 inches deep in non-cropland away from water supplies.

NET WEIGHT 1.11 POUNDS
1 1/4

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

BRUSSELS SPROUTS (7): Aphids: Dust plant thoroughly.

CARROTS (7): Aphids, Leafhoppers: Dust plant thoroughly.

CAULIFLOWER (7): Aphids: Dust plant thoroughly.

CLOVER (0): Aphids, Mites: Dust plant thoroughly. May be applied on day of harvest or grazing.

CRANBERRIES (3): Leafhoppers, Black Hops: Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1): Apply thoroughly. **Note:** Do not apply on cucurbits.

GRAPES (3): Drosophila Spp. (Vinegar or Pinworm): Dust plant thoroughly.

LETTUCE (Leaf Lettuce 14) (Head Lettuce 14): Dust plant thoroughly.

ONIONS (Green 3): Thrips: Dust plant thoroughly.

ORNAMENTALS: Aphids, Spider Mites, Black Flies: This material may cause injury on certain ferns, as on certain species of Crassula, and on Cana.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Dust plant thoroughly. **Note:** Make thorough application necessary to maintain control.

PEA (3): Pea Aphid: Dust plant thoroughly. Vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Dust plant thoroughly.

SPINACH (7): Aphids: Dust plant thoroughly.

STRAWBERRIES (3): Aphids, Spider Mites: Dust plant thoroughly.

TOBACCO (7): Aphids: Dust plant thoroughly.

TOMATO (1): Aphids, Spider Mites, Russet Mites: Dust plant thoroughly.

POULTRY, CA

DIRECT APPLICATION TO BEEF AND SWINE: Apply on back and neck of the animal before milking or during milking. Ideal time for application is 10-14 day intervals. **Note:** When applying dusts avoid contamination of feed. Do not use spray or dip dairy animals in malathion.

DIRECT APPLICATION TO BEEF CATTLE: Treat animal thoroughly. Repeat in 10 days per animal. Apply on back and neck at 10-14 day intervals. **Note:** Do not apply to lactating dairy cows during 2 weeks of freshening.

POULTRY: (Chickens, Ducks, Geese and Turkeys) **Chicken Red Mite:** Dust individual birds by net application in 4-8 weeks or when necessary. **Chicken Red Mite:** should be a supplemental treatment. **PREMISE TREATMENT:** Apply 1 lb. 50 g per roost and adjacent areas.

SWINE: Lice. Treat animals, pens and litter thereafter as needed.

NOTICE: Seller warrants that this product conforms to the label. Since conditions of use, such as weather, and application equipment will vary, Seller makes no

ACCEPTED

AUG 8 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, recommended for the pesticide registered under FIFRA-128

DUST

SWINE
BLES

BY WEIGHT

..... 4.0%
..... 96.0%
100.0%

OF CHILDREN

ing of dust. Avoid contact of feed and foodstuffs. Keep

on.

imp and crab may be killed where these are important marshes and estuaries. Do not conditions favor drift from

DIRECTIONS FOR USE

Begin applications when insects first appear and repeat as required. Dust plants thoroughly. Observe day interval between last application and harvest indicated by number in () following crop.

ALFALFA (0): Armyworms: Dust plant thoroughly. Make thorough application when larvae are small. **Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittle Bugs, Vetch Bruchids:** Dust plant thoroughly. **Spotted Alfalfa Aphid:** Dust plant thoroughly. May be applied on day of harvest or grazing. **Note:** During bloom apply only in the evening or early morning when bees are not active.

BEANS (1): Mexican Bean Beetle, Leafhoppers, Aphids: Dust plant thoroughly.

BLACKBERRIES (1), BOYSENBERRIES (1), DEWBERRIES (1), LOGANBERRIES (1), RASPBERRIES (1): Aphids, Spider Mites, Thrips, Leafhoppers, Rose Scale: Dust plant thoroughly.

BLUEBERRIES (0): Blueberry Maggots: Dust plants thoroughly.

BROCCOLI (3), CABBAGE (7), KALE (7), TURNIP (3), MUSTARD (7): Aphids, Imported Cabbage Worm, Cabbage Loopers: Dust plant thoroughly. **Note:** For control of Cabbage Loopers Malathion should be used in combination with other recommended insecticides.

BRUSSELS SPROUTS (7): Aphids: Dust plant thoroughly.

CARROTS (7): Aphids, Leafhoppers: Dust plant thoroughly.

CAULIFLOWER (7): Aphids: Dust plant thoroughly.

CLOVER (0): Aphids, Mites: Dust plant thoroughly. **Note:** Do not apply to clover in bloom. May be applied on day of harvest or grazing.

CRANBERRIES (3): Leafhoppers, Black Headed Fireworms: Dust plant thoroughly. **Note:** Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1): Aphids, Spider Mites, Leafhoppers: Dust plant thoroughly. **Note:** Do not apply on cucurbits unless plants are dry.

GRAPES (3): Drosophila Spp. (Vinegar or pomace fly): Dust plant thoroughly. **Spider Mites:** Dust plant thoroughly.

LETTUCE (Leaf Lettuce 14) (Head Lettuce 7): Aphids, Mites, Cabbage Loopers: Dust plant thoroughly.

ONIONS (Green 3): Thrips: Dust plant thoroughly.

ORNAMENTALS: **Aphids, Spider Mites, Bagworms:** Dust thoroughly. **Note:** Applications of this material may cause injury on certain ferns, such as Boston Maidenhair and Pteris, as well as on certain species of Crassula, and on Galearia Junipers.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Dust plant thoroughly. **Armyworms:** Dust plant thoroughly. **Note:** Make thorough applications when larvae are small and repeat as necessary to maintain control.

PEA (3): Pea Aphid: Dust plant thoroughly. **Note:** Do not apply within 7 days of harvest if vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Dust plant thoroughly.

SPINACH (7): Aphids: Dust plant thoroughly.

STRAWBERRIES (3): Aphids, Spider Mites: Dust plant thoroughly. **Thrips:** Dust plant thoroughly.

TOBACCO (1): Aphids: Dust plant thoroughly.

TOMATO (1): Aphids, Spider Mites, Russet Mites: Dust plant thoroughly.

POULTRY, CATTLE AND SWINE

DIRECT APPLICATION TO BEEF AND DAIRY CATTLE: Horn Fly: Use 1 Tbsp. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Ideal time for application is immediately after milking.

FILE — SWINE VEGETABLES

BY WEIGHT

ate of diethyl	4.0%
.....	96.0%
TOTAL	100.0%

REACH OF CHILDREN

Avoid breathing of dust. Avoid contact with contamination of feed and foodstuffs. Keep

sterase inhibition.

Toxic to fish. Shrimp and crab may be killed. Do not apply where these are important. Do not use near out of tidal marshes and estuaries. Do not use when weather conditions favor drift from application. Clean, store, and dispose of equipment, or disposal of wastes.

Do not apply to treatment or residues on crops. Protective clothing. Agricultural Extension Service.

Destroy it by burying with waste. Bury at least 18 inches deep in non-crop land.

POUNDS

IAL CORPORATION
orado 80022

- BROSSELS SPROUTS (7): Aphids:** Dust plant thoroughly.
- CARROTS (7): Aphids, Leafhoppers:** Dust plant thoroughly.
- CAULIFLOWER (7): Aphids:** Dust plant thoroughly.
- CLOVER (0): Aphids, Mites:** Dust plant thoroughly. **Note:** Do not apply to clover in bloom. May be applied on day of harvest or grazing.
- CRANBERRIES (3): Leafhoppers, Black Headed Fireworms:** Dust plant thoroughly. **Note:** Do not apply to wet cranberry vines or foliage.
- CUCUMBERS (1), SQUASH (1), MELONS (1): Aphids, Spider Mites, Leafhoppers:** Dust plant thoroughly. **Note:** Do not apply on cucurbits unless plants are dry.
- GRAPES (3): Drosophila Spp. (Vinegar or pomace fly):** Dust plant thoroughly. **Spider Mites:** Dust plant thoroughly.
- LETTUCE (Leaf Lettuce 1-1) (Head Lettuce 7): Aphids, Mites, Cabbage Loopers:** Dust plant thoroughly.
- ONIONS (Green 3): Thrips:** Dust plant thoroughly.
- ORNAMENTALS: Aphids, Spider Mites, Bagworms:** Dust thoroughly. **Note:** Applications of this material may cause injury on certain ferns, such as Boston, Maidenhair and Pteris, as well as on certain species of Crassula, and on Canaert Junipers.
- GRASS AND GRASS HAY (0): Aphids, Leafhoppers:** Dust plant thoroughly. **Armyworms:** Dust plant thoroughly. **Note:** Make thorough applications when larvae are small and repeat as necessary to maintain control.
- PEA (3): Pea Aphid:** Dust plant thoroughly. **Note:** Do not apply within 7 days of harvest if vines are to be fed to animals.
- POTATOES (0): Aphids, Leafhoppers:** Dust plant thoroughly.
- SPINACH (7): Aphids:** Dust plant thoroughly.
- STRAWBERRIES (3): Aphids, Spider Mites:** Dust plant thoroughly. **Thrips:** Dust plant thoroughly.
- TOBACCO (7): Aphids:** Dust plant thoroughly.
- TOMATO (1): Aphids, Spider Mites, Russet Mites:** Dust plant thoroughly.

POULTRY, CATTLE AND SWINE

DIRECT APPLICATION TO BEEF AND DAIRY CATTLE: Horn Fly: Use 4 Tbsp. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Ideal time for application is immediately after milking. Repeat applications at 10-14 day intervals. **Note:** Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs. Do not use spray or dip dairy animals for malatation.

DIRECT APPLICATION TO BEEF CATTLE AND NON-LACTATING DAIRY CATTLE: Lice: Treat animal thoroughly. Repeat in 10 days and thereafter as needed. **Horn Fly:** Use 4 Tbsp. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals. **Note:** Do not apply to lactating dairy animals or non-lactating dairy animals with 2 weeks of freshening.

POULTRY: (Chickens, Ducks, Geese and Turkeys): **Northern Fowl Mite, Poultry Lice and Chicken Red Mite:** Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4-8 weeks or when necessary. Treatment of poultry for Northern Fowl Mite and Chicken Red Mite should be a supplemental treatment to premise treatment.

PREMISE TREATMENT: Apply 1 lb. 50-60 sq. ft. of floor and floor space and 1/2 parts, roosts and adjacent areas.


SWINE: Lice: Treat animals, pens and litter thoroughly. Repeat application in 10 days and thereafter as needed.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-128

EPA Est. 4715-CO-1

2(0f2)



4% MALATHION DUST

Poultry - Cattle - Swine
and Listed Vegetables

ACTIVE INGREDIENTS:	By Weight
Malathion (O,O dimethyl dithiophosphate of diethyl mercaptosuccinate)	4.0%
INERT INGREDIENTS:	96.0%
Total	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or absorbed through the skin. Avoid breathing of dust. Avoid contact with the skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. ATROPINE is antidotal.

ENVIRONMENTAL CAUTION: This product is toxic to fish, Shrimp, and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams or ponds. Keep out of tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes; bury ashes at least 18 inches deep in non cropland away from water supplies.

NET WEIGHT: POUNDS

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

DIRECTIONS FOR USE

See application when insect first appears and repeat as required. Dust plants thoroughly. Observe days interval between last application and harvest indicated by label for each crop.

ALFALFA (0): Armyworms: Apply 4 lbs./acre. Make thorough application when larvae are small. Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Little Bluet, Vetch Bruchids: Apply 3 lbs./acre.

SPOTTED ALFALFA APHID: Ground Equipment: Apply 15 lbs./acre. Air Equipment: Apply 25 lbs./acre. Note: Do not apply to alfalfa to be grazed. Do not apply to alfalfa to be baled.

BEANS (1): Mexican Bean Beetle, Leafhoppers, Aphids: Apply 1 lb./acre.

BLACKBERRIES (1), BOYSENBERRIES (1), DEWBERRIES (1), LOGANBERRIES (1), RASPBERRIES (1): Aphids, Spider Mites, Thrips, Leafhoppers, Rose Scale: Apply 40 lbs./acre. On raspberries apply 45 lbs./acre.

BLUEBERRIES (0): Blueberry Maggots: Apply 25 lbs./acre by ground or air equipment.

BROCCOLI (2), CABBAGE (2), KALE (1), TURNIP (1), MUSTARD (1): Aphids, Imported Cabbage Worm, Cabbage Loopers: Apply 40 lbs./acre. Note: For control of Cabbage Loopers, Malathion should be used in combination with other recommended insecticides.

BRUSSEL SPROUTS (7): Aphids: Apply 30 lbs./acre.

CARROTS (1): Aphids, Leafhoppers: Apply 40 lbs./acre.

CAULIFLOWER (7): Aphids: Apply 50 lbs./acre.

CLOVER (0): Aphids, Mites: Apply 25-30 lbs./acre. Note: Do not apply to clover in bloom. May be applied on the ground or grazing.

CRANBERRIES (2): Leafhoppers, Black Headed Fireworms: Apply 30-35 lbs./acre (50 lbs. by aircraft). Note: Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1): Aphids, Spider Mite, Leafhoppers: Apply 25-35 lbs./acre. Note: Do not apply on cucurbits if cucurbit are dry.

GRAPES (3): Drosophila Spp. (Vinegar Flies), Spider Mites: Apply 25-30 lbs./acre. Note: Do not apply on grapes if grapes are to be fed to animals.

LETTUCE (Leaf Lettuce (1) (Head Lettuce (1)): Aphids, Mites, Cabbage Loopers: Apply 20-30 lbs./acre.

ONIONS (Green (3)): Thrips: Apply 10 lbs./acre.

ORNAMENTALS: Aphids, Spider Mites, Bagworms: Dust thoroughly. Note: Applications of this material may cause injury on certain ferns, such as Boston Maidenhair and Pteris, as well as on certain species of Chussea, and on Carolina Junipers.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Apply 30-40 lbs./acre. Armyworms: Apply 25-40 lbs./acre. Note: Make thorough applications when larvae are small and repeat as necessary to maintain control.

PEAS (3): Pea Aphid: Apply 25 lbs./acre. Note: Do not apply within 7 days of harvest if vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Apply 25-30 lbs./acre.

SPINACH (7): Aphids: Apply 30-35 lbs./acre.

STRAWBERRIES (3): Aphids, Spider Mites: Apply 40 lbs./acre. Thrips: Apply 20 lbs./acre.

TOBACCO (7): Aphids: Apply 25-30 lbs./acre. Note: Use maximum dosage on plant buds.

TOMATO (1): Aphids, Spider Mites, Russet Mites: Apply 25-35 lbs./acre.

POULTRY, CATTLE AND SWINE

DIRECT APPLICATION TO BEEF AND DAIRY CATTLE - Horn Fly: Use 4 Tbsp. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 14 days before milking or during milking. Repeat application is indicated after milking. Repeat application at 10-14 day intervals. Note: Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs. Do not use spray or dip during lactation.

DIRECT APPLICATION TO BEEF CATTLE AND NON LACTATING DAIRY CATTLE - Lice: Treat animal thoroughly. Repeat 10 days and thereafter as needed. Horn Fly: Use 3 Tbsp. per animal. Apply on the back and neck of the animal and repeat applications at 10-14 day intervals. Note: Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

POULTRY (Chickens, Ducks, Geese and Turkeys): Northern Fowl Mite, Poultry Lice and Chicken Red Mite: Dust individual birds by means of a shaker, tin or rotary hand duster. Repeat application in 4-8 weeks or when necessary. Treatment of poultry for Northern Fowl Mite and Chicken Red Mite should be a supplemental treatment to premise treatment.

PREMISE TREATMENT: Apply 1 lb./50 sq. ft. of floor and floor space and to nests, roosts and adjacent areas.

SWINE: Lice: Treat animals, pens and litter thoroughly. Repeat application in 10 days and thereafter as needed.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, seller makes no claim other than those stated on this label.

EPA Reg. No. 4715-128
EPA Est. 4715 CO-1

2(0f2)



Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, this pesticide registered under EPA Reg. No. 4715-128

4% MALATHION DUST

Poultry - Cattle - Swine
and Listed Vegetables

ACTIVE INGREDIENTS:	By Weight
Malathion: O,O dimethyl dithiophosphate of diethyl mercaptosuccinate	4.0%
INERT INGREDIENTS:	96.0%
Total	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or absorbed through the skin. Avoid breathing of dust. Avoid contact with the skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. ATROPINE is antidotal.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams or

DI
Begin applications when insects first
Observe days interval between last ap
ing crop.

ALFALFA (0): Armyworms: App
are small. **Aphids, Potato Leafhop**
Vetch Bruchids: Apply 30 lbs./acre.
Spotted Alfalfa Aphid: Ground Equi
12-20 lbs./acre. Use the higher dose
or grazing.

Note: During bloom apply only in
BEANS (1): Mexican Bean Beetle, L

BLACKBERRIES (1), BOYSENBERR
RASPBERRIES (1): Aphids, Spide
lbs./acre. On raspberries, apply 45 lb

BLUEBERRIES (0): Blueberry Ma

BROCCOLI (3), CABBAGE (7): KAL
Cabbage Worm, Cabbage Loopers: A
Note: For control of Cabbage Loope
recommended insecticides.

BRUSSEL SPROUTS (7): Aphids:

CARROTS (2): Aphids, Leafhopper

CAULIFLOWER (7): Aphids: Appl

CLOVER (0): Aphids, Mites: Appl
Note: Do not apply to clover in l

CRANBERRIES (2): Leafhoppers, B
by aircraft.)

Note: Do not apply to wet cranberri

CUCUMBERS (1), SQUASH (1), ME
30-35 lbs./acre.

Note: Do not apply on cucurbits un

GRAPES (3): Drosophila Spp (V
Mites: Apply 20-40 lbs./acre.

LETTUCE (Leaf Lettuce 14) (Head
20-40 lbs./acre.

ONIONS (Green 3): Thrips: Apply

ORNAMENTALS: Aphids, Spider M

Note: Applications of this materi
Maidenhair and Pteris, as well as or

GRASS AND GRASS HAY (0): Aph
Apply 35-40 lbs./acre.

Note: Make thorough applications
rain control.

PEAS (3): Pea Aphid: Apply 25 lbs
Note: Do not apply within 7 days o

POTATOES (0): Aphids, Leafhoppi

SPINACH (7): Aphids: Apply 30-3

STRAWBERRIES (3): Aphids, Sp
lbs./acre.

TOBACCO (7): Aphids: Apply 25-
Note: Use maximum dosage in plant

TOMATO (1): Aphids, Spider Mites

and Listed Vegetables

ACTIVE INGREDIENTS:	By Weight
Malathion: O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate	4.0%
INERT INGREDIENTS:	96.0%
Total	100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or absorbed through the skin. Avoid breathing of dust. Avoid contact with the skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. ATROPINE is antidotal.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams or ponds. Keep out of tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL: Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes; bury ashes at least 18-inches deep in non-cropland away from water supplies.

NET WEIGHT: POUNDS

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

CRANBERRIES (3): Leaf (by aircraft.)
Note: Do not apply to water.

CUCUMBERS (1), SQUASH (1): 30-35 lbs./acre.
Note: Do not apply on cucumbers.

GRAPES (3): Drosophila Mites: Apply 20-40 lbs./acre.

LETTUCE (Leaf Lettuce): 30-40 lbs./acre.

ONIONS (Green 3): Thrips

ORNAMENTALS: Aphids
Note: Applications of this product to Maidenhair and Pteris, as well as other ferns.

GRASS AND GRASS HAY: Apply 35-40 lbs./acre.
Note: Make thorough application for pest control.

PEAS (3): Pea Aphid: Apply 20-40 lbs./acre.
Note: Do not apply within 14 days of harvest.

POTATOES (0): Aphids, Colorado Potato Beetle

SPINACH (7): Aphids: Apply 20-40 lbs./acre.

STRAWBERRIES (3): Apply 20-40 lbs./acre.

TOBACCO (7): Aphids: Apply 20-40 lbs./acre.
Note: Use maximum dosage.

TOMATO (1): Aphids, Spider Mites

DIRECT APPLICATION: animal. Apply on back and before milking or during milking. Repeat applications at 10-day intervals.
Note: Do not treat animal with malathion if feed, food contains malathion.


DIRECT APPLICATION: Lice: Treat animal thoroughly. 3 Tbsp. per animal. Apply every 3-4 days.
Note: Do not apply to animal during freshening.

POULTRY (Chickens, Ducks, Geese): Chicken Red Mite: Dust with Malathion Mite and Chicken Red Mite. Apply to roosts and adjacent areas.

SWINE: Lice: Treat animal with Malathion thereafter as needed.

NOTICE: Seller warrants that this product conforms to the label. Since conditions of application vary, the user should read the label on this label.

EPA Reg. No. 4715-128



Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, this pesticide registered under EPA Reg. No. **4715-128**

Swine
Vegetables

	By Weight
.....	4.0%
.....	96.0%
Total	100.0%

OF REACH OF CHILDREN

h the skin. Avoid breathing of dust.
thoroughly after using. Avoid
Keep out of reach of children and

may cause cholinesterase inhibition

product is toxic to fish. Shrimp and
recommended on this label. Do not

DIRECTIONS FOR USE

Begin applications when insects first appear and repeat as required. Dust plants thoroughly. Observe days interval between last application and harvest indicated by number in () following crop.

ALFALFA (0): Armyworms: Apply 35 lbs./acre. Make thorough application when larvae are small. **Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittle Bugs, Vetch Bruchids:** Apply 30 lbs./acre.

Spotted Alfalfa Aphid: Ground Equipment - Apply 12-15 lbs./acre. Air equipment: Apply 12-20 lbs./acre. Use the higher dosage on vetch alfalfa. May be applied on day of harvest or grazing.

Note: During bloom apply only in the evening or early morning when bees are not active.

BEANS (1): Mexican Bean Beetle, Leafhoppers, Aphids: Apply 30-35 lbs./acre.

BLACKBERRIES (1), BOYSENBERRIES (1), DEWBERRIES (1), LOGANBERRIES (1), RASPBERRIES (1): Aphids, Spider Mites, Thrips, Leafhoppers, Rose Scale: Apply 40 lbs./acre. On raspberries, apply 45 lbs./acre.

BLUEBERRIES (0): Blueberry Maggots: Apply 25 lbs./acre by ground or air equipment.

BROCCOLI (3), CABBAGE (7) KALE (7), TURNIP (3), MUSTARD (7): Aphids, Imported Cabbage Worm, Cabbage Loopers: Apply 30 lbs./acre.

Note: For control of Cabbage Loopers, Malathion should be used in combination with other recommended insecticides.

BRUSSEL SPROUTS (7): Aphids: Apply 30 lbs./acre.

CARROTS (7): Aphids, Leafhoppers: Apply 30-40 lbs./acre.

CAULIFLOWER (7): Aphids: Apply 50 lbs./acre.

CLOVER (0): Aphids, Mites: Apply 25-30 lbs./acre.

Note: Do not apply to clover in bloom. May be applied on day of harvest or grazing.

CRANBERRIES (2): Leafhoppers, Black Headed Fireworms: Apply 30-35 lbs./acre. (50 lbs. by aircraft.)

Note: Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1): Aphids, Spider Mites, Leafhoppers: Apply 30-35 lbs./acre.

Note: Do not apply on cucurbits unless plants are dry.

GRAPES (3): Drosophila Spp (Vinegar or pomace fly): Apply 25-30 lbs./acre. **Spider Mites:** Apply 20-40 lbs./acre.

LETTUCE (Leaf Lettuce 14) (Head Lettuce 7): Aphids, Mites, Cabbage Loopers: Apply 30-40 lbs./acre.

ONIONS (Green 3): Thrips: Apply 30-40 lbs./acre.

ORNAMENTALS: Aphids, Spider Mites, Bagworms: Dust thoroughly.

Note: Applications of this material may cause injury on certain ferns, such as Boston, Maidenhair and Pteris, as well as on certain species of Crassula, and on Canaerti Junipers.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Apply 30-40 lbs./acre. **Armyworms:** Apply 35-40 lbs./acre.

Note: Make thorough applications when larvae are small and repeat as necessary to maintain control.

PEAS (3): Pea Aphid: Apply 25 lbs./acre.

Note: Do not apply within 7 days of harvest if vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Apply 25-30 lbs./acre.

SPINACH (7): Aphids: Apply 30-35 lbs./acre.

STRAWBERRIES (3): Aphids, Spider Mites: Apply 40 lbs./acre. **Thrips:** Apply 25-50 lbs./acre.

TOBACCO (7): Aphids: Apply 25-30 lbs./acre.

Note: Use maximum dosage in plant beds.

TOMATO (1): Aphids, Spider Mites, Russet Mites: Apply 35-45 lbs./acre.

Vegetables

By Weight

phosphate of diethyl	4.0%
.....	96.0%
Total	100.0%

OF REACH OF CHILDREN

h the skin. Avoid breathing of dust.
h thoroughly after using. Avoid
. Keep out of reach of children and

may cause cholinesterase inhibition.

product is toxic to fish. Shrimp and
recommended on this label. Do not
urces. Keep out of lakes, streams or
estuaries. Do not apply where runoff
weather conditions favor drift from
water by cleaning of equipment, or
uct only as specified on this label.

posed to direct treatment or residues
be obtained from your Cooperative

-use empty container. Destroy it by
y away from smoke or fumes; bury
cropland away from water supplies.

POUNDS

TIONAL CORPORATION
, Colorado 80022

CRANBERRIES (3): Leafhoppers, Black Headed Fireworms: Apply 30-35 lbs./acre. (50 lbs. by aircraft.)

Note: Do not apply to wet cranberry vines or foliage.

CUCUMBERS (1), SQUASH (1), MELONS (1) Aphids, Spider Mites, Leafhoppers: Apply 30-35 lbs./acre.

Note: Do not apply on cucurbits unless plants are dry.

GRAPES (3): Drosophila Spp. (Winegar or pomace fly): Apply 25-30 lbs./acre. **Spider Mites:** Apply 20-40 lbs./acre.

LETTUCE (Leaf Lettuce 14) (Head Lettuce 7): Aphids, Mites, Cabbage Loopers: Apply 30-40 lbs./acre.

ONIONS (Green 3): Thrips: Apply 30-40 lbs./acre.

ORNAMENTALS: Aphids, Spider Mites, Bagworms: Dust thoroughly.

Note: Applications of this material may cause injury on certain ferns, such as Boston, Maidenhair and Pteris, as well as on certain species of Crassula, and on Canaerti Junipers.

GRASS AND GRASS HAY (0): Aphids, Leafhoppers: Apply 30-40 lbs./acre. **Armyworms:** Apply 35-40 lbs./acre.

Note: Make thorough applications when larvae are small and repeat as necessary to maintain control.

PEAS (3): Pea Aphid: Apply 25 lbs./acre.

Note: Do not apply within 7 days of harvest if vines are to be fed to animals.

POTATOES (0): Aphids, Leafhoppers: Apply 25-30 lbs./acre.

SPINACH (7): Aphids: Apply 30-35 lbs./acre.

STRAWBERRIES (3): Aphids, Spider Mites: Apply 40 lbs./acre. **Thrips:** Apply 25-50 lbs./acre.

TOBACCO (7): Aphids: Apply 25-30 lbs./acre.

Note: Use maximum dosage in plant beds.

TOMATO (1): Aphids, Spider Mites, Russet Mites: Apply 35-45 lbs./acre.

POULTRY, CATTLE AND SWINE

DIRECT APPLICATION TO BEEF AND DAIRY CATTLE: Horn Fly: Use 4 Tbsp. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Ideal time for application is immediately after milking. Repeat applications at 10-14 day intervals.

Note: Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs. Do not use spray or dip dairy animals in malathion.

DIRECT APPLICATION TO BEEF CATTLE AND NON-LACTATING DAIRY CATTLE: Lice: Treat animal thoroughly. Repeat in 10 days and thereafter as needed. **Horn Fly:** Use 3 Tbsp. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals.

Note: Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

POULTRY (Chickens, Ducks, Geese and Turkeys): Northern Fowl Mite, Poultry Lice and Chicken Red Mite: Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4-8 weeks or when necessary. Treatment of poultry for Northern Fowl Mite and Chicken Red Mite should be a supplemental treatment to premise treatment.

PREMISE TREATMENT: Apply 1 lb./50-60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas.

SWINE: Lice: Treat animals, pens and litter thoroughly. Repeat application in 10 days and thereafter as needed.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-128

EPA Est. 4715 CO-1