

DIRECTIONS FOR USE

**FLY MAGGOTS:** To control fly maggots in such maggot breeding areas as garbage dumps and manure piles, mix 1 quart of DDVP 15% Concentrate with 7-1/2 gallons of water and apply the diluted solution in a coarse, wet spray. Apply one quart of the diluted solution to each 100 square feet.

**ANIMAL BUILDINGS:** To control flies, mosquitoes, roaches, ants and sowbugs in Horse Barns, Dairy Barns, Shelter Sheds, Milk Sheds and Feedlots, mix 1 quart of DDVP 15% Concentrate with 7-1/2 gallons of water and apply the diluted solution to each 1000 square feet. Treat around doorways, feed storage rooms, alleyways and window sills. Animals may be present during treatment. Do not contaminate feed, drinking water, milk or milking utensils.

**DAIRY AND BEEF CATTLE:** For control of House Fly, Horn Fly, Stable Fly, Face Fly, and mosquitoes mix one gallon of 15% Concentrate with 15 gallons water. Apply 2 fluid ounces of this diluted solution per animal as a mist spray daily with hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply in excess of 2 fluid ounces per day.

**IN TREATING DOG KENNELS:** Apply to outside dog runways, window sills and ledges. For control of fleas and grown dog ticks, mix one quart of DDVP 15% Concentrate with 7-1/2 gallons water and spray one quart of this diluted solution to each 500 square feet.

**WARNING:** Poisonous if swallowed, inhaled or absorbed through skin or eyes; Do not get on skin, in eyes, or on clothing. Do not breathe fumes. Avoid breathing of spray mist. In case of contact, remove contaminated clothing and wash skin thoroughly with soap and water. For eyes, flush with plenty of water, get medical attention. If poisoning should occur, CALL A PHYSICIAN IMMEDIATELY. ATROPINE IS ANTIDOTAL. Wear natural rubber gloves, protective clothing and goggles when handling concentrate to prepare emulsions. During fogging or mist spraying applications, wear a mask or respirator approved by the U. S. Dept. of Agriculture for use with DDVP. Wash thoroughly after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Do not contaminate feed, foodstuffs or drinking water. Do not use inside homes or where milk is stored or processed.

This product is toxic to fish and wildlife. Keep out of lakes,

# DDVP 15% CONCENTRATE

FOR CONTROL OF FLIES, GNATS, MOSQUITOES,  
MAGGOTS, ROACHES, FLEAS, ANTS, SOWBUGS,  
BROWN DOG TICKS

DANGER



POISON

## KEEP OUT OF REACH OF CHILDREN

See Antidote Statement and other required warning statements on the side panels.

ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate (DDVP)	13.95%*
Related Compounds	1.05%*
Xylene	80.00%
INERT INGREDIENTS:	5.00%
Total	100.00%

\* Equivalent to 15% of DDVP Insecticide

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022  
ACCEPTED

AUG. 8, 1969

streams or ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.



ANTIDOTE AND FIRST AID TREATMENT: Atropine is the emergency antidote for DDVP poisoning. Consult your physician about obtaining an adequate supply of 1/100 grain Atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100 grain Atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See Warning Symptoms Below.) Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.