



READ ENTIRE LABEL
FOLLOW DIRECTIONS CAREFULLY

GENERAL INFORMATION: When used as directed this product controls many annual and perennial weeds in: corn, wheat, sorghum, barley, rye, oats, roadsides, pastures and turf/lawn areas. It is more effective when used on small, actively growing weeds. Use low-pressure equipment (40 lbs.) on crops and turf. Garden-type pressure sprayers work well on spot spraying. Double-check your equipment to be sure it is properly calibrated: improper tractor speed, pump pressure, nozzle tip size, or rusty equipment will result in poor weed kill or possibly heavy crop damage. Use separate spray equipment for insecticides and fungicides. Under certain high temperature conditions, vapors from this product will injure nearby susceptible plants. Apply when temperatures range from 70-90°F. Coarse sprays are less likely to drift. Do NOT make aerial applications near cotton or sensitive plants such as: vegetables, legumes, grapes or flowers. Consult your local Cooperative Agricultural Extension Service for additional information consistent with label directions and cautions.

DIRECTIONS FOR USE

Use 5-20 gallons of water per acre for uniform spraying. Do not apply if rain is expected within the hour.

SUSCEPTIBLE WEEDS (Use lower dosage rate):

Arrowweed	Lambsquarter	Ragweed
Buckhorn	Mustard	Sunflower
Cocklebur	Morning Glory	Wild Carrot
Dandelion	Pigweed	Wild Radish
Dock	Plantain	Wild Sweet Potato

HARD-TO-KILL WEEDS (Use higher dosage rate):

Bindweed	Hoary Cress	Milkweed
Buckbrush	Horse Nettle	Sheep Sorrel
Buckwheat	Locoweed	Wild Onion
Canada Thistle	Milkweed	Willow

CORN (Post-Emergence): May be applied at several stages of growth. Use drop nozzles when corn is 10 inches or over. Mix 1 cupful in 3 gals. water and wet weeds thoroughly. **NOTE:** Do not apply from tasseling to dough stage.

Early Spraying: (Corn is 2-4 inches high) Apply as soon as possible after most weeds have emerged. Use 1 pt. acre in sufficient water to give uniform coverage. Corn drop nozzles are not necessary at this time.



AMINE 400 2,4-D WEED KILLER

Contains: 4 lbs. 2,4-D Acid per gallon.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic acid*

INERT INGREDIENTS:

Total
*2, 4 Dichlorophenoxyacetic Acid equivalent . . . 41.0%

By Weight

49.3

50.7

100.0

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact, wash with soap and water. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs. Do not store near seed, feed, insecticides, fungicides or fertilizers. Keep out of reach of children and domestic animals.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

ENVIRONMENTAL CAUTION: Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Rinse equipment & dispose of wastes by burying in non cropland away from domestic water supplies. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not re-use empty container; destroy by perforating or crushing, and, bury with wastes.

NET CONTENTS 4.5 GALLONS

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

Lay By Spraying: (Corn is 10 inches high) Apply as soon as possible after most weeds have emerged. Use 1 pt. acre in sufficient water to give uniform coverage. Corn drop nozzles are not necessary at this time. **NOTE:** Do not apply from tasseling to dough stage.

High-Clearance Spraying: This spray is most useful when weeds get started after Lay By. Use 1 pt./acre; adjust spray nozzles to hit highest weeds.

SORGHUM (Post-Emergence ONLY): Use Drop nozzles. Sorghums are most tolerant to 2, 4 D when 4-12 inches high. Spray a check plot first. Use 1 pt. acre. **NOTE:** Do not apply during flowering or early dough stage.

HARD-TO-KILL WEEDS in NON-CROPLAND: For perennial weeds, such as Sow Thistle and Russian Knapweed (see list) in drainage ditchbanks, fence rows and farm roadsides. Use 2-3 qts. in 20 gals. water with 5-8 oz. liquid detergent; thoroughly wet weed foliage. Treat in Spring and again at bloom time before weeds go to seed. Repeat the following year. **NOTE:** Do not let spray drift.

SMALL GRAINS (WHEAT, BARLEY, RYE, OATS)

Winter Wheat and Rye. Treat only in the Spring: Use 1 1/2 pts. in sufficient water to cover one acre. Apply from the tillered to the jointing stage. **Spring Wheat, Oats and Barley:** Use 1 - 1 1/3 pt./acre. Use higher dosage for hard to kill weeds. Treat when crop is fully tillered (4-8 inches high) up until early-boot stage. **NOTE:** Do not spray during seedling stage, or, boot, to dough stage.

PASTURES and TURF: Use 2-3 pts. in 10 gals. water/acre. Apply early in Summer when plants are actively growing. Treat again in bud stage before weeds go to seed. **NOTE:** Remove cows from dairy pastures during spray application and do not graze treated areas for 7 days.

LAWNS: Use 1 Tbsp. gal. for 1,000 sq. ft. Repeat as needed for control. **NOTE:** Do not spray new lawns until after first cutting. Do NOT use on: Bentgrass, St. Augustine grass, Dichondra, or, Carpetgrass lawns.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment may vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-168 EPA F. 4715 CO-1