### DIRECTIONS FOR CLEANING UTENSILS AND EQUIPMENT:

- 1. Scrape and prewash dishes, glasses, utensils and equipment whenever possible.
- 2. Wash with Sani-Udder using 1 ounce to each 5 gallons of warm water.
- Rinse with clean water.
- 4. Sanitize in a solution of 1 ounce Sani-Udder to 5 gallons of water. Immerse all utensils for at least two minutes, or for contact time specified by governing sanitary code.
- 5. Place sanitized utensils on rack or drain board to air dry.

Change washing and sanitizing solutions when the brown iodine color disappears. Make fresh solution each time used.

#### TABLE OF DILUTIONS:

1 ounce Sani-Udder to 5 U.S. Gallons . . . . . 25 ppm titratable iodine



# DETERGENT & SANITIZER

FOR UTENSILS - EQUIPMENT - UDDERS

**CAUTION:** KEEP OUT OF REACH OF CHILDREN

See Side Panels for Additional Cautions

## **ACTIVE INGREDIENTS:**

Nonylphenoxypoly (ethyleneoxy) ethanol -Iodine Complex (providing 1.75% titratable iodine) ...... 14.75% Phosphoric Acid ........ 6.75% 14.00% Glycolic Acid ........ INERT INGREDIENTS: ...... 64.50% 100.00% Total

Net Contents 15 U.S. Gallons

## DIRECTIONS FOR UDDER WASHING AND SANITIZING:

- 1. For each cow to be milked, place one clean towel in plastic bucket containing fresh, warm 25 ppm titratable iodine solution.
- 2. Wash udder and teats, then discard towel in second bucket.
- 3. Examine each quarter with strip cup.
- 4. After milking, dip each teat in dip cup containing 25 ppm titratable iodine solution. An efficient automatic proportioning sprayer may be used to wash the udder in place of towel or sponge.

CAUTION: Sani-Udder is a concentrated product-Harmful if Sw. llowed. Causes skin irritation. Avoid contamination of foods. Use in recommended dilutions not to exceed temperatures of 115°F. Avoid contact with skin and eyes. In case of contact, flush skin or eyes with plenty of water: for eyes, get medical attention.

U.S.D.A. Reg. No. 4715-171