

DIRECTIONS

BACKRUBBERS: For control of Horn Flies and as an aid in reduction of Face Flies on Beef Cattle dilute 1 quart with 11 quarts of fuel oil. Apply the diluted material to saturate the backrubber mechanism. Recharge the reservoir or treating surface as needed. For Horn Flies locate the backrubber to give the cattle free access to all parts of the body particularly the back.

For Face Flies, one end of the backrubber must be low enough (not more than 15 to 20 inches above ground) to allow animals to treat their heads. Dilute 1 quart with 2 gallons fuel oil. Effective only for those cattle treating face and head. Specially designed canvas face treaters may be available in your area.

Do not allow calves under 6 months of age to treat themselves. Do not allow dairy cattle access to backrubbers. Do not allow beef cattle access within 30 days of slaughter.

BEEF CATTLE AND HORSES: For control of Lone Star Tick, Ear Tick, Cattle Fever Tick and Horn Flies, dilute one quart in 25 gallons of water. For Ear Ticks treat infested ears, neck and head. Dip or apply to animal thoroughly with enough spray force to part hair and wet the skin. Do not spray within 30 days of slaughter or dip within 60 days of slaughter. Do not spray animals less than three months old or dip animals less than six months old. Do not treat dairy cattle.

For Lice on Beef Cattle, dilute 1 quart in 48 gallons of water. Thoroughly spray. If calves are infested and must be treated, do not dip if they are under 6 months of age but apply spray sparingly or sponge on infested areas. Do not add infested animals to the treated herd. Do not spray within 30 days of slaughter or dip within 60 days of slaughter.

SHEEP AND GOATS: Fleece Worms - Sheep Scab - Lice and Sheep Ticks. For Fleece Worms, mix 1 quart to 25 gallons of water. For Sheep Scab, mix 1 quart with 48 gallons of water. For Lice and Sheep Ticks (Keds) mix 8 ounces with 60 gallons of water. Dip or spray thoroughly, preferably in Spring or Summer when wool is short to give insecticide spray penetration. Replacement Ewes or Rams should be treated before adding to the flock. For Sheep Scab, dip animals making sure that material penetrates wool, especially around head. If dipping vat is intermittently used, agitate and recharge to the desired strength before next dipping. Do not spray within 30 days nor dip within 60 days of slaughter. Do not use on Dairy Goats. Do not spray or dip animals less than three months old.

HOGS: For Lice mix 1 quart in 25 gallons of water. A single application will give control. Spray Sows 30 to 45 days before farrowing.



PROTEX - TOXADANE

One Quart Makes 25 to 240 Gallons of Spray or Dip to Control Lice, Ticks, Flies, Sarcoptic Mange Mites, Fleas

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene (containing 67% to 69% Chlorine)	45%
Gamma Isomer of Benzene Hexachloride - Lindane	1%
Insecticidally Active Petroleum Hydrocarbon	44%
INERT INGREDIENTS:	10%
Total	100%

WARNING:

KEEP OUT OF REACH OF CHILDREN

See Side Panels for Additional Warnings

or three weeks thereafter. Do not spray or dip pigs that are less than weaning age or less than 3 months old. Do not spray within 30 days of slaughter or dip animals within 60 days of slaughter.

UTILITY SPRAY for Mosquitoes - Flies - Fleas and Gnats: Treat out-buildings, loading sheds, barns, fences, and waste foliage. Dilute 1 gallon to 8-1/2 gallons of water and thoroughly spray to point of runoff. Make application to walls, doors, ground areas, ceilings and places where insects rest or breed. Do not use in Dairy Barns, Milk Houses, or Poultry Houses. Do not spray foliage that may be eaten by domestic animals. Repeat applications as needed for control. For fleas, treat animal bedding and replace with new. Do not use in dwellings. Do not contaminate feed.

SARCOPTIC MANGE MITE: On Hogs, Beef Cattle and Horses, mix 1 quart in 25 gallons of water. Thoroughly spray animals with power sprayer when mange is found. Do not dip. On Horses, the preferred treatment is sponging thoroughly the infested areas. Do not spray within 30 days of slaughter. Do not spray animals less than 3 months of age. If infestation is severe, plan on a second spray treatment 15 to 21 days later.

CAUTION: This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label. Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

WARNING: May be fatal if swallowed. Do not breathe spray mist. Do not get into eyes, on skin or clothing. Wash thoroughly after using. Do not contaminate feed or food stuffs. In case of contact with eyes or skin, wash immediately and get medical attention for eyes. May be absorbed through the skin.

Avoid treating overheated or sick animals. Avoid treating animals in cold stormy weather.

Do not apply this product to Dairy Animals. Do not apply in Dairy Barns, Milk Houses or Poultry Houses nor to pets or humans. This product cannot be used on food or forage to be fed to Dairy Animals or animals being finished for slaughter. Water animals before dipping.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

4715-176

Net Contents

Mar 10, 1969

475-176