

FOR CONTROL OF ROOTWORMS AND CERTAIN OTHER
INSECTS ON CORN

SOIL INSECTS IN CORN:

FOR ROOTWORMS: Apply 1-1/3 pint per acre as a row spray treatment and 2-2,3 pints per acre for broadcast spraying. Treat just before planting and disc in immediately. Row treatment is done at time of planting with nozzles adjusted so sprayed area is covered with soil along with seed.

FOR CORN BILLBUGS, WIREWORMS, SEED CORN MAGGOT, WHITE GRUBS, GREEN JUNE BEETLE LARVAE, as well as CORN ROOTWORMS, make a broadcast application of 2/3 gallon in sufficient water to cover an acre (3-10 gallons for aerial application, 10-20 gallons for ground application). Apply before planting and disc in immediately.

SOIL INSECTS IN OTHER CROPS:

SOYBEANS: For Grape Colapsis, Seed Corn Maggot, White Grubs and Wireworms, apply 2/3 to 1 gallon per acre in sufficient water (3-10 gallons for aerial application, 10-20 gallons for ground application), treating soil prior to or at the time of planting. Cover sprayed area with soil immediately.

SMALL GRAINS: BARLEY, OATS, RYE, WHEAT - For White Grubs and Wireworms, treat same as for Soybean insects but as a pre-planting treatment only.

GARDEN INSECTS: For Wireworms on Tomatoes and Peppers, use 2/3 to 1 gallon per acre in 10-20 gallons of water (6 tbsp. in 2 gallons of water per 1,000 sq. ft.) Spray soil around new sets or seedlings before edible parts have started to form. Work into top few inches of soil.

HEPTACHLOR 300

EMULSIFIABLE CONCENTRATE

(Contains Three Pounds Actual Heptachlor per gallon)

WARNING: KEEP OUT OF REACH OF CHILDREN

See Side Panels for Additional Warnings

ACTIVE INGREDIENTS:

Heptachlor *	31.08%
Related Compounds	12.08%
Petroleum Hydrocarbons	48.84%
<u>INERT INGREDIENTS:</u>	<u>8.00%</u>
Total	100.00%

* Heptachlorotetrahydro-4,7-methanoindene

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

For Wireworms on Snap Beans, use 5-1/2 pints per acre in 10-20 gallons of water (4-3/4 tbsp. in 2 gallons of water per 1,000 sq. ft.) Apply only as a preplanting treatment. Work into top few inches of soil.

LAWN INSECTS: For Ants, Chiggers, Ear Grubs, White Grubs and Wireworms, use 4 tbsp. in 2 gallons of water (2/3 gallon per acre). For Fleas and Ticks, treat grass and foliage. Wash insects off. Keep children and pets off treated areas until material is washed into the soil and grass is cut.

WARNING: May be fatal if swallowed. Do not inhale or spray mist. Do not get on skin. In case of contact, wash immediately with soap and water. Do not eat or drink foodstuffs. Do not use on any food crops as a spray. Soil treatments only.

This product is toxic to fish and wildlife. Do not apply to streams, rivers, or other bodies of water. Do not apply to any area not sprayed. Rinse equipment and containers and dispose of in non-crop lands away from water supply. Do not dispose of by punching holes in the container.

NOTICE: Seller warrants that this product contains the ingredient statement on the label. Since conditions of use, weather, compatibility with other chemicals, and application equipment will vary, Seller makes no other warranty other than those stated on this label.

U.S.D.A. Reg. No. 4715-183 Net

**ROOTWORMS AND CERTAIN OTHER
INSECTS ON CORN**

IN:
Apply 1-1/3 pint per acre as a row spray
pints per acre for broadcast spraying.
ling and disc in immediately. Row treat-
planting with nozzles adjusted so sprayed
oil along with seed.

**S. WIREWORMS, SEED CORN MAGGOT,
EN JUNE BEETLE LARVAE**, as well as
make a broadcast application of 2/3 gallon
cover an acre (3 - 10 gallons for aerial
llons for ground application). Apply be-
in immediately.

IER CROPS:

pe Colapsis, Seed Corn Maggot, White
, apply 2/3 to 1 gallon per acre in suffi-
ons for aerial application, 10 - 20 gallons
, treating soil prior to or at the time of
yed area with soil immediately.

RLEY, OATS, RYE, WHEAT - For White
, treat same as for Soybean insects but as
ent only.

For Wireworms on Tomatoes and Peppers,
r acre in 10 - 20 gallons of water (6 tbsp.
per 1,000 sq. ft.) Spray soil around new
fore edible parts have started to form.
es of soil.

HEPTACHLOR 300

EMULSIFIABLE CONCENTRATE

(Contains Three Pounds Actual Heptachlor per gallon)

WARNING: KEEP OUT OF REACH OF CHILDREN

See Side Panels for Additional Warnings

ACTIVE INGREDIENTS:

Heptachlor *	31.08%
Related Compounds	12.08%
Petroleum Hydrocarbons	48.84%
INERT INGREDIENTS:	8.00%
Total	100.00%

* Heptachlorotetrahydro-4,7-methanoindene

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

For Wireworms on Snap Beans, use 5-1/4 to 6-1/2 pts. per
acre in 10 - 20 gallons of water (4-3/4 tbsp. in 2 gallons of water
per 1,000 sq. ft.) Apply only as a preplanting treatment and
work into top few inches of soil.

LAWN INSECTS: For Ants, Chiggers, Earwigs, Fleas, Ticks,
White Grubs and Wireworms, use 4 tbsp. in 2 gallons of water
for 1,000 sq. ft. (2/3 gallon per acre). For Chiggers, Fleas
and Ticks, treat grass and foliage. Wash into soil for soil in-
sects. Keep children and pets off treated areas until this ma-
terial is washed into the soil and grass is dry.

WARNING: May be fatal if swallowed. Do not breathe vapor
or spray mist. Do not get on skin. In case of contact, wash
immediately with soap and water. Do not contaminate feed and
foodstuffs. Do not use on any food crops as a direct application
spray. Soil treatments only.

This product is toxic to fish and wildlife. Birds feeding on
treated areas may be killed. Keep out of lakes, ponds and
streams. Do not apply to any area not specified on the label.
Rinse equipment and containers and dispose of wastes by bury-
ing in non-crop lands away from water supplies. Containers
should be disposed by punching holes in them and burying with
wastes.

NOTICE: Seller warrants that this product conforms to the
ingredient statement on the label. Since conditions of use, such
as weather, compatibility with other chemicals, and condition
of application equipment will vary, Seller makes no claims
other than those stated on this label.

U.S.D.A. Reg. No. 4715-183

Net Contents Gal.