

Broad Spectrum FLOWER & GARDEN SPRAY

Contains 2 lbs. Methoxychlor and 2 lbs. Malathion per gallon

CAUTION:
KEEP OUT OF REACH OF CHILDREN

See side panel for other cautions.

ACTIVE INGREDIENTS:

- * Methoxychlor, Technical 24.0%
- * Malathion 24.0%
- Aromatic Petroleum Derivative Solvent 47.0%

INERT INGREDIENTS: 5.0%

Total 100.0%

- * Equivalent to 21% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 3% other isomer and reaction products.
- ** O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

Net Contents _____

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

STRAWBERRY - For control of leafhoppers, Colorado potato beetle, leafhoppers, leaf beetles, aphids, thrips, spider mites, white fly, and other insects. Use 2-3 qts. per acre. For control of strawberry leaf beetles, strawberry root weevil & whiteflies. Use 2-3 qts. per acre. For control of aphids. **CAUTION** - For use above 3-4-2 qts. per acre should not be applied within 14 days of harvest.

FLOWERS & ORNAMENTAL SHRUBS
Cystic Shell Scale & Eucalyptus Scale - Use 2-4-1 Pt.
Aphids, Spider Mites, White Fly, Thrips, Mealy Bugs, Japanese Beetle, Adult, Four-lined Plant Bug, Fernished Plant Bug, Rose Leafhopper, Potato Leafhopper - Use 3-3-1 Pt.
Bugworms, Juniper Scale - Use 2-1-2 qt. Soft-Bodied Scale - Use 1-1-1 Pt.
Pine Needle Scale - Use 5 qts.

Apply in summer when infestation is first. Repeat in fall. For severe infestations, apply when crawlers have settled on foliage.

CAUTION - Do not use on certain ferns, including Boston Maiden Hair Ferns, as well as some species of Crassula, because of possible injury to these plants. Do not spray plants of viburnum and avoid repeated applications to evergreens. Do not spray when temperatures are abnormally high.

SHRUB & TREE INSECTS - For control of May Beets, Leafhoppers, Japanese Beetles, Sawflies, Nymphs, Leafhoppers, Japanese Beetle, Forest Tent Caterpillar, Gypsy Moth, Gypsy Moth, Elm Leaf Beetle, and Cankerworms. For use on trees, use 1 to 2 parts per gallon of water in conventional sprayers. When using a mist blower, dilute one part of Flower & Garden Spray to 3 parts water using approximately 1-2 parts per gallon of tree.

CAUTION - Chinese plants may be injured by foliage sprays of high degree Flower & Garden spray when applied with conventional spraying equipment, especially under conditions of drought and high temperature. Do not spray when temperatures are abnormally high. Avoid repeated applications to evergreens.

SMALL QUANTITY DILUTION TABLE

Use spray nozzle and use following dilution table:

Quantity of Label Shows	Use This Amount for Each Gallon of Water	Quantity of Label Shows	Use This Amount for Each Gallon of Water
1 qt.	2 tsp.	1-1/2 qt.	3 tsp.
2-1/2 Pt.	2-1/2 tsp.	2 qt.	4 tsp.
1-1/2 qt.	3 tsp.	3-1/2 Pt.	5-1/4 tsp.
3-3/4 Pt.	4-3/4 tsp.	4-1/2 qt.	7 tsp.
5 qt.	5 tsp.	5 qt.	8 tsp.

CAUTION - Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing fumes of spray mist. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water after using. Keep away from children and domestic animals. Destroy container when empty by perforating or crushing and burying in a safe place. Do not reuse. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE - Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label. U.S.P.A. Reg. No. 4715-201