

Application

Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Scale Insects. Apply Broad Spectrum Orchard Spray at the specified amount with an expert or good operator and direct the spray stream into the nearly filled spray tank, up to the leaves of the tree. Spray early in the morning, spray when the temperature is in the 60's or 70's. Coverage of all exposed plant surfaces is essential. The spray should be mixed with 100 gallons of water. The spray should be applied to the leaves of the tree mixed with sufficient water for 2 hours before harvest. Do not apply more than once every 7-day interval between last application and harvest. Do not mix with the crop. Observe use limitations.

Broad Spectrum ORCHARD SPRAY

Contains: 24.0% Methoxychlor, 24.0% Malathion, 5.0% Inert

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for other cautions.

ACTIVE INGREDIENTS:

• Methoxychlor, Technical	24.0%
• Malathion	24.0%
• Aromatic Petroleum Derivative Solvent	47.0%

INERT INGREDIENTS:	5.0%
Total	100.0%

- Equivalent to 21% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 3% other isomer and reaction products.
- 0.0% dimethyl dithiophosphate of diethyl mercaptosuccinate

Net Contents _____

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

Broad Spectrum ORCHARD SPRAY

Contains 24.0% Methoxychlor and 24.0% Malathion per gallon

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for other cautions.

ACTIVE INGREDIENTS:

• Methoxychlor, Technical	24.0%
• Malathion	24.0%
Aromatic Petroleum Derivative Solvent	47.0%

INERT INGREDIENTS:	5.0%
Total	100.0%

- Equivalent to 21 1/2% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 3% other isomer and reaction products.
- O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

Net Contents _____

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

days if needed. Japanese Beetles - Use 1 qt. See Application directions. Rose Chater - Use 2 Qt. When adults first appear, usually near bloom of Concord grapes. Two-spotted Spider mite - Use 1 qt. When mites first appear, and 7 days later if infestation is heavy. CAUTION: May injure the Rubier, Italia, Cardinal & Almeria varieties if applied off clusters appear.

BLACKBERRY, BOYSENBERRY, DEWBERRY, LOGANBERRY, RASPBERRY (14): Use 3-1/2 Qt. on foliage as needed to control Aphids, Japanese Beetle, Rose Chater, Rose leafhopper, Rose Scale and Strawberry Weevil.

BLUEBERRY (14): Cherry Fruitworm - Use 1 qt. Three applications at 10 day intervals beginning just after petal fall. Japanese Beetle - Use 2 Qt. On foliage when adults appear, repeat as needed.

CURRENT (14) & GOOSEBERRY (14): Currant Aphid - Use 4 qt. As leaf buds are opening or later if infestation occurs. Imported Currantworm - Use 4 Qt. When worms appear.

STRAWBERRY (3): Use 2-4 Qts. on foliage as needed to control Field Cricket, Lygus Bugs, Spittlebugs & Thrips. Use 2-3-1/2 Qts. on foliage as needed to control Flea Beetles, Strawberry Leaf Beetle, Strawberry Weevil. Use 2-3 Qts. on foliage as needed to control Potato Leafhoppers, Strawberry Leaf Roller, Strawberry Root Weevil and Whiteflies. Use 3-1/2 Qts. on foliage as needed to control Aphids. Caution: Dosages above 3-1/2 Qts. per acre should not be applied within 14 days of harvest.

SHADE TREE INSECTS: To control May Beetle, Tree Bugs on oak and sycamore, Tussock Moth, Fall Webworm, Japanese Beetle, Forest Tent Caterpillar, Eastern Tent Caterpillar, Gypsy Moth, Elm Leaf Beetle and Cankerworms on shade trees, use 1 to 2 quarts per 100 gallons of water in conventional sprayers. When using a mist blower, dilute one part of Orchard Spray to 3 parts water using approximately 1-2 pints per 35-50 foot tree.

Caution: Chinese Elms may be injured by foliar spray. Chinese elms are injured by spray when applied with conventional spraying equipment, especially under conditions of drought and high temperature. Do not spray when temperatures are abnormally high. Avoid repeated applications to evergreens.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing fumes or spray mist. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and water after using. Keep away from children and domestic animals. Destroy container when empty by perforating or crushing and burying in a safe place. Do not re-use. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 111-020