

DIRECTIONS FOR USE

This is a concentrate and must be diluted with base oil before fogging. To each one-half gallon of concentrate add 90 gallons of No. 2 diesel oil and mix thoroughly. Follow the directions in the manual of your fogging equipment for best results. Do not mix with water as this is not an emulsifiable concentrate.

OUTDOOR FOGGING: Apply diluted solution of fogging concentrate at the rate of one gallon per acre. The fog should be directed over the entire area. Particular attention should be given to areas where flies congregate such as fences, walls of buildings. Animals may be present during treatment of feedlots. Avoid direct application to exposed feed and water. Do not apply the fog where food or feed crops are growing. Not for use on home lawns.

FOR CONTROL OF FLY MAGGOTS: In maggot breeding areas such as garbage dumps and manure piles, apply at the rate of 1/2 gallon of diluted solution per 100 sq. ft. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Repeat as necessary. Birds feeding on treated areas may be killed.

FOR ANIMAL BUILDINGS: Apply as a fog inside buildings at rate of 1 gallon of diluted solution per 32,000 cubic feet. Dairy and beef cattle and horses may be present during application. Do not use this solution for fogging purposes in areas where animals have received direct application within eight hours. Do not contaminate water, feed, milk or milking utensils.

POISONOUS TO FISH AND WILDLIFE: This product is toxic to fish and wildlife. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

COMBUSTIBLE MIXTURE: DO NOT POUR, SPILL OR STORE NEAR FIRE OR HEAT.

DANGER: May be fatal if swallowed, inhaled or absorbed through the skin or eyes. Do not get on skin, in eyes, or on clothing. Do not breathe fumes. Wear rubber gloves, protective clothing and goggles when handling concentrate to prepare solution. In case of contact, remove contaminated clothing and wash skin thoroughly with soap and water; for eyes, flush with plenty of water; get medical attention. If poisoning should occur, CALL A PHYSICIAN IMMEDIATELY. ATROpine AND 2-PAM ARE ANTIDOTES. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed, foodstuffs or drinking water. Do not use inside homes where milk is stored or processed. Avoid breathing of spray mist. During fogging or mist spraying applications wear a mask or respirator approved by the U. S. Bureau of Mines for use with DDVP. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operational area.



Burn or destroy containers, never reuse. Stay out of smoke or fumes. Bury spillage in non-crop area away from water supplies. Clean up area with strong lye solution.

ANTIDOTE AND FIRST AID: Emergency antidote for DDVP is about obtaining an adequate dose of emergency use. 2-PAM should be administered in conjunction with IMMMEDIATELY IN ALL POISONING.

INTERNAL: If the material has been吞咽 (swallowed), immediately. This may be done by throat or by giving warm salt water (one glassful of water). Repeat until vomit ANYTHING BY MOUTH TO THE VOMIT.

EXTERNAL: If the material has been absorbed through the skin, immediately remove patient from contact with the material. Remove all contaminated clothing and wash with running water for at least 15 minutes.

IF WARNING SYMPTOMS OCCUR: Atropine tablets, minimum dose of 1/2 grain. ATROpine OR 2-PAM (UNLIQUIDATED) See Warning Symptom section. Start artificial respiration if necessary. Transport the patient to a hospital.

NOTE TO PHYSICIAN: Watch for signs of respiratory distress. If patient shows signs of respiratory distress, give oxygen. If patient shows signs of convulsions, give morphine. If patient shows signs of shock, give epinephrine. If patient shows signs of heart trouble, give digitalis. If patient shows signs of kidney trouble, give diuretics. If patient shows signs of liver trouble, give cholestyramine.

TREATMENT: Attention is the following first aid complications of para-oxonathene stimulate. Injected at 0 minute intervals, complete respiration and oxygenation and may be administered. NOT GIVE MORPHINE OR other sedatives. Oxygen administration. Oxygen administration. Sedatives may be susceptible to very small doses. Allow a sufficient exposure time to be determined.

NOTICE: Seller warrants the ingredient statement on the label, weather, compatibility with other application equipment will vary, those stated on this label.

DDVP 10 POUND OIL SOLUTION

**To Be Used Only By Public Health Officials,
Mosquito Abatement Districts
And Pest Control Operators.**

DANGER ! **POISON !**

**KEEP OUT OF REACH OF CHILDREN. See Side Panels for
Antidote and Other Required Warning Statements.**

ACTIVE INGREDIENTS:

| | |
|---|---------|
| 2,2-dichlorovinyl dimethyl phosphate (DDVP) | 83.78%* |
| Related Compounds | 6.31%* |
| Aromatic Petroleum Derivatives..... | 9.91% |
| Total | 100.00% |

*Equivalent to 90.09% w Technical DDVP Insecticide.
Contains 10 lbs. active ingredient per gallon.

EPA REG. NO. 4715-232

Net Contents 10 Gal

DDVP 10 POUND OIL SOLUTION

To Be Used Only By Public Health Officials,
Mosquito Abatement Districts
And Pest Control Operators.

DANGER !  POISON !

KEEP OUT OF REACH OF CHILDREN. See Side Panels for
Antidote and Other Required Warning Statements.

ACTIVE INGREDIENTS:

| | |
|---|---------|
| 2,2-dichlorovinyl dimethyl phosphate (DDVP) | 83.78%* |
| Related Compounds | 6.31%* |
| Aromatic Petroleum Derivatives..... | 9.91% |
| Total | 100.00% |

*Equivalent to 90.09% w Technical DDVP Insecticide.
Contains 10 lbs. active ingredient per gallon.

ANTIDOTE AND FIRST AID TREATMENT. Atropine is the emergency antidote for DDVP poisoning. Consult your physician about obtaining an adequate supply of 1/100 gram atropine tablets for emergency use. 2-PAM is also antidotal and may be administered in conjunction with atropine. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). Repeat until vomit fluid is clear. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash them immediately with running water for at least ten minutes; get medical attention.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100 gram atropine tablets immediately. NEVER ADMINISTER ATROPINE OR 2-PAM UNLESS WARNING SYMPTOMS APPEAR. (See Warning Symptoms below. Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.)

NOTE TO PHYSICIAN: Warning Symptoms - Symptoms include weakness, cardiac tightness in chest, diarrhea, confusion, twitching, convulsions, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of nerve agent-type sympathetic nervous stimulation. If there are signs of post-sympathetic stimulation, atropine sulfate should be injected at 10 mg. intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. 2-PAM chloride is also an antidote and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 4715-232

Net Contents 1 Gal