

Cameraman's Note

Poor Copy

DIRECTIONS FOR USE

This is a concentrated and must be diluted prior to use before fogging. To dilute one gallon of concentrate add 90 gallons of water. Dilute one gallon of concentrate to 90 gallons of water. Follow the directions on the label of your fogging equipment for best results. Do not mix with water as this will reduce the effectiveness.

OUTDOOR FOGGING To control flies, mosquitoes and gnats apply diluted solution of fogger to generate a fine mist at the rate of 3-5 gallons per acre. Fogging should be done over the entire area. Particular attention should be given to areas where flies or mosquitoes are present such as around waste disposal areas. Animals may be present during treatment of feedlot. Avoid direct application to exposed feed and water. Do not apply the fog when there is a strong wind or rain. Do not use in confined areas.

FOR CONTROL OF FLY MAGGOTS Dilute concentrate at the rate of 1-2 gallons of diluted solution per 100 sq. ft. of area. Apply to water, feedlot, and waste disposal areas. Repeat applications as needed.

FOR ANIMAL BUILDINGS To control flies and mosquitoes in animal buildings, dilute concentrate at the rate of 1-2 gallons per 100 sq. ft. of area. Apply to water, feedlot, and waste disposal areas. Repeat applications as needed.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

DANGER - This product is highly toxic. It is fatal if swallowed, inhaled, or absorbed through the skin. It causes severe irritation to the eyes, nose, and throat. It is also a strong irritant to the respiratory system. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling. Do not breathe vapors. Do not get in eyes, on clothing, or on food.

ANTIDOTE AND FIRST AID TREATMENT - ATROPINE
In case of poisoning, give 1-2 mg. of atropine immediately. If necessary, repeat every 30 minutes until symptoms subside. Do not induce vomiting. Do not give anything by mouth to an unconscious person.


INTERNAL - In case of poisoning, give 1-2 mg. of atropine immediately. If necessary, repeat every 30 minutes until symptoms subside. Do not induce vomiting. Do not give anything by mouth to an unconscious person.

ACCEPTED
JUL 24 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4-715-233



80% DDVP OIL SOLUTION

Contains 8.4 lbs. Technical DDVP Insecticide Per Gallon
To Be Used Only By Public Health Officials
Mosquito Abatement Districts and Pest Control Operators



DANGER: KEEP OUT OF REACH OF CHILDREN

See Side Panels for Antidote and Other Required Warning Statements

ACTIVE INGREDIENTS	PERCENTAGE
2,2-dichlorovinyl dimethyl phosphite (DDVP)	74.4%
Related Compounds	5.6%
Aromatic Petroleum Derivative	20.0%
TOTAL	100.0%

*Equivalent to 80%w Technical DDVP insecticide

NET CONTENTS: 1 5 30 55 GALLONS

EXTEN
immedi
remove
nap and
minutes
IF WA
1-100-gr
MINIST
APPEA
patient
separate
NOTE
and the
vision
TREAT
gonist
lation
study
Treatm
at 12
MORPH
Ob
ENVIR

4

ACCEPTED
JUL 24 1975
Federal Insecticide, Fungicide, and Rodenticide Act
Registered under Reg. No. 4715-233

80% DDVP OIL SOLUTION

Contains 8.4 Lbs. Technical DDVP Insecticide Per Gallon
To Be Used Only By Public Health Officials,
Mosquito Abatement Districts and Pest Control Operators.



DANGER: KEEP OUT OF REACH OF CHILDREN

See Side Panels for Antidotes and Other Required Warning Statements.

ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate (DDVP)	74.4%
Related Compounds	5.6%
Aromatic Petroleum Derivatives	20.0%
Total	100.0%

*Equivalent to 80%w Technical DDVP insecticide.

NET CONTENTS GALLONS
1 30 35

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the DDVP, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, flush immediately with running water for at least 15 minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1-100-grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** See **WARNING SYMPTOMS** below. Keep patient prone on floor. Start artificial respiration immediately if patient is not breathing. Transport to the nearest hospital immediately.

NOTE TO PHYSICIAN: Warning Symptoms: Symptoms include weakness, headache, nausea, vomiting, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific antidote for the parasympathetic stimulation caused by DDVP. Administer 1-100-grain atropine tablets in emergency use. If there are signs of parasympathetic stimulation, atropine tablets should be injected at 10-minute intervals in doses of 1-2 milligrams until complete atropinization is achieved. 2-3 milligrams may be given intravenously. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Oxygen should be administered. Give artificial respiration if necessary. Administer 1-100-grain atropine tablets every 15-30 minutes until the patient is breathing spontaneously. Administer 1-100-grain atropine tablets every 15-30 minutes until the patient is breathing spontaneously.

ENVIRONMENTAL CAUTION: Do not use in areas where water is used for drinking. Do not apply where it may be carried off by wind or water. Do not use where it may be carried off by wind or water. Do not use where it may be carried off by wind or water.

This product is flammable. Keep away from fire. Treatment should be given immediately. Antidotes may be obtained from the manufacturer, Entonox, Inc., Extension Station.

Decorate and destroy used containers. Do not reuse. Do not reuse. Do not reuse. Do not reuse. Do not reuse.

Net weight 8.4 lbs. per gallon. Contains 8.4 lbs. per gallon. Contains 8.4 lbs. per gallon.

EP-1715

Camerman's Note
Poor Copy

DIRECTIONS FOR USE

This is a concentrate and must be diluted with base oil before fogging. To use, one-half gallon of concentrate add 80 gallons of No. 2 diesel oil and mix thoroughly. Follow the directions in the manual of your fogging equipment for best results. Do not mix with water as this is not an emulsifiable concentrate.

OUTDOOR FOGGING: To control flies, mosquitoes and gnats, apply diluted solution of fogging concentrate at the rate of 3 1/4 gallons per acre. The fog should be directed over the entire area. Particular attention should be given to areas where flies congregate such as fences, walls of buildings. Animals may be present during treatment of feedlots. Avoid direct application to exposed feed and water. Do not apply the fog where food or feed crops are growing. Not for use on home lawns.

FOR CONTROL OF FLY MAGGOTS: In maggot breeding areas such as garbage dumps and manure piles, apply at the rate of 1/2 gallon of diluted solution per 100 sq. ft. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Repeat as necessary. Birds feeding on treated areas may be killed.

FOR ANIMAL BUILDINGS: To control flies, mosquitoes, roaches, ants and sowbugs, apply as a fog inside buildings at rate of 1 gallon of diluted solution per 32,000 cubic feet. Dairy and beef cattle and horses may be present during application. Do not use this solution for fogging purposes in areas where animals have received direct application within eight hours. Do not contaminate water, feed, milk or milking utensils.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DANGER Poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or in eyes. Do not breathe fumes. Avoid exposure to fumes. Wear clean rubber gloves, waterproof protective clothing and goggles when handling this material.

Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for DD-2 protection. In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

ANTIDOTE AND FIRST AID TREATMENT ATROPINE is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine. Call a physician in all cases of suspected poisoning.

INTERNAL: Induce vomiting immediately. This may be done by introducing a finger to the throat or by giving 1 Tbsp. of salt in a glass of warm water. Repeat until vomit fluid is clear. Never given anything by mouth to an unconscious person!

ACCEPTED

JUL 24 1970

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 4715

Con

Mos

DA

See Side

ACTIVE

2,2 dic

Related

Aroma

*Equivalent

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the DDVP; remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, flush immediately, with running water for at least 10 minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See **WARNING SYMPTOMS** below.) Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport to the nearest hospital immediately.

NOTE TO PHYSICIAN: Warning Symptoms. Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against para-sympathetic nervous stimulation. Consult your physician about obtaining an adequate supply of 1/100-grain atropine tablets for emergency use. If there are signs of para-sympathetic stimulation atropine sulfate should be injected at 10-minute intervals, in doses of 1-2 milligrams, until complete atropinization has occurred. 2-PAM chloride is also an antidote and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Clear airway by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to even small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

ENVIRONMENTAL CAUTION: This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. For more information, may be obtained from your Cooperative Agricultural Extension Service.

Decontaminate and destroy used containers by washing thoroughly with a dilute solution of caustic (less than 5%) and detergent. **CAUTION:** Avoid skin and eye contact with caustic solutions. Rinse containers with water and dispose of the washings in such a way that they do not constitute a hazard to, or contaminate, water supplies. Destroy containers by crushing and burning, or bury in non-cropland away from water supplies.

NOTICE: Bayer assumes no risk of the storage or handling of this material not in the original unopened containers given herewith.

EPA Reg. No. 4715-207

EPA Est. 4715-CO

ments.

4%*
6%*
0%
0%