



**CAUTION:** Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

**IMPORTANT:** Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides and fungicides. Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

**DIRECTIONS**

**WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS:** To kill Poison Ivy, Rag Weed, Morning-glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if re-growth develops. Use 1.5 fl.oz. in 1 gallon of water or 1 cupful in 5 gallons of water.

**LAWN WEEDS:** For the most effective control at the lowest dosage, apply when the weeds are growing vigorously, after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank

**DANDELION DESTROYER WITH BANVEL D<sup>®</sup>**

For Control of Chickweed and Certain Broadleaf Turf Weeds On Lawns and Golf Courses

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**  
See Side Panel For Additional Cautions.

**ACTIVE INGREDIENTS:**

|  |                |
|--|----------------|
| Diethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)* | 3.43%          |
| Diethylamine Salts of Related Acids                        | 0.55%          |
| Diethylamine Salt of 2,4-Dichlorophenoxyacetic acid**      | 17.22%         |
| <b>INERT INGREDIENTS:</b>                                  | <b>78.80%</b>  |
| <b>Total</b>   | <b>100.00%</b> |

\* Equivalent to 2.84% 3,6-dichloro-o-anisic acid (dicamba) or 0.26 pounds per gallon.  
\*\* Equivalent to 14.37% 2,4-Dichlorophenoxyacetic acid or 1.25 pounds per gallon.

**DIRECTIONS**

type or hose-end sprayer adjusted to deliver For Dandelions, Chickweed, Clover, Plantain, Dandelion, Wild Onion and many other broadleaf weeds. . . . FOLLOW DILUTION IN SCHEDULE A. For late applications and for hard-to-kill weeds such as Oxalis, Yarrow and nutcracker knotweed, INCREASE SCHEDULE B. If weeds persist, repeat application three weeks. Wild Onions will require repeat in late fall and early spring.

**DILUTION CHART**

| SCHEDULE A       | SCHEDULE B       | U.S. Gal. |
|------------------|------------------|-----------|
| 1 Tablespoonful  | 3.5 Teaspoonfuls |           |
| 1/2 fl. oz.      | 0.6 fl. oz.      |           |
| 5 Tablespoonfuls | 6 Teaspoonfuls   |           |
| 2 1/2 fl. oz.    | 1 fl. oz.        |           |
| 1 Pint           | 19 fl. oz.       | 10        |

**NOTICE:** Seller warrants that this product contains the ingredient statement on the label. Since conditions such as weather, compatibility with other chemicals and condition of application equipment will vary, claims other than those stated on this label.

EPA REG. NO. 4715-234 NET CONTENTS

7-26-72  
4715-234  

---

40X31-70



## DANDELION DESTROYER WITH BANVEL D\*

For Control of Chickweed and Certain  
Broadleaf Turf Weeds On Lawns and  
Golf Courses

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**  
See Side Panel For Additional Cautions.

**ACTIVE INGREDIENTS:**

|   |                |
|---|----------------|
| Diethylamine Salt of Dicamba (3,4-dichloro-o-anisic acid)*..... | 1.43%          |
| Diethylamine Salts of Related Acids.....                        | 0.55%          |
| Diethylamine Salt of 2,4-Dichlorophenoxyacetic acid**.....      | 17.22%         |
| <b>INERT INGREDIENTS.....</b>                                   | <b>78.80%</b>  |
| <b>Total</b>  | <b>100.00%</b> |

\* Equivalent to 2.84% 3,4-dichloro-o-anisic acid (dicamba) or 0.26 pounds per gallon.  
\*\* Equivalent to 16.27% 2,4-dichlorophenoxyacetic acid or 1.25 pounds per gallon.

### DIRECTIONS

Type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Herbit, Wild Onion and many other broadleaf weeds in the Spring ... FOLLOW DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Oxalis, Yarrow and nutcracker, INCREASE THE DOSAGES TO SCHEDULE B. If weeds persist, repeat application in two or three weeks. Wild Onions will require repeated applications in late fall and early spring.

### DILUTION CHART

| SCHEDULE A                     | SCHEDULE B                      | Water Gallons | Area Sq. Ft. |
|--------------------------------|---------------------------------|---------------|--------------|
| 1 Tablespoonful<br>4 fl. oz.   | 1.5 Teaspoonfuls<br>0.6 fl. oz. | 1             | 200          |
| 5 Tablespoonfuls<br>24 fl. oz. | 6 Teaspoonfuls<br>3 fl. oz.     | 5             | 1000         |
| 1 Pint                         | 19 fl. oz.                      | 30            | 6500         |

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 4715-234      NET CONTENTS

and clothing. In case  
er; for eyes, get med-  
Keep out of the reach

sprays are less likely  
ornamental shrubs, trees  
treat under drip line  
ornamental shrubs and  
irrigation waters. Do  
near desirable trees  
water. Do not store in  
insecticides and fungi-  
t. Augustine grass or  
airable bent-grass or  
seeded lawns or turf.  
thoroughly and do not  
for applying 2,4-D

**USUAL AREAS:** To kill  
weeds and many other  
are actively growing.  
if re-growth develops,  
1 cupful in 5 gallons

control at the lowest dos-  
age possible after a Spring  
treated and apply the  
area. Use either a tank

1-70