



# LAWN WEED KILLER

## Contains 2,4-D & Silvex

Kills **CHICKWEED, SPOTTED SPURGE** and other Broadleaf Lawn Weeds

**ACTIVE INGREDIENTS:** BY Weight  
Mixed Octyl Ester of 2,4-Dichlorophenoxyacetic Acid (2,4-D) 2.8%  
Isooctyl Ester of Silvex (2,2,4,5-Trichlorophenoxy Propionic Acid) 17.0%

**INERT INGREDIENTS:** 77.2%

\*Equivalent to 10% 2,4-D Acid  
\*\*Equivalent to 5% Silvex

Contains the equivalent of 9.8 lbs. of 2,4-D Acid and 0.4 lbs. of Silvex per gallon.

### CAUTION: Keep out of Reach of Children.

Harmful if swallowed. Avoid contact with skin, eyes, and hair. Wash hands after using. Keep out of reach of children and domestic animals.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

**ENVIRONMENTAL CAUTION:** This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water used for domestic irrigation purposes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift of spray from treated areas. Apply this product only as specified on this label.

Do not reuse empty container. Rinse smaller container from sprayer and place in trash can for disposal. Empty container by pouring perforation and burn in non-polluted area away from water supply.

NET CONTENTS: ONE GALLON  
COLORADO INTERNATIONAL CORPORATION  
Commerce City, Colorado 80022

# ACCEPTED

## JUL 3 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **4715-225**

**DO NOT** store near seeds, fertilizers, insecticides or fungicides

### GENERAL INFORMATION

For most effective control, apply when weeds are growing vigorously after a rain in the Spring or Fall. Poor results will be obtained during periods of drought and turf injury may occur. Do not apply during the heat of the day. If rain is expected within 24 hours, delay application.

### DIRECTIONS FOR USE

Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Apply as a coarse spray to prevent spray drift to desirable plants. Vapors from the product may injure susceptible plants in immediate vicinity. Do not use same spray equipment for purposes other than application of insecticides and herbicides. **NOTE:** Do not permit children or domestic animals to go onto the sprayed grass until the spray has completely dried.

### HARD-TO-KILL WEEDS

**Ragweed, Dock, Thistle, Morning Glory in Lawns, Fence Rows and Rights-of-Way.** Use 3 lbs. per gallon of water (1-100 dilution). Spray to thoroughly wet all plant surfaces. Repeat as necessary.

### LAWN WEEDS

**Clover, Chickweed, Chicory, Dandelion, Henbit, Plantain, Spotted Spurge and Wild Onions:** Mix 4 Tbsp. (1/2 oz.) in 2 gallons of water. Apply to 500 sq. ft. of lawn area. If lawn begins to grow, apply application, and repeat. **Wild Onions** may require repeated applications in late fall and early spring. **NOTE:** Do not use on Dandelion, Poppy, or the Yellow and White Perennial Ryegrass. Do not apply to new lawn until after at least second cutting. Do not reseed lawn during week period after use of this material.

### WOODY PLANTS and WEEDS

**Poison Ivy, Poison Oak, Honeysuckle, Sumac, Brambles** - Use general directions with control in industrial fields, Fence Rows and Rights-of-Way. Use (1-100) dilution of water. Spray to thoroughly wet all plant surfaces. Application may be repeated in 2 weeks.

**NOTICE:** Seller warrants that the product conforms to the implied state of the art for the use intended. Seller disclaims any liability for injury or damage with the exception of those specifically stated on the label. Seller disclaims any liability for injury or damage with the exception of those specifically stated on the label.