



25% LINDANE

WETTABLE POWDER

FOR CERTAIN GARDEN INSECTS
AND CONTROL OF FLIES, LICE, TICKS AND MANGE ON BEEF CATTLE

ACTIVE INGREDIENT:	
Gamma Isomer of Benzene Hexachloride from Lindane	25%
INERT INGREDIENTS:	
	75%
TOTAL	100%

WARNING: KEEP OUT OF REACH OF CHILDREN
See Side Panel for required warning statements

DIRECTIONS

LIVESTOCK PESTS—For Lice on Beef Cattle, Horses and Hogs, use 1 pound per 100 gallons of water (2 level teaspoons per gallon). Treat entire animal or infested areas. May be necessary to repeat treatment within 15 to 20 days. For Sarcoptic Mange on Beef Cattle, Horses, Hogs and Sheep, use 1½ pounds per 100 gallons of water (3 level teaspoons per gallon). Use about 2 gallons of spray per animal for cattle and horses and proportionately less for smaller livestock. Use a power sprayer capable of developing sufficient pressure to part the hair and wet the skin. Apply two sprays one week apart. For ticks on beef cattle and horses, use 1 pound per 100 gallons of water. Thoroughly treat all areas, wetting the skin. Make two applications about 20 days apart or as needed for continued control. For lice and ticks on Sheep, use 1 pound per 100 gallons of water. Apply in the fall. It may be necessary to repeat treatment after shearing. Spray against grain of wool along back and around ears. The spray machine should develop enough pressure to penetrate the wool and wet the skin. Do not dip animals. Do not spray animals within 30 days of slaughter. Do not use on Dairy Animals.

CAUTION: Do not treat animals under 3 months of age. Sprays for young animals (over 3 months of age) should not exceed a concentration of 0.03% lindane (1 pound of this product in 100 gallons of water). Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather. Avoid treating overheated or sick animals.

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

Net Contents _____ Pounds

DIRECTIONS

RESIDUAL SPRAYS—To control flies, mosquitoes, fleas, roaches and ants in farm buildings, use 1 pound per 10 gallons of water. Apply as a coarse wet spray at the rate of 1 gallon diluted spray to 500 square feet or until surface is thoroughly wet. For flies and mosquitoes, spray ceilings, walls, posts, window and door frames where flies prefer to rest. Treat adjacent buildings to reduce the number of flies entering the barn. For fleas, make applications to walls, baseboards, floors and to bedding in animal sleeping quarters. For roaches and ants, treat floors, baseboards, cracks, and crevices and other hiding places. Direct spray on to as many insects as possible at time of application. Repeat as needed. Do not use inside homes, poultry houses, dairy barns or milk rooms. Do not contaminate milk, milking equipment, feed or drinking water.

V. GETABLES: For Aphids or Plant Lice mix 3 lbs. 25% Lindane Wettable Powder in sufficient water to cover one acre and spray both top and under sides of all leaves on the following: **CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, BROCCOLI**—Do not apply after heads or edible parts begin to form; **CUCUMBERS**—Do not apply after blossoming begins; **SPINACH**—Do not apply after seedling stage; **TOMATOES**—Do not apply after blossoms fall. For **CELERY**, spray seedling stage only using 1½ lbs. 25% Lindane per acre. For **PEPPERS**, do not spray after blossoms fall, using 1½ lbs. 25% Lindane per acre.

HOME GARDENS: For Aphids or Plant Lice on Cabbage, Cauliflower, Brussels Sprouts, Broccoli, Cucumbers, Tomatoes, Celery and Peppers, mix 1 teaspoon 25% Lindane with one gallon of water and spray as above, observing all cautions given.

SQUASH BUGS: Mix 2 lbs. 25% Lindane Wettable Powder and 4 lbs. 50% Methoxychlor Wettable Powder in sufficient water to cover one acre (1 teaspoon 25% Lindane & 2 teaspoons 50% Methoxychlor with one gallon of water for Home Gardens) and apply so as to cover affected pumpkin and squash plants thoroughly. Repeat every two weeks. Do not apply within 7 days of harvest. Application in late season may cause some reduction in quality of squash.

CAUTION

CAUTION: Lindane should not be used in soils where root crops are to be grown. Do not use on household pets or humans.

Burn bags when empty; never re-use. Keep out of smoke. Bury any spillage or spray waste in non-crop areas away from water supplies.

This product is toxic to fish and wildlife. Keep out of any body of water. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

WARNING: May be fatal if swallowed! May be absorbed through skin! Do not breathe dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Avoid contamination of feed and foodstuffs.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 4715-252

