

DIRECTIONS FOR USE
SOIL TREATMENT

For control of soil insects such as Earth larvae, White Grubs, and Wireworms, apply 5 to 10 lbs. per acre on top dressings, corn, soybeans, alfalfa, timothy, rye, and other feed crops. For control of Green June beetle larvae, alfalfa, timothy, rye, and other feed crops, add 5 to 12 lbs. in sufficient water to cover one acre (4 1/2 oz. per 1000 square feet). For Green June beetle larvae use 12 to 20 lbs. per acre (7 1/2 oz. per 1000 square feet). Apply as a soil treatment before or at the time of planting. On beans do not exceed 10 lbs. per acre.


ANTIDOTE: Emetic of 1 tablespoon mustard in tumbler of water. Call a physician. In case of spilling on skin, wash promptly with soap and water.

WARNING: May be fatal if swallowed. Do not breathe vapor, dust or spray mist. Do not get on skin. In case of contact wash immediately with soap and water. Do not contaminate feed and foodstuffs.

CAUTION: Highly toxic to fish and wildlife. Do not contaminate water by cleaning of equipment or disposal of wastes. Treated seed exposed on the soil surface will be hazardous to birds.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 4715-257 Net Weight _____ lbs.



NO-STIK
HEPTACHLOR
25% SEED TREATER

With Graphite Added

For Use in the Control of False Wireworms, Wireworms, Corn Rootworms, Seed Corn Maggots, and Certain Other Soil Insects as Named Below

ACTIVE INGREDIENTS:

Heptachlor*	25.00%
Related Compounds	9.72%
INERT INGREDIENTS:	65.28%
Total	100.00%

*Heptachloro-4,5-methanotetrahydroindene

WARNING:

KEEP OUT OF THE REACH OF CHILDREN.

For Antidote and Other Warnings See Back Panels.

COLORADO CHEMICAL CORP., DENVER, COLORADO 80202

DIRECTIONS FOR USE
SEED TREATMENT

No-Stik Heptachlor 25% Seed Treater is applied by drill or planter box application made through dust treaters.

To insure best mixing when applied by planter box proceed as follows:


Place half of the seed in the drill or planter box and half of the total amount of No-Stik Heptachlor 25% Seed Treater uniformly over the seed. Stir the seed and No-Stik by stirring with a paddle until the seed are uniformly covered. Add the No-Stik and again mix thoroughly.

DOSAGE

Crop	Insect
Wheat	False Wireworm
Wheat	Wireworm
Oats	False wireworm
Oats	Wireworm
Barley	False wireworm
Barley	Wireworm
Sorghum	False wireworm
Sorghum	Kafir ant, wireworm
Corn	Wireworm, Southern corn rootworm, Seed corn maggot
Soybeans	Seed corn maggot

Apply Full

— DO NOT RE-USE TRAY
— DESTROY WHEN EMPTY



NO-STIK HEPTACHLOR 25% SEED TREATER

With Graphite Added

For Use in the Control of False Wireworms, Wireworms, Corn Rootworms, Seed Corn Maggots, and Certain Other Soil Insects as Named Below

ACTIVE INGREDIENTS:

Heptachlor* 25.00%
Related Compounds 9.72%

INERT INGREDIENTS: 65.28%
Total 100.00%

*Heptachloro-4,5-methanotetrahydroindene

WARNING:

KEEP OUT OF THE REACH OF CHILDREN.

For Antidote and Other Warnings See Back Panels.

COLORADO CHEMICAL CORP., COLO. 80022

DIRECTIONS FOR USE SEED TREATMENT

No-Stik Heptachlor 25% Seed Treater is designed for drill or planter box application; application can also be made through dust treaters.

To insure best mixing when applying through the drill or planter box proceed as follows:

Place half of the seed in the drill or planter box. Sprinkle half of the total amount of No-Stik Heptachlor 25% Seed Treater uniformly over the seed surface. Mix seeds and No-Stik by stirring with a paddle or stick until the seeds are uniformly covered. Add the remainder of the seed and No-Stik and again mix thoroughly.

DOSAGES

Crop	Insect	Dosage	
		Oz./100 lbs.	Oz./Bu.
Wheat	False Wireworm	8-12	4 3/4-7 1/4
Wheat	Wireworm	4-6	2 1/2-3 1/2
Oats	False wireworm	8-12	4-6
Oats	Wireworm	4-6	1 1/2-2
Barley	False wireworm	8-12	4-6
Barley	Wireworm	4-6	2-3
Sorghum	False wireworm	8	4
Sorghum	Kafir ant, wireworm	4-6	2-3
Corn	Wireworm, Southern corn rootworm Seed corn maggot	2-3	
Soybeans	Seed corn maggot	2-3	

Apply Full Dosage

— DO NOT RE-USE THIS CONTAINER —
DESTROY WHEN EMPTY

ACCEPTED
MAY 02
UNDER THE FEDERAL
FUNGICIDE AND ROU-
TINE FOR ECONOMIC
ED UNDER NO. 111
TO ATTACHED LABEL