


**WARNING:**

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin! In case of contact, flush with plenty of water; for eyes, get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces.

Do not use treated seed for food or feed purposes.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.



**POLYRAM<sup>®</sup>**  
**POTATO SEED TREATER**  
**No. 7**

**WARNING:**

**KEEP OUT OF REACH OF CHILDREN**

See Left Panel for other cautions

**ACTIVE INGREDIENTS:**

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis (dithiocarbamate) - zinc with 1 part by weight (16.1%) ethylenebis - [dithiocarbamic acid] bimolecular and trimolecular cyclic anhydrosulfides and disulfides. . . . . 7.0%

**INERT INGREDIENTS** . . . . . 93.0%  
Total . . . . . 100.0%

Zinc expressed as metallic - 1.23%  
U. S. Patent Pending  
Polyram is a registered Trademark of Badische Anilin & Soda Fabrik A. G.

NET WEIGHT 50 POUNDS

**COLORADO INDUSTRIAL CORP., COMMERCE CITY, COLO. 80012**

**DIRECTIONS FOR USE**

**POTATO-CUT SEED:** Fusarium Seed Piece Decay, Black Leg, Seed-borne Common Scab--Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed pieces. For effective control, Seed Treater must be applied to entire surface of each piece immediately after cutting. If planting is delayed, follow the recommended storage practices of cut seed pieces in your area.

**POTATO-WHOLE SEED:** Seed-borne Common Scab--Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed. Seed Treater should be applied evenly to entire surface. If the treated whole seed is cut, a second application will be necessary if Black Leg and Fusarium Seed Piece Decay are to be controlled.

EPA REG. NO. 1715-282