DIRECTIONS FOR USE

SEED TREATMENT – A Planter Box Treatment

For temporary protection of germinating seed and seedlings from seed corn maggots, seed corn beetles, cutworms, Asiatic garden beetle larvae, European chafer larvae, green June beetle larvae, Japanese beetle larvae, white grubs, wireworms, flea beetle larvae, and for the control of seed and soil-borne diseases, including damping-off and seed decay organisms, and seedling blights. Prepared for use directly in the planter box, or may be mixed in a rotating container such as a cement mixer. Best protection will be obtained where every seed is coated with the chemical. DO NOT USE TREAT-ED SEED FOR FOOD, FEED OR OIL PURPOSE.


If the temperature is high and soil moisture is low, reduce dosage by half, especially for mid-season planting.

Apply dust out of doors only and plant seed promptly after treating.

*NOTE: Captan-Heptachlor Seed Protectant has been mixed with graphite as an aid in safe handling and lubrication of equipment.

TREATMENT OF SMALLLOTS OF SEED (Plant - er Box Rates)

Corn, sorghum, beans, soybeans 4 oz. per Bushel Wheat, barley and oats 6 oz. per cwt.



CAPTAN-HEPTACHLOR

SEED PROTECTANT

| Planter Box Application with Graphite* Added FOR SEED DISEASE AND SOIL INSECT CONTROL WARNING: |
|--|
| KEEP OUT OF THE REACH OF CHILDREN |
| See Side Panel for Additional Warnings |
| ACTIVE INGREDIENTS: |
| Heptachlor (heptachlorotetrahydro-4,7-methanoindene) |
| · · · · · · · · · · · · · · · · · · · |
| Related Compounds |
| Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicar - |
| boximide) |
| INERT INGREDIENTS: |
| Total 100.00% |
| Protect From Excessive Heat - Store In A Cool, Dry Place. |
| Net WeightLbs. |


WARNING: May be fatal if swallowed. Do not breathe vapor or dust. Wear a respirator. Do not get on skin. In case of contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs.

This product is toxic to fish, birds and other wildlife. Treated seeds exposed on soil surface will be hazardous to birds and other wildlife. Dispose of excess treated seeds by burial. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Container should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 4715-290



1