



INDOOR-OUTDOOR INSECT SPRAY

Cameraman's Note
Poor Copy

A. ...

CILLS: Houseflies, Mosquitoes, Gnats, Wasps, Roaches, Earwigs, Two-Spotted Mites, Sowbugs, Spiders, Centipedes, Ants, Biting Flies, Aphids (Plant Lice), Leafhoppers, Exposed Thrips, Adult Whiteflies, Plant Bugs, Japanese Beetles, Spittlebugs.

| ACTIVE INGREDIENTS: | 4 | By Weight |
|---|---|---------------------|
| (5-Benzyl-3-furyl) methyl 2, 2-dimethyl-3- | | |
| (2-methylprope "/i) cyclopro | panecarboxylate | *+ 0 200% |
| Related Compounds | *** ********** | 0.028% |
| d-trans Allethrin (ally) homolog of Cinerin I) 0.150% | | |
| Related Compounds | · · · · · · · · · · · · · · · · · · · | 0.0125 |
| Aromatic Petroleum Hydrocarbo | ons | 0.272 % |
| INERT INGREDIENTS: | • | , . 99 .338' |
| · | Fotal | 100.000% |

Sis/trans isomers ratio: max. 30% (土) cis and miri. 70% (土) trans. Patent No. 3,465,007-SBP-1382円 S.5. Penick & Co., New York

SHAKE BEFORE USING

CAUTION: KEEP OUT OF REACH OF CHILDREN

See back punel for complete caution statement,

WARNING: Contents under pressure. Do not poncture Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

NET WEIGHT: [] [] [] POUNDS

COLORADO INTERNATIONAL CORP.
Commerce City, Colo. 80022

DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

TO CONTROL HOUSEHOLD PESTS

For Houseflies, Mosquitoes, and Grats: Close all doors and windows. Point nozzle upward; direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

For Crawling Insects: Foaches, Earwigs, Grain Mites, Sowbugs, Spiders and Centipedes: Spray thoroughly into Hiding places, cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible.

Ants: Spray traits, nests and points country. Spray on ants where possible. Repeat as necessary.

OUTDOORS—TO REDUCE ANNOYANCE FROM FLYING INSECTS:
Reduce Mosquitoes, Gnats (Midges), Houseflies, Biting Flies and Pyralid and Noctuid Flying Moths by spraying their resting areas. Follow precautions listed above. Hold container 18-24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow, sweeping motion at the rate of from 1-3 seconds per cubic yard of space while moving away from the treated areas. Freat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

OUTDOORS—TO PROTECT ORNAMENTAL PLANTS
To control Aphids, Eynosad Thrips, Adult Whiteflies, Leafhoopers,
Two-spotted Mites, Piantaus, Spittledgs and Japanese Beetles on smaller
Rhododandrons, Pines, Sprice, Hemlock, Spirea, Dogwood, Palm, Mimosa,
Toyan, Viburnum, Euonymus, Oaks, Maples, Locust, Forsythia and Althea:
and Mimosa Webworm Larvae on Mimosa: Spray directly on insects. Hold
container 18 inches from the infested plants. Spray in short spray bursts of
1-3 seconds or spray at the rate of one linear foot of growth per second.
Direct spray to all sides, covering foliage until slightly moist. Avoid heavy
wetting. Repeat as necessary to obtain control.

OUTDOORS -- TO PROTECT FLOWERS

Spray Peionies, Iris, Zinnia, Snapuragons, Ageratums, Chrysanthemums, Nasturtiums, Daisy, Geraniums, Marigolds, Daphne, Fosshedera, Ivv. Tobira, Queen Sago and Broad Leaf Acacia for control of: Aphids, Exposed Thrips, Adult White Flies, Leafhoppers, Plant Bugs. Spittlebugs and Japanese Beetles. Spray as directed above.

USE WITH CAUTION ON: Azaleas: Junipers; Roses, Including Hybrid Tea Roses; Flowering Quince; Ash and Camellias. These plants may be subject to leaf or flower burn under certain conditions of use.

For Adult White Flies, Two-Spotted Mites and Mealybugh: Be suite to direct spray to contact these pests. Thoroughly cover the infested area. Use several repeat applications at 2-day intervals, if necessary.

TO PROTECT HOUSE PLANTS.

Spray infested African Violets, Philodendron, Geraniums, Begonias, and Coleus for the pests listed. Leave the sprayed area when cone indoors; and, ventilate area after treatment is completed.

MUISANCE FESTS — ON or ENTERING BUILDINGS: WASPS: Applications should be made in laterevening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary, Spray directly on stray wasps that enter buildings.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible, Repeat as necessary.

SPIDERS: Direct short burst of spray on spiders and repeat as necessary. EARWICS: Spray directly on Facelings in crevices and other hiding places on contdoor porches, patio foor and doorway entrances.

CAUTION: Harmful If swallowed. Avoid contamination of feed and foodstuffs. Do not use on edible crops. Remove pets, birds and cover fish aquariums before spraying

NOTICE: BUYER ASSUMES ALL RISK OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS OF VENTUREWITH.

EPA Rug No. 471 294

EPA ESI.