



#### GARDEN VEGETABLES AND BERRIES

To control insects as shown with each of the following crops: Use 2½ tsp. 50% MALATHION SPRAY per gallon of water. Spray on foliage every 10-14 days throughout the growing season or as needed. Observe time limitations set before harvest as shown by the number of days in ( ) after each listed crop.

CABBAGE (7); CAULIFLOWER (7); BROCCOLI (3); BRUSSELS SPROUTS (7); KALE (7); KOHLRABI (7); TURNIPS (3); PARSNIPS (7); RUTABAGAS (3): To control: **Aphids, Imported Cabbageworms.**

ASPAPAGUS (1): To control **Asparagus Beetle and Thrips.**

TOMATOES (1); EGGPLANT (3): To control **Spider Mites and Aphids.**

BLACKBERRIES (1); BLUEBERRIES (1); BOYSENBERRIES (1); LOGANBERRIES (1); RASPBERRIES (1); GOOSEBERRIES (3): To control: **Aphids, Spider Mites, Japanese Beetles, Leafhoppers.**

STRAWBERRIES (3): To control **Aphids.**

BEANS (1): To control **Mexican Bean Beetles and Aphids.**  
CUCUMBERS (7); MELONS (1); SQUASH (1); PUMPKINS (3): For control of **Cucumber Beetle, Aphids and Spider Mites.**

POTATOES (3); SPINACH (7); SWISS CHARD (7); COLLARDS (7); PEAS (3); PEPPERS (3); OKRA (1): To control **Aphids.**

GREEN ONIONS (3): To control **Onion Thrips, Onion Maggots.**

CARROTS (7): To control **Leafhoppers.**

#### FRUIT TREES

APPLES, PEARS: To control **European Red Mite, Clover Mite, Two-Spotted mite, Woolly Apple Aphids, Forbes Scale, Green Apple Aphid & Rosy Apple Aphid:** Use 2½ tsp. 50% Malathion per gallon of water. Make two or more applications as needed for mites, one or more for aphids at pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest.

## 50% MALATHION SPRAY

For Control of Specified Insects on: **FLOWERS, ORNAMENTALS, SHRUBS, VEGETABLES and FRUIT TREES** as Listed in Label Directions.

### CAUTION:

### KEEP OUT OF REACH OF CHILDREN

See Side Panels For Additional Cautions

#### ACTIVE INGREDIENTS:

Malathion\* ..... 50.00%

Xylene ..... 42.36%

INERT INGREDIENTS: ..... 7.64%

Total ..... 100.00%

\*O, O-dimethyl dithiophosphate of methyl mercaptosuccinate

Net Contents:

**COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022**

# MALATHION MAY

Insects on: FLOWERS,  
VEGETABLES and FRUIT  
Label Directions.

ION:  
CH OF CHILDREN  
Additional Cautions

.....50.00%  
.....42.36%  
..... 7.64%  
Total ..... 100.00%

e of r' yl mercaptosuccinate

, COMMERCE CITY, COLO. 80022

## FLOWERS, ORNAMENTALS, SHRUBS

Add 2½ t. p. of 50% Malathion to each gallon of water used for spraying. Will control Aphids; Spider Mites; Whiteflies; Thrips; Mealy Bugs; Japanese Beetle; Star Four-Lined Plant Bug; Tarnished Plant Bug; Tree Hopper; Potato Leafhopper; Bagworms; Juniper Scale; Oyster Shell Scale and Eucynymus Scale. For **Soft Brown Scale**: Use 3 tsp. per gallon of water. For **Pine Needle Scale**: Use 4½ tsp. per gallon of water. Apply in summer when infestation occurs. Repeat in 10-12 days. For scales, apply when crawlers have settled on foliage. **CAUTION**: Do not use on certain ferns, including Boston Maidenhair and Pteris, as well as, some species of Crassula, because of possible injury to these ornamentals.

**CAUTION**: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist or fumes. Avoid getting in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling. Change all contaminated clothing and wash it with soap and hot water. Keep away from children and domestic animals. If poisoning occurs, GET PROMPT MEDICAL AID.

**NOTE TO PHYSICIAN**: Malathion may cause cholinesterase inhibition. ATROPINE is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

**ENVIRONMENTAL CAUTION**: This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift of spray from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Avoid contamination of feed and foodstuffs. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not re-use empty container; wrap smaller containers in newspaper and place in trash can for disposal; destroy larger containers by perforating or crushing and bury in safe place away from water supplies.

**NOTICE**: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-299 EPA Est. 4715-CO-1