

ACCEPT
 MAR 28 1973
 UNDER THE FEDERAL INSECT
 FUNGICIDE AND ROENTICIDE
 FOR ECONOMIC ACTION REGIS
 ED UNDER THE FIFTEEN
 TO ALL OTHERS.



**CHLORDANE
 40 SPRAY**

CAUTION: Harmful if swallowed. Contact with skin causes toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Avoid breathing dust or spray mist.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

DIRECTIONS FOR USE

Apply the amount of CHLORDANE 40 SPRAY indicated in sufficient water to treat one acre. For cutworm control, apply to the surface of the soil particularly the areas around the base of the plants. For corn rootworm, root maggots, wireworm and white grub control apply as a broadcast treatment to the soil and immediately work it into the top few inches prior to planting.

BEANS, BEETS: Cutworms — Use 4 to 5 pounds per acre. On green or snap beans do not apply after pods start to form. On beets do not apply after seedling stage if tops are to be used for food.

BROCCOLI, CABBAGE, CAULIFLOWER, PEPPERS, TURNIPS: Wireworms — Use 6 pounds per acre. Apply only as a preplanting soil application.

ACTIVE INGREDIENT:

*Technical Chlordane40.0%

INERT INGREDIENTS:60.0%

Total100.0%

*Equivalent to 24% Octachloro-4,7-Methanotetrahydroindane and 16% related compounds

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See Side Panels For Additional Cautions

CORN, POTATOES: White Grubs, Wireworms, Root Maggots — Use 6½ pounds per acre. Apply pre-planting soil application.

CORN: Corn Rootworms — Use 5 pounds per acre only as a pre-planting soil application.

LAWNS, TURF: Ants, European Earwigs — 4 pounds per acre (1 to 1½ ounces per 1000 sq ft). Thoroughly wash down into turf and soil. Beetle grubs, White-fringed Beetle larvae, C. Beetle Larvae, White Grubs, Wireworms — Use 4 pounds per acre (2½ to 5 ounces per 1000 sq ft).

For 2 to 3 years control use 25 pounds per acre (5 ounces per 1000 square feet) in sufficient water to work insecticide 4 to 6 inches into soil with water. In seeded areas, cultivate insecticide into soil to 4 to 6 inches immediately after applying.

Keep children and domestic pets off lawns and lawns it has been thoroughly watered and completely dry.

ORNAMENTALS, NON-CROP AREAS: Grasshoppers — 2½ pounds per acre. Apply when insects appear and as required.

NOTICE: Seller warrants that this product contains the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other products and conditions of application equipment will vary, seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 4715-311 Net Weight _____

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

ACCEPTED
 MAR 28 1973
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND ROENTICIDE ACT
 FOR ECONOMIC ACTION REGISTER-
 ED UNDER THE FIFTEENTH SUBJECT
 TO APPLICABLE LAWS.



**CHLORDANE
 40 SPRAY**

ACTIVE INGREDIENT:
 *Technical Chlordane40.0%

INERT INGREDIENTS:60.0%

Total100.0%

*Equivalent to 24% Octachloro-4,7-Methanotetrahydroindane and 16% related compounds

CAUTION:

KEEP OUT OF REACH OF CHILDREN
 See Side Panels For Additional Cautions

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022

If swallowed. Contact with skin
 In case of contact with skin, wash
 Avoid contamination of feed and
 thing dust or spray mist
 fish and wildlife. Keep out of any
 feeding on treated areas may be
 act only as specified on this label.
 rying in non-crop lands away from
 iner should be disposed of by
 by burning. Keep out of smoke.
 rial to drift onto neighboring crops
 ise in a manner or at a time other
 th label directions because animal,
 illegal residues or other undesirable
CTIONS FOR USE
CHLORDANE 40 SPRAY indicated
 eat one acre. For cutworm control,
 of the soil particularly the areas
 e plants. For corn rootworm, root
 nd white grub control apply as a
 o the soil and immediately work it
 is prior to planting.
 pods — Use 4 to 5 pounds per acre.
 ns do not apply after pods start to
 it apply after seedling stage if tops
 d.
GE, CAULIFLOWER, PEPPERS,
 — Use 6 pounds per acre. Apply
 soil application.

CORN, POTATOES: White Grubs, Wireworms. **RADISH:**
 Root Maggots — Use 6½ pounds per acre. Apply only as a
 pre-planting soil application.

CORN: Corn Rootworms — Use 5 pounds per acre. Apply
 only as a pre-planting soil application.

LAWNS, TURF: Ants, European Earwigs — Use 2½ to
 4 pounds per acre (1 to 1½ ounces per 1000 square feet).
 Thoroughly wash down into turf and soil. Japanese
 Beetle grubs, White-fringed Beetle larvae, Green June
 Beetle Larvae, White Grubs, Wireworms — Use 6 to 12
 pounds per acre (2½ to 5 ounces per 1000 square feet).

For 2 to 3 years control use 25 pounds per acre (9½
 ounces per 1000 square feet) in sufficient water. Wash
 insecticide 4 to 6 inches into soil with water. For non-
 seeded areas, cultivate insecticide into soil to a depth of
 4 to 6 inches immediately after applying.

Keep children and domestic pets off lawns and turf until
 it has been thoroughly watered and completely dry.

ORNAMENTALS, NON-CROP AREAS: Grasshoppers —
 2½ pounds per acre. Apply when insects appear and repeat
 as required.

NOTICE: Seller warrants that this product conforms to
 the ingredient statement on the label. Since conditions of
 use, such as weather, compatibility with other chemicals,
 and conditions of application equipment will vary, Seller
 makes no claims other than those stated on this label.

E.P.A. Reg. No. 4715-311 Net Weight _____ Lbs.