

**CAUTION:** Inhalation is dangerous, may be absorbed through the skin and may cause eye damage. Do not breathe dust. Always wash thoroughly with soap and warm water after handling. Never eat tampons coated with soap and hot water before re-use. Do not contaminate food or foodstuffs. If non-fogging glasses are used, pay attention immediately. Prolonged exposure will result in cholinesterase depression. **NOTE TO PHYSICIAN:** Atropine Sulfate - antidote. 2 PAM and salbutamol and may be used in conjunction with Atropine.

**ENVIRONMENTAL CAUTION:**

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not spray when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply the product only as specified in this label.

This product is toxic to birds and should not be applied when persons are actively visiting the area.


Do not use empty container. Destroy empty container by puncturing, crushing, or burying in safe place.

**DIRECTIONS**

**NAVUNEX** is a selective insecticide which controls a wide variety of insect pests but does not destroy beneficial insects. It provides protection to your crops against many pests.

For best results, mix NAVUNEX with kaolin (talc) or lime and then sulfur or with aluminum and ferric oxide. Mix until the mixture is uniform. Apply at approximately one pound per acre. However, the amount of NAVUNEX Dust to use is less than one pound per acre. This is effective.

Begin applications when insects first appear and repeat as required. Dust plants thoroughly. Observe day intervals between dust application and harvest indicated by number of days following a crop.



## NAVUNEX 5 DUST

**Contains 5% Trichlorfon**

A Selective Insecticide  
for Use on Certain Vegetable Crops

ACTIVE INGREDIENT:

Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by leaching of equipment or disposal of waste. Apply the product at the rate specified on this label.

This product is not to be used around flowers or shrubs which bees frequent very visibly on the ground.

Do not use empty containers. Do not empty or contain by putting in the crushing or burying in a safe place.

#### DIRECTIONS

NAVUNEX is a selective insecticide which controls a wide variety of insect pests that does not destroy the field crop. It has the following uses: (1) Control of Japanese Beetle on lawns.


Do not use NAVUNEX 5 Dust in combination with alkaline materials such as lime and lime sulfur or with summer oil and formalin. Is NAVUNEX 5 known to be non-patible with all nitrated and phosphated insecticides. However, the effect of a mixture of NAVUNEX 5 Dust is lost when it is applied in combination with other insecticides which are also selective.

Begin treatments when seeds are up and repeat as required. Dust can be thoroughly applied by day. Intervene between dust applications and harvest indicated by number of days to a crop.

**ARTICHOKE** (14) - 3 to 5 lbs. per acre. Apply to 10 to 50 lbs. per acre.

**ALFALFA, CLOVER (7)** - Alfalfa Cutworm - Alfalfa Webworm - Leafworm - Cowpea Bugs - Leaf Bugs - Grasshopper - Bean Maggots - Cutworms. Weight - 50 lbs. per acre. (14) - 20 lbs. per acre. Only one application per cutting. No cutting or grazing until 7 days after treatment.

**BEANS (DRY & SNAP), COWPEAS, SOUTHERN PEAS, BLACK-EYED PEAS, CROWDER PEAS (14)** - Armyworm - Cotton Bollworm - Imported Cabbageworm - Imported Mexican Bean Borer - Striped Borer - Imported Cutworm - Western Bean Cutworm - 20 to 30 lbs. per acre. Do not feed treated vines to meat or dairy animals. Do not apply more than twice per season for cowpeas.



# NAVUNEX 5 DUST

## Contains 5% Trichlorfon


#### A Selective Insecticide

for Use on Certain Vegetable Crops

#### ACTIVE INGREDIENT

Trichlorfon (Dimethyl (2,2-dichloro-  
hydroxyethyl) phosphonate)

#### INERT INGREDIENTS




#### CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR REQUIRED CAUTION STATEMENT.

NET WEIGHT

POUNDS



RECEIVED  
MARCH 13 1975  
for the  
and  
agriculture  
CPA

## NAVUNEX 5 DUST

Contains 5% Trichlorfon

A Selective Insecticide  
for Use on Certain Vegetable Crops

ACTIVE INGREDIENT

Trichlorfon (Methyl 2-Chloro-

BEEFTEAK 20 Btu/lb  
100 mg/L (ppm) 100 mg/L (ppm)  
100 mg/L (ppm) 100 mg/L (ppm)

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER  
(21) 100 mg/L (ppm) 100 mg/L (ppm)  
100 mg/L (ppm)

COTTON 75 mg/L (ppm) 75 mg/L (ppm)  
100 mg/L (ppm) 100 mg/L (ppm)  
Black  
Flameless Compaction Cotton 100 mg/L (ppm)  
Cotton 100 mg/L (ppm) 100 mg/L (ppm)

LEAVES 100 mg/L (ppm)  
Eucalyptus, Pepper, Tomato, A. C. C. and  
Vinegar 100 mg/L (ppm) 100 mg/L (ppm)

PEPPERS 21 100 mg/L (ppm) 100 mg/L (ppm)

PUMPKINS 140 100 mg/L (ppm) 100 mg/L (ppm)

SAFFLOWER 100 mg/L (ppm) 100 mg/L (ppm)  
100 mg/L (ppm) 100 mg/L (ppm)  
100 mg/L (ppm) 100 mg/L (ppm)

SUGAR BEETS 140 100 mg/L (ppm) 100 mg/L (ppm)  
Beets 100 mg/L (ppm) 100 mg/L (ppm)  
100 mg/L (ppm) 100 mg/L (ppm)  
100 mg/L (ppm) 100 mg/L (ppm)

TOBACCO 30 100 mg/L (ppm) 100 mg/L (ppm)  
20 100 mg/L (ppm) 100 mg/L (ppm)

TONS TREATED 100 mg/L (ppm) 100 mg/L (ppm)

