



DANGER—POISON—PRECAUTIONS 

POISONOUS IF SWALLOWED

Small amount of this product can be fatal if swallowed. Do not use if you are ill or have a fever.



POISONOUS BY SKIN CONTACT

Do not use if you have a skin condition. Do not use if you are pregnant or nursing. Wash hands thoroughly after use.



POISONOUS IF BREATHED

Do not use in enclosed spaces. Do not use near heat or open flame. Do not use near children or pets.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. COMBUSTIBLE MIXTURE.

**MEVINPHOS EM 4**  
ORGANOPHOSPHORUS INSECTICIDE EMULSIFIABLE LIQ  
Contains 4 Lbs. Mevinphos Per Gallon

**STOP READ LABEL BEFORE USE**  
**NOT FOR HOME USE**

ACTIVE INGREDIENTS:

Alpha Isomer of 2-Carbomethoxy-1-Methylvinyl

Dimethyl Phosphate

Related Compounds

Aromatic Petroleum Solvent Derivatives

INERT INGREDIENTS:

 **POISON**  Total ...

**DANGER KEEP OUT OF REACH OF CHILDREN**

See side panel for Antidote and other required Warning statements.  
Read all directions carefully before using.

NET CONTENTS: 4.0 LBS. (1.818) GALLONS

COLORADO INTERNATIONAL CORPORATION  
COMMERCE CITY, COLORADO 80022

ACCEPTED

# MEVINPHOS EM 4

DIPHOSPHORUS INSECTICIDE EMULSIFIABLE LIQUID

Contains 4 Lbs. Mevinphos Per Gallon

**READ LABEL BEFORE USING!  
NOT FOR HOME USE**

INGREDIENTS:	By Weight
2-Carbomethoxy-1-Methylvinyl phosphate .....	27.6%
Surfactants .....	18.4%
Hydrocarbon Solvent Derivatives .....	47.0%
EMULSIFIERS: .....	7.0%
Total .....	100.0%



**KEEP OUT OF REACH OF CHILDREN**

See separate panel for Antidote and other required Warning statements.  
Read all directions carefully before using.

NET CONTENTS:     GALLONS  
1 5 10 55

COLORADO INTERNATIONAL CORPORATION  
COMMERCE CITY, COLORADO 80022

**DANGER:** Poisonous if Swallowed. Toxic if Inhaled. Absorbed Through Skin. Repeated Inhalation Through Nose. Repeat Inhalation of vapors may, without symptoms, progressively increase susceptibility to mevinphos insecticide poisoning. Do not swallow or get in eyes, on skin or on clothing. Do not breathe vapors or spray mist. Wear clean natural rubber gloves and clean waterproof protective clothing and goggles. Replace gloves frequently and discard used gloves. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Phosphorus Insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing in soap and hot water before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with strong eye solution.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected humans and animals should be kept out of treated fields for 24 hours after treatment. Do not contaminate food or feed products. Keep away from heat and open flames. Rinse equipment and containers and dispose of empty wastes and dry sprays by spraying into crops/lands away from water supplies.

**ANTIDOTE AND FIRST AID TREATMENT**  
**ATROPINE IS THE EMERGENCY ANTIDOTE FOR MEVINPHOS INSECTICIDE POISONING.** Consult your physician about obtaining an immediate supply of 1-100 grain atropine tablets. For emergency use, use **WALL-PHOS** Antidote.

**IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

**INTERNAL:** If the material has been swallowed, induce vomiting immediately. This may be done by irritating the back of the throat or pouring 150 ml (5 fl. oz.) of 1-tablespoonful of 1% glass of water. **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

**EXTERNAL:** If the material has been spilled on the skin, immediately remove patient from vicinity of the insecticide; remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

**IF WARNING SYMPTOMS APPEAR:** Administer two 1-100 grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See **WARNING SYMPTOMS** below). Keep patient propped up and breathe. Start artificial respiration immediately if patient is not breathing. Transport patient immediately to the nearest physician.

### NOTE TO PHYSICIAN

**WARNING SYMPTOMS:** Weakness, headache, tightness in chest, blurred vision, dry mouth, dilated pupils, skin flush, sweating, nausea, vomiting, diarrhea and abdominal cramps.

**TREATMENT:** Atropine is the specific antidote. Antidote of choice against mevinphos parasympathetic stimulation. If there is a lot of parasympathetic stimulation, atropine should be injected at 10 minute intervals doses of 1-2 milligrams, until complete parasympathetic stimulation has occurred. 2-PAM is a specific antidote for 2-PAM. 2-PAM is a specific antidote for 2-PAM. 2-PAM is a specific antidote for 2-PAM.

**DANGER.** Poisonous if Swallowed. Irritant if Absorbed. Irritant if Skin Contact. Irritant if Inhaled. Repeat Inhalation or Irritant if Contact may, without symptoms, progressively increase susceptibility to mevinphos insecticide poisoning. Do not swallow or get on eyes, or on or on clothing. Do not breathe vapor or spray mist. Wear clean, natural rubber gloves and clean water-proof protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of type passed by the U.S. Bureau of Mines for Phosphorus Insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing in soap and hot water before reuse. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with strong lye solution.

Do not apply or allow to drift to area occupied by unprotected humans or beneficial animals. Unprotected humans and animals should be kept out of treated fields for 24 hours after treatment. Do not contaminate food or feed products, keep away from heat and open flame. Rinse equipment and containers and dispose of any wastes and any sprays by spraying on non-crop lands away from water supplies.

**ANTIDOTE AND FIRST AID TREATMENT**  
**ATROPINE IS THE EMERGENCY ANTIDOTE FOR MEVINPHOS INSECTICIDE POISONING.** Consult your physician about obtaining an immediate supply of 1-100 grain tablets. See also instructions by label. **TEL. PH 760, A. 115**

**IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

**INTERNAL:** If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

**EXTERNAL:** If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide; remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

**IF WARNING SYMPTOMS APPEAR:** Administer two 1-100 grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See **WARNING SYMPTOMS** below). Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

**NOTE TO PHYSICIAN**

**WARNING SYMPTOMS:** Weakness, headache, tightness in chest, blurred vision, nausea, pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

**TREATMENT:** Atropine is the specific therapeutic antagonist of choice against the sympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals in doses of 1-2 milligrams, until complete atropinization has occurred. 2-4 AM is also antidotal and may be administered in combination with atropine. 1-2 mg. atropine should be given every 10

minutes by postural or i.v. route. Oxygen administration may be necessary. Observe patient continuously for 48 hours. A cholinesterase inhibitor may, without warning, cause profound respiratory depression. Small doses of any cholinesterase inhibitor will allow no further exposure until time for cholinesterase regeneration has been allowed, as determined by blood test.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**


**ENVIRONMENTAL CAUTION:** This product is toxic to fish, shrimp, crabs and other wildlife. Keep out of lakes, streams, ponds, swamps and estuaries. Do not apply where runoff is likely to occur. Do not apply where other conditions favor drift to crops. Do not apply. Shrimp and crabs may be killed if applied on rates recommended on this label. Do not apply where these are important resources. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. For more information, may be obtained from your Cooperative Agricultural Extension Service.

**CONTAINER DISPOSAL:** Do not reuse empty container. Destroy by crushing and burying or incinerating away from water supplies.

**PRECAUTIONS:** See label carefully that this product conforms to the ingredients listed on the label. Since conditions of use, use of other chemicals, compatibility with other chemicals and conditions of application in equipment vary, always wear masks and other protection when using this label.

REG. PAT. NO. 17-1194 EPA REG. NO. 133-1011



**ING!**

Weight

- 27.6%
- 18.4%
- 47.0%
- 7.0%

---

- 100.0%

**DREN**

21-55

# MEVINPHOS EM 4

## READ ALL PRECAUTIONS AND DIRECTIONS BEFORE USING

### GENERAL INFORMATION

Application can be made by aircraft or ground power equipment by trained personnel only using approved protective equipment. **Do not apply with hand equipment.**

Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

### AIRCRAFT AND GROUND APPLICATION

The actual rate required to provide thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are therefore intended to cover a broad range of conditions.

Crop	Gal. Water/Acre	
	Aircraft	Ground
Vegetable and Field Crops	5-20	20-125
Grasses	7-25	20-200
Orchard Crops	10-20	40-800

### DIRECTIONS FOR USE

OBSERVE DAYS INTERVAL BETWEEN LAST APPLICATION AND HARVEST AS INDICATED BY THE NUMBER IN FOLLOWING THE CROP.

#### FRUIT AND NUT CROPS

##### APPLES (1), PEACHES (1), PEARS (1), PLUMS (1)

**Aphids, Mites:** Use 1/2 to 1 pt. 100 gal. water. **Grasshoppers, Lygus Bugs and Red-banded Leafroller:** Use 1/2 to 2/3 pt. 100 gal. water.

**NOTE:** Do not apply more than 5 pts. / acre on peaches and plums, or 6 1/2 pts. / acre on apples and pears. **PEACHES ONLY:** Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the re-entry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

##### CHERRIES (SOUR ONLY) (2)

**Aphids, Mites:** Use 1/2 to 2/3 pt. 100 gal. water.

**NOTE:** Do not apply more than 5 pts. / acre.

##### CITRUS (ORANGES (1), LEMONS (1), GRAPEFRUIT (1))

**Aphids:** Use 1/2 to 2 pts. / acre in 200 or more gal. water. **Fruit Tree Leafroller, Orange Tortrix, Omnivorous Leafroller:** Use 1 pt. / acre in 500 gal. water. **Western Tussock Moth Larvae, Citrus Cutworm, Variegated Cutworm, Pink Scavenger Caterpillar:** Use 2 pt. / acre in 1200 gal. water. Allow 7 days / acre between applications.

**NOTE:** Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the re-entry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

#### GRAPES

**Aphids:** Use 1/2 to 1 pt. / acre. Do not treat within 2 days of harvest. **Leaf Folder, Leafhoppers, Mites, Red banded Leafroller and Lygus Bugs:** Use 1 to 2 pt. / acre. Do not treat within 2 days of harvest. Do not re-enter within 5 days of harvest.

Grapes  
Orchard Crops

7-25  
10-20

20-200  
40-800

### DIRECTIONS FOR USE

OBSERVE DAYS INTERVAL BETWEEN LAST APPLICATION AND HARVEST AS INDICATED BY THE NUMBER FOLLOWING THE CROP

#### FRUIT AND NUT CROPS

##### APPLES (1), PEACHES (1), PEARS (1), PLUMS (1)

Aphids, Mites: Use 1 to 2 pt. 100 gal. water to Grasshoppers, Lygus Bugs and Red-banded Leafroller: Use 1 to 2.3 pt. 100 gal. water

NOTE: Do not apply more than 50% of one or more of the following pesticides to any one acre. **PEACHES ONLY:** Do not permit bees to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the reentry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

##### CHERRIES (SOUR ONLY) (2)

Aphids, Mites: Use 1 to 2.3 pt. 100 gal. water \*

NOTE: Do not apply more than 5 pt. / acre

##### CITRUS (ORANGES (1), LEMONS (1), GRAPEFRUIT (1))

Aphids: Use 1 to 2 pt. / acre in 200 to 400 gal. water. Fruit Tree Leafroller, Orange Tortrix, Omnivorous Leafroller: Use 1 to 2 pt. / acre in 600 gal. water. Western Tussock Moth Larvae, Citrus Cutworm, Variegated Cutworm, Pink Scavenger Caterpillar: Use 1 to 2 pt. / acre in 1200 gal. water. Allow at least 7 days between applications.

NOTE: Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the reentry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

#### GRAPES

Aphids: Use 1 to 2 pt. / acre. Do not treat within 2 days of harvest. Leaf Folder, Leafhoppers, Mites, Red-banded Leafroller and Lygus Bugs: Use 1 to 2 pt. / acre within 2 pt. / acre do not treat within 5 days of harvest.

NOTE: Do not permit workers to be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the reentry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

##### STRAWBERRIES (1)

Aphids, Mites: Use 1 to 2 pt. 100 gal. water. Grasshoppers, Strawberry Leafroller, Salt marsh Caterpillar and Lygus Bugs: Use 1 to 2 pt. 100 gal. water. Earliest of Aphids and Mites: Use 1 to 2 pt. 100 gal. water

NOTE: Do not apply more than 5 pt. / acre

##### WALNUTS (1)

Aphids: Use 1 to 2 pt. / acre in 100 gal. water. Mites, Omnivorous Looper, Orange Tortrix, Fruit Tree Leafroller, and Western Tussock Moth: Use 1 to 2 pt. / acre in 200 gal. water

NOTE: Do not apply more than 5 pt. / acre

#### FIELD CROPS

##### ALFALFA (1), CLOVER (1)

Aphids, Alfalfa Caterpillar: Use 1 to 2 pt. / acre. Grasshoppers, Leafhopper, Cutworms (cumbing), Mites, Alfalfa Weevil Larvae, Lygus Bugs: Use 1 to 2 pt. / acre

NOTE: Do not exceed 5 pt. / acre

##### MUSTARD GREENS (3), TURNIP TOPS (3)

Aphids: Use 1 to 2 pt. / acre. Cabbage Lopper, Imported Cabbage Worm, False Chinch Bug, Diterous Leafminer (adult), Grasshoppers, Leafhoppers and Mites: Use 1 to 2 pt. / acre

NOTE: Do not exceed 5 pt. / acre

**SQUASHES (FOR VEGETABLES)**

Aphids: Use 1/2 to 1 qt. per acre.

**EGGPLANT (1)**

**ARTICHOKES**

Aphids: Use 1/2 to 1 qt. per acre. Plum-Moth

**BEANS (1)**

Aphids: Use 1/2 to 1 qt. per acre. Grasshopper, Leafhopper, Mites and Salt Marsh Caterpillar

NOTE: Do not exceed 1 qt. per acre.

**BEETS (INCLUDING TOPS) (1)**

Aphids: Use 1/2 to 1 qt. per acre. Cabbage Looper, Cutworm (climbing), Dipterous Leafminer (adult), False Chinch Bug, Grasshoppers, Imported Cabbage Worm, Leafhopper, Mites and Salt Marsh Caterpillar

**BROCCOLI (1), CABBAGE (1)**

Aphids: Use 1/2 to 1 qt. per acre. Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhopper, Salt Marsh Caterpillar, Mites, Cutworms (climbing), Dipterous Leafminer (adult) and Lygus Bugs

NOTE: Do not exceed 1 qt. per acre. For head to leaf Aphids: Use 1 qt. per acre, but do not treat within 5 days of harvest.

**BRUSSELS SPROUTS (3), CAULIFLOWER (3), COLLARDS (7), KALE (7)**

Aphids: Use 1/2 to 1 qt. per acre. Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt-Marsh Caterpillar, Mites, Cutworms (climbing), Dipterous Leafminer (adult) and Lygus Bugs: Use 1/2 to 1 qt. per acre. For head to leaf Aphids: Use 1 qt. per acre, but do not treat collards and kale within 7 days of harvest, 2 days on Brussels sprouts and cauliflower.

**CARROTS (1)**

Aphids: Use 1/2 to 1 qt. per acre. Leafhoppers, Lygus Bugs, Mites, Cabbage Looper, Dipterous Leafminer (adult), Cutworm (climbing) and Salt-Marsh Caterpillar: Use 1/2 to 1 qt. per acre.

**CELERY (3)**

Aphids: Use 1/2 to 1 qt. per acre. Dipterous Leafminer (adult), Lygus Bugs, Salt Marsh Caterpillar, Leafhoppers, Cabbage Loopers and Mites. Use 1/2 to 1 qt. per acre. For head to leaf Aphids: Use 1 qt. per acre, but do not treat within 5 days of harvest.

**CORN (FIELD, SWEET, AND POPCORN) FOR GRAIN ONLY (1)**

Aphids: Use 1/2 to 1 qt. per acre.

NOTE: Do not exceed 1 qt. per acre.

**CUCUMBERS (1), EGGPLANT (1), PEPPERS (1)**

Aphids: Use 1/2 to 1 qt. per acre. Grasshoppers, Leafhoppers, Mites: Use 1/2 to 1 qt. per acre.

NOTE: Do not exceed 1 qt. per acre.

**LETTUCE (10)**

Aphids: Use 1/2 to 1 qt. per acre. Corn Earworm, Cutworms (climbing), Dipterous Leafminer (adult), Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Mites, Lygus Bugs, Salt Marsh Caterpillar, False Chinch Bug and Thrips: Use 1/2 to 1 qt. per acre. For head to leaf Aphids: Use 1 qt. per acre, but do not treat within 5 days of harvest.

**MELONS (INCLUDING CANTALOUPE) (1), HONEY DEW MELONS (1),**

**MUSKMELONS (1), WATERMELONS (1)**

Aphids: Use 1/2 to 1 qt. per acre. Cabbage Looper, Dipterous Leafminer (adult), Leafhoppers, Lygus Bugs, Mites, False Chinch Bug, Salt Marsh Caterpillars, Cutworms (climbing), and Grasshoppers: Use 1/2 to 1 qt. per acre.

Caterpillar, Mites, Cutworms (climbing), Dipterous Leafminer (adult) and Lygus Bugs: Use 1/2 to 1 pt./acre. For  
Cabbage Looper, Imported Cabbag Worm, Grasshoppers, Leafhoppers, Mites, Lygus Bugs, Salt Marsh Caterpillar, but do not treat collards and kale within 7 days of harvest, 3 days on Brussels  
sprouts.

#### CARROTS (1)

Aphids: Use 1/2 to 1 pt./acre. Leafhoppers, Lygus Bugs, Mites, Cabbage Looper, Dipterous Leafminer (adult), Cutworm  
(climbing), Salt Marsh Caterpillar, Grasshoppers: Use 1/2 to 1 pt./acre.

#### CELERY (3)

Aphids: Use 1/2 to 1 pt./acre. Dipterous Leafminer (adult), Lygus Bugs, Salt Marsh Caterpillar, Leafhoppers, Cabbage  
Looper, Imported Cabbag Worm, Grasshoppers, False Chinch Bug, Aphids: Use 1/2 to 1 pt./acre, but do not treat within 5 days of harvest.

#### CORN (FIELD, SWEET, AND POPCORN) FOR GRAIN ONLY (1)

Aphids: Use 1/2 to 1 pt./acre.  
NOTE: Do not exceed 1 pt./acre.

#### CUCUMBERS (1), EGGPLANT (1), PEPPERS (1)

Aphids: Use 1/2 to 1 pt./acre. Grasshoppers, Leafhoppers, Mites: Use 1/2 to 1 pt./acre.  
NOTE: Do not exceed 1 pt./acre.

#### LETTUCE (10)

Aphids: Use 1/2 to 1 pt./acre. Corn Earworm, Cutworms (climbing), Dipterous Leafminer (adult), Cabbage Looper,  
Imported Cabbage Worm, Grasshoppers, Mites, Lygus Bugs, Salt Marsh Caterpillar, False Chinch Bug and Thrips: Use  
1/2 to 1 pt./acre. For hard to kill Aphids: Use 1 pt./acre, but do not treat within 14 days of harvest.

#### MELONS (INCLUDING CANTALOUPE (1), HONEY-DEW MELONS (1), MUSKMELONS (1), WATERMELONS (1))

Aphids: Use 1/2 to 1 pt./acre. Cabbage Looper, Dipterous Leafminer (adult), Leafhoppers, Lygus Bugs, Mites, False  
Chinch Bugs, Salt Marsh Caterpillars, Cutworms (climbing), and Grasshoppers: Use 1/2 to 1 pt./acre.

#### ONIONS (INCLUDING GREEN ONIONS) (1)

Thrips and Cutworms (climbing): Use 1/2 to 1 pt./acre.

#### PEAS (INCLUDING VINES) (1)

Aphids: Use 1/2 to 1 pt./acre. Grasshoppers, Leafhoppers, Mites and Cutworms (climbing): Use 1/2 to 1 pt./acre.

#### POTATOES (1)

Aphids: Use 1/2 to 1 pt./acre. Grasshoppers, Leafhoppers and Mites: Use 1/2 to 1 pt./acre.

#### SPINACH (4)

Aphids: Use 1/2 to 1 pt./acre. Cabbage Looper, Grasshoppers, Leafhoppers, Mites, Dipterous Leafminer (adult), Cutworms  
(climbing), Salt Marsh Caterpillar and False Chinch Bug: Use 1/2 to 1 pt./acre. For hard to kill Aphids: Use 1 pt./acre,  
but do not treat within 7 days of harvest.

#### SUMMER SQUASH (1)

Aphids: Use 1/2 to 1 pt./acre. Cabbage Looper, Dipterous Leafminer (adult), Leafhoppers, Lygus Bugs, Mites, False  
Chinch Bugs, Salt Marsh Caterpillar, Cutworms (climbing) and Grasshoppers: Use 1/2 to 1 pt./acre.

#### TOMATOES (1)

Aphids: Use 1/2 to 1 pt./acre. Grasshoppers, Leafhoppers and Mites: Use 1/2 to 1 pt./acre.

#### TURNIPS (3)

Aphids: Use 1/2 to 1 pt./acre. Cabbage Loopers, Imported Cabbage Worms, Grasshoppers, Leafhoppers, Mites, False  
Chinch Bugs and Dipterous Leafminers (adult): Use 1/2 to 1 pt./acre.