



MALATHION POULTRY, HOG & CATTLE DUST

Contains 5% Malathion

ACTIVE INGREDIENT:	
Malathion: (*O, O-dimethyl phosphorodithioate of diethyl mercaptosuccinate	5%
INERT INGREDIENTS:	95%
Total	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Back Panel for additional required cautions.

NET WEIGHT: 1 3 4 50 POUNDS

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

DIRECTIONS FOR USE

POULTRY

Northern Fowl Mite, Poultry Lice: Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4-8 weeks or when necessary. Treatment of poultry for Northern fowl mite should be a supplemental treatment to the treatment of the premises.

Premise Treatment: Apply 1 lb. per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent area.

HOGS

Lice: Treat animals thoroughly. Repeat in 10 days and thereafter as needed.

DOGS & CATS

Fleas: Direct Application: Apply sufficient dust to treat animal thoroughly. Use any garden-type duster, puff duster or shaker-type can. If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatment. If the animal is confined to an area free of fleas, one treatment may be sufficient.

CATTLE – Beef & Non-Lactating

Lice: Treat animal thoroughly. Repeat in 10 days and thereafter as needed.

Horn Fly: Use 3 Tbsp. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals.

NOTE: Do not treat animals under one month of age. When applying dusts avoid contaminations of feed, food containers, and water troughs.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing of dust or fumes. Avoid getting in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling. Change all contaminated clothing and wash it with soap and hot water. Keep away from children and domestic animals. If poisoning occurs, GET PROMPT MEDICAL AID.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. ATROPINE is antidotal, 2-PAM is also antidotal and may be used in conjunction with atropine.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift of spray from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Avoid contamination of feed and foodstuffs. Apply this product only as specified on this label.

Do not re-use empty container; wrap smaller containers in newspaper and place in trash can for disposal; destroy larger containers by perforating or crushing and bury in safe place away from water supplies.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-3'

EPA Est. 4715-CO-1