CAPTAN 5% DUST

. .

ACCEPTED

JUN 10 1975

FUNCIONE AND NOBLATIONS AND POR ECONOMIC POISON REGISTERS ED UNDER NO. 47/5 - 340

SPECIAL FORMULATION POTATO SEED PIECE TREATMENT

CAUTION:

KEEP OUT OF REACH OF CHILDREN
See back panel for required caution statements.

NET WEIGHT: \$\bigcup_{10} \bigcup_{50} \text{POUNDS}\$

COLORADO INTERNATIONAL CORPORATION Commerce City, Colorado 80022

Cameraman's Note
Poor Copy

DIRECTIONS FOR USE

POTATO SEED PIECE TREATMENT

To control Damp-Off, Seed Rot Fungi, Brown Eye Disease: Apply at the rate of 20 oz. per bushel cut seed pieces or 2 lbs. (32 oz.) to 100 lbs. of cut seed pieces. Apply in convenient container or by duster attachment over belt. Coat each seed evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, stole treated seed pieces in open crates two days before bagging.

NOTE: Do not feed treated potato seed to livestock or use for human food. Dispose unused treated potato seed by burying or plowing under the soil.

CAUTION: Harmful if taken internally. Avoid inhaling dust Avoid contact with skin. Avoid contamination of feed and food. Store in a cool dry place; protect from excessive heat.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label

CONTAINER DISPOSAL: Do not re use empty containers. Destroy by burying with wastes or by burning. Keep away from smoke or fumes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715 340

EFA Est 4715-CO-1