

Cameraman's Note

Poor Copy

FOR USE BY PROFESSIONAL PEST CONTROL OPERATORS ONLY

DANGER: VAPONA insecticide is poisonous if swallowed, inhaled or absorbed through the skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Wear natural rubber gloves, protective clothing and goggles when handling. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP (VAPONA insecticide) protection. Keep all unprotected persons out of operational area. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before smoking or eating. Wash all contaminated clothing with soap and hot water before reuse.

ANTIDOTE AND FIRST AID TREATMENT: ATROPINE is the emergency antidote for VAPONA Insecticide poisoning. Consult your physician about an adequate supply of 1/100-grain atropine tablets for emergency use. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoonful of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See warning symptoms below).

KEEP patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN: WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

PRECAUTIONS IN USING: VAPONA is poisonous; therefore precautions must be observed when handling. (See antidote and warnings.) Store in a safe place. Children or animals must not be able to reach or handle. Care should be used in handling empty containers. Do not contaminate feed, water or foodstuffs.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F. may cause bursting. Never throw container into fire or incinerator.

Manufactured By:
PET CHEMICALS, INC.
P.O. Box 656, Miami Springs, Fla. 33166



DIRECTIONS:
area of 100
forging and
insecticide p
All equipment
building and
under way. R
should be w
tanks or ren
of tender p
Place man
can use wh
free flow of
base of t
the p
and VAPONA
and
VAPONA

OPERATORS ONLY

through the skin or eyes. Do protective clothing and goggles. Bureau of Mines for DDVP (VAPONA). In case of contact, wash immediately with soap and water before smoking or eating. Do not reuse.

Antidote for VAPONA Insecticide poisoning: Tablets for emergency use. CALL A

This may be done by introducing a small amount of water. NEVER GIVE

patient from the vicinity of the material. If the material gets into

eyes immediately NEVER ADMINISTER

if not breathing. Transport the

headache, tightness in chest, blurred vision, nausea, vomiting, and cramps.

antidote: parasympathetic nervous stimulant. Administer at 10-minute intervals. Use is contraindicated. Clear chest

with atropine. Clear chest with atropine. Clear chest with atropine. Clear chest

with atropine. Clear chest with atropine. Clear chest with atropine. Clear chest

with atropine. Clear chest with atropine. Clear chest with atropine. Clear chest

with atropine. Clear chest with atropine. Clear chest with atropine. Clear chest

with atropine. Clear chest with atropine. Clear chest with atropine. Clear chest

3166



NET WEIGHT 12 OZ.



KILLS EXPOSED STAGES OF FOLLOWING INSECTS: ROACHES, FLIES, SILVERFISH, FLYING MOTHS, WEEVILS IN HOMES, BUILDINGS AND RAILROAD CARS

DIRECTIONS: To be used by licensed pest control operators only. Use one 12 oz. can for each unobstructed closed area of 10,000 cubic feet but not less than 7,000 cubic feet. Use one 6 oz. can for each 5,000 cubic feet. During fogging and to release contents use a mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP insecticide protection. Open cabinets and doors to areas to be treated. Shut off air conditioners and fans before using.

All occupants must be removed from the house or building before treatment is started and should not re-occupy building for at least 2 hours. Open doors and windows and allow to be ventilated before re-occupying. Unwrapped or unsealed cans should be removed or placed in refrigerator during fogging. Food handling equipment, dishes, should be washed after application. Do not contaminate water, feed or foodstuffs. Remove all pets, cover fish tanks or remove. Cover house plants within 8 feet of point of release with newspaper to prevent possible burning of tender plants.

Place cans in center of area to be treated, or near point of greatest infestation. Place newspapers under cans. Use one or more cans in combinations of 6 and 12 ounce sizes to give the best distribution of vapors where free flow of mist is not assured. Snap flip over lip of container. Leave building at once. The entire contents will be released. Do not use in edible products areas of food processing plants, restaurants, or other areas where food is usually prepared or processed. Do not use in serving areas while food is being served.

ACTIVE INGREDIENTS: 1,1-dichloro-2,2-dimethyl phosphoramide 6.51%, related compounds 0.49%, Petroleum distillates 85.00%

FOR USE WHERE A LONG LASTING RESIDUAL IS NOT DESIRED



DANGER: KEEP OUT OF REACH OF CHILDREN.

See antidote statement and other cautions on back panel

EPA Reg. No. 47

ACC
12
UNDER THE
FUNGICIDE
FOR B.O.
ED UNDER